

Class: XII Session: 2020-21

Subject: Mathematics

Practice Question Paper 3 (Theory)

Time Allowed: 3 Hours

Maximum Marks: 80

**General Instructions:**

1. This question paper contains two **parts A and B**. Each part is compulsory. Part A carries **24** marks and Part B Carries **56** marks
2. **Part-A** has Objective Type Questions and **Part-B** has Descriptive Type Questions.
3. Both Part A and Part B have choices.

**Part-A:**


1. It consists of two sections- **I and II**
2. Section **I** comprises of 16 very short answer type questions.
3. Section **II** contains **2** case studies. Each case study comprise of 5 case-based MCQs. An examinee is to Attempt **any 4 out of 5 MCQs**.

**Part –B:**

1. It consists of three sections- **III, IV and V**.
2. Section **III** comprises of 10 questions of **2 marks** each.
3. Section **IV** comprises of 7 question of **3 marks** each.
4. Section **V** comprises of 3 questions of **5 marks** each.
5. Internal choice is provided in **3** questions of Section –III, **2** questions of Section- IV and **3** questions of Section-V. You have to attempt only one of the alternatives in all such questions.

| Sr. No. | Part-A  | Marks |
|---------|---|-------|
|         | <b>Section I</b><br><b>All questions are compulsory. In case of internal choices attempt any one.</b>               |       |
| 1.      | Consider $f: R_+ \rightarrow [4, \infty]$ given by $f(x) = x^2 + 4$ . Check whether the function is one-one or not. | 1     |



|   |   |   |
|---|---|---|
| 11  | Find the area of a parallelogram whose two adjacent sides are $\hat{j} + 2\hat{k}$ and $\hat{i} + 2\hat{j}$ .   | 1 |
| 12  | For what value of 'k', the matrix $\begin{pmatrix} 2 & 5 \\ k & 10 \end{pmatrix}$ is a singular matrix?   | 1 |
| 13  | If a plane has the intercepts a, b, c and is a distance of 'p' units from the origin, then $\frac{1}{a^2} + \frac{1}{b^2} + \frac{1}{c^2} = \dots\dots\dots$ fill in the blank.   | 1 |
| 14  | Find the coordinates of the point where the line $\frac{x-5}{-2} = \frac{y-1}{3} = \frac{z-6}{-5}$ crosses the ZX- plane.   | 1 |
| 15  | Given two independent events A and B such that $P(A)=0.3$ and $P(B)=0.6$ . find $P(A \text{ and not } B)$ .   | 1 |
| 16  | Whether true or false.<br>If A and B are events such that $P(A B)=P(B A)$ , then $A \cap B = \emptyset$ .   | 1 |
| <b>SECTION II</b><br><b>Both the case study questions are compulsory.</b><br><b>Attempt any 4 sub parts from each question. Each question carries 1 mark.</b> |   |   |
| 17  | <p>An open toy box with a square base is to be made out of a given quantity of metal sheet of area <math>c^2</math>.</p>  <p>Based on the above information answer following.</p> <p>a) If x represents the side of square base and y represents the height of the toy box then the relation between the variables</p> <p>a) <math>66xy = c^2</math><br/> b) <math>x^3 = c^2</math><br/> c) <math>x^2 + 4xy = c^2</math><br/> d) <math>2xy + 4x^2 = c^2</math></p> | 4 |

|    |  |   |
|----|--|---|
|    | <p>b) The volume of the toy box V expressed as a function x is</p> <p>a) <math>V = xy^2</math></p> <p>b) <math>V = \frac{c^2x - x^3}{4}</math></p> <p>c) <math>V = \frac{x^3 - c^2x}{4}</math></p> <p>d) <math>V = \frac{x^2(c^2x - x^2)}{4}</math></p> <p>c) The maximum volume of the box is.</p> <p>a) <math>\frac{c^2}{6\sqrt{3}}</math></p> <p>b) <math>\frac{6c^2}{\sqrt{3}}</math></p> <p>c) <math>\frac{\sqrt{3}c^2}{6}</math></p> <p>d) <math>\frac{c^3}{6\sqrt{3}}</math></p> <p>d) If the box were to be closed then the relation between x and y would be</p> <p>a) <math>2x^2 + 4xy = c^2</math></p> <p>b) <math>4x^2 + 2xy = c^2</math></p> <p>c) <math>6xy = c^2</math></p> <p>d) <math>6x^2 = c^2</math></p> <p>e) If the box were to be closed then the volume of the box expressed as a function of x.</p> <p>a) <math>\frac{x^2(c^2 - 2x^2)}{4}</math></p> <p>b) <math>V = \frac{c^2x - 2x^3}{4}</math></p> <p>c) <math>V = x^3</math></p> <p>d) <math>V = \frac{2x^3 - c^2x}{4}</math></p> |   |
| 18 | <p>By examining the Covid-19 test report of some laboratory, the probability that a person is diagnosed Covid-19 positive when he is actually suffering from it is 0.99. The probability that the report incorrectly diagnosed a person to be Covid-19 positive on the basis of report is 0.001. In a certain city 300 of 1000 persons suffer from Covid-19.</p> <p>Based on the above information answer the following.</p> <p>i) The conditional probability that a person is diagnosed Covid-19 positive given that he actually has Covid-19</p>  | 4 |

|  |  |  |
|--|--|--|
|  | <p>a) 0.7<br/>b) 0.99<br/>c) 0.001<br/>d) 0.3</p>  |  |
|  | <p>ii) A person is selected at random and is diagnosed with Covid-19. What is the chance that he actually has Covid-19</p> <p>a) 0.99<br/>b) 0.91<br/>c) 297/304<br/>d) 304/297</p>  |  |
|  | <p>iii) MCD wants to keep a check, so an officer during checking selects the report randomly. According to the report, person is diagnosed Covid-19 positive. What is the probability that the person doesn't have actually Covid-19.</p> <p>a) 0.001<br/>b) 7/297<br/>c) 7/2977<br/>d) 0.99</p> |  |
|  | <p>iv) Total probability of a person being diagnosed Covid-19 positive.</p> <p>a) 1<br/>b) 0.6933<br/>c) 0.2977<br/>d) 22%</p>   |  |
|  | <p>v) Let E be the event of person being diagnosed Covid-19 positive and let E<sub>1</sub> and E<sub>2</sub> be the events that he actually has Covid-19 and he actually doesn't have Covid-19 then find <math>\sum_{i=1}^2 P(E_i E)</math></p> <p>a) 0.991<br/>b) 1<br/>c) 92%<br/>d) 0.989</p> |  |

| <b>Part- B</b>     |   |   |
|--------------------|---|---|
| <b>Section III</b> |   |   |
| 19                 | Find the value of<br>$\tan^{-1}(1) + \cos^{-1}\left(-\frac{1}{2}\right) + \sin^{-1}\left(-\frac{1}{2}\right)$   | 2 |
| 20                 | $(x \quad -5 \quad -1) \begin{pmatrix} 1 & 0 & 2 \\ 0 & 2 & 1 \\ 2 & 0 & 3 \end{pmatrix} \begin{pmatrix} x \\ 4 \\ 1 \end{pmatrix} = 0$ <p>Solve for 'x'</p> <p style="text-align: center;"><b>Or</b></p> $2 \begin{pmatrix} 3 & 4 \\ 5 & x \end{pmatrix} + \begin{pmatrix} 1 & y \\ 0 & 1 \end{pmatrix} = \begin{pmatrix} 7 & 0 \\ 10 & 5 \end{pmatrix}$ <p style="text-align: center;">Find <math>(x - y)</math>.</p> | 2 |
| 21                 | Find K so that the function $f(x) = \begin{cases} kx + 1, & \text{if } x \leq \pi \\ \cos x, & \text{if } x > \pi \end{cases}$<br>Is continuous at $x = \pi$ .  | 2 |
| 22                 | Find the slope of the normal to the curve $x = 1 - a \sin \theta, y = b \cos^2 \theta$ at $\theta = \frac{\pi}{2}$ .  | 2 |
| 23                 | Find $\int e^x \frac{\sin^4 x - \cos^4 x}{\sin x - \cos x} dx$<br><br><b>Or</b><br>Evaluate $\int_{-\frac{\pi}{2}}^{\frac{\pi}{2}} \sin^7 x dx$   | 2 |
| 24                 | What is the area lying in the first quadrant and bounded by the circle $x^2 + y^2 = 4$ and the lines $x = 0, x = 2$ .   | 2 |
| 25                 | Solve the differential equation<br>$\frac{dy}{dx} = x^3 \operatorname{cosec} y$ , given that $f(0) = 0$   | 2 |

|  |  |   |
|--|--|---|
| 26   | Find the vector equation of a plane passing through A(2, 5, -3), B(-2,-3, 5) and c(5, 3, -3).  | 2 |
| 27   | Find the distance between lines<br>$\rightarrow$<br>$r = \hat{i} + 2\hat{j} - 4\hat{k} + \lambda(2\hat{i} + 3\hat{j} + 6\hat{k})$<br>$\rightarrow$<br>and $r = 3\hat{i} + 3\hat{j} - 5\hat{k} + \mu(2\hat{i} + 3\hat{j} + 6\hat{k})$ .   | 2 |
| 28   | The random variable x has a probability distribution P(x) of the following form, where k is a number,<br>$P(X=x) = \begin{cases} k, & \text{if } x = 0 \\ 2k, & \text{if } x = 1 \\ 3k, & \text{if } x = 2 \\ 0. & \text{other wise} \end{cases}$ Determine the value of $p(X \leq 2)$ . | 2 |
| <b>Section IV</b><br><b>All questions are compulsory. In case of internal choices attempt any one.</b> |  |   |
| 29   | Show that the relation R in the set<br>$A = \{x \in Z : 0 \leq x \leq 12\}$ , given by<br>$R = \{(a, b :  a - b ) \text{ is a multiple of } 4\}$<br>Is an equivalence relation.  | 3 |
| 30   | If $y = x^a + x^{\sin x}$ , find $\frac{dy}{dx}$   | 3 |
| 31   | If $y = 3e^{2x} + 2e^{3x}$<br>Prove that $\frac{d^2y}{dx^2} - 5\frac{dy}{dx} + 6y = 0$<br><b>Or</b><br>If $x = a(\cos\theta + \theta\sin\theta)$<br>$y = a(\sin\theta - \theta\cos\theta)$<br>Find $\frac{d^2y}{dx^2}$ .   | 3 |

|  |  |   |
|--|--|---|
| 32   | Find the intervals in which the function $f$ given by $f(x) = 4x^3 - 6x^2 - 72x + 30$ is,<br><br>(a) Strictly increasing<br>(b) Strictly decreasing.   | 3 |
| 33   | Find $\int \frac{x^2}{(x^2+1)(x^2+4)} dx$  | 3 |
| 34   | Find the area of the ellipse $x^2 + 9y^2 = 36$ using integration   | 3 |
| 35   | Find the general solution of the differential equation<br><br>$e^x \tan y dx + (1 - e^x) \sec^2 y dy = 0$<br><br><b>Or</b><br>Find the general solution of the differential equation<br>$x \log x \frac{dy}{dx} + y = \frac{2}{x} \log x$  | 3 |
| <b>Section V</b><br><b>All questions are compulsory. In case of internal choices, attempt any one.</b> |  |   |
| 36   | Using matrices solve the following system of equations.<br>$x + 2y - 3z = -4$<br>$2x + 3y + 2z = 2$<br>$3x - 3y - 4z = 11$<br><br><b>Or</b><br><br>Given $A = \begin{pmatrix} 2 & 2 & -4 \\ -4 & 2 & -4 \\ 2 & -1 & 5 \end{pmatrix}$ , $B = \begin{pmatrix} 1 & -1 & 0 \\ 2 & 3 & 4 \\ 0 & 1 & 2 \end{pmatrix}$<br><br>Find $BA$ and use this to solve the system of equation.<br>$y + 2z = 7, x - y = 3, 2x + 3y + 4z = 17$ | 5 |
| 37   | Find the foot of perpendicular from the point $(2, 3, -8)$ to the line   | 5 |



$\frac{4-x}{2} = \frac{y}{6} = \frac{1-z}{3}$ . Also find the perpendicular distance from the given point to the line.

**Or**

Find the equation of the plane through the intersection of the planes  $\vec{r} \cdot (i + 3j) - 6 = 0$  and  $\vec{r} \cdot (3\hat{i} - \hat{j} - 4\hat{k}) = 0$  whose perpendicular distance from the origin is unity.


38

Maximize  $Z = x + y$   
subject to the constraints

$$\begin{aligned} x + 4y &\leq 8 \\ 2x + 3y &\leq 12 \\ 3x + y &\leq 9 \\ x \geq 0, y &\geq 0 \end{aligned}$$

**Or**

The corner points of the feasible region determined by the system of linear equations are as share below:



Answer each of the following

5

|  |  |  |
|--|--|--|
|  | <p>i. Let <math>Z = x + 2y</math> be the objective function. Find the maximum and minimum value of <math>Z</math> and also the corresponding points at which the maximum and minimum values occurs.</p> <p>ii. Let <math>Z = px + qy</math>, where <math>p, q &gt; 0</math> be the objective function. Find the condition on <math>p</math> and <math>q</math> so that the maximum <math>Z</math> occurs at <math>Q\left(\frac{3}{2}, \frac{15}{4}\right), R\left(\frac{7}{2}, \frac{3}{4}\right)</math>. Also mention the number of optimal solutions in this case.</p> |  |
|--|--|--|