CHEMISTRY THEORY (043)

PRACTICE PAPER 1

M.M 70 Time : 3 hours

General Instructions:

Read the following instructions carefully.

- a) There are 33 questions in this question paper. All questions are compulsory.
- b) Section A: Q. No. 1 to 16 are objective type questions. Q. No. 1 and 2 are passage based questions carrying 4 marks each while Q. No. 3 to 16 carry 1 mark each.
- c) Section B: Q. No. 17 to 25 are short answer questions and carry 2 marks each.
- d) Section C: Q. No. 26 to 30 are short answer questions and carry 3 marks each.
- e) Section D: Q. No. 31 to 33 are long answer questions carrying 5 marks each.
- f) There is no overall choice. However, internal choices have been provided.
- g) Use of calculators and log tables is not permitted. SECTION A (OBJECTIVE TYPE)

SECTION A: Objective Questions (1 Mark)

Passage Based Questions

1. Read the passage given below and answer the following questions: (1x4=4)

Oxygen is the most abundant of all the elements on earth. Oxygen forms about 46.6% by mass of earth's crust. Dry air contains 20.946% oxygen by volume .However, the abundance of sulphur in the earth's crust is only0.03-0.1%. Combined sulphur exists primarily as sulphates such as gypsum CaSO₄.2H₂O, epsom salt MgSO₄.7H₂O, baryte BaSO₄ and sulphides such as galena PbS, zinc blende ZnS, copper pyrites CuFeS₂ Traces of sulphur occur as hydrogen sulphide in volcanoes. Organic materials such as eggs, proteins, garlic, onion, mustard, hair and wool contain sulphur.Selenium and tellurium are also found as metal selenides and tellurides in sulphide ores. Polonium occurs in nature as a decay product of thorium and uranium minerals. Livermorium is a synthetic radioactive element.

The following questions (I - iv) are multiple choice questions. Choose the most appropriate answer:

| | | 40.000. | | ., | | | 40.000. | one choose and most appropriate another |
|-------|-------|---------------------------------|----------|----------|----------|-------------------|---------|---|
| (i) | Which | of the f | followin | g group | 16 ele | ments s | how ma | aximum catenation property? |
| | (a) | 0 | (b) | S | (c) | Se | (d) | Те |
| (ii) | Which | of the f | followin | g group | 16 ele | ments c | loes no | t exist in -2 oxidation state? |
| | (a) | 0 | (b) | S | (c) | Se | (d) | Те |
| (iii) | Which | of the f | followin | g is mo | st volat | ile? | | |
| | (a) | H ₂ O | (b) | H_2S | (c) | H ₂ Se | (d) | H ₂ Te |
| (iv) | Which | of the f | followin | g is the | wrong | statem | ent? | |
| | (a) | a) Ozone is diamagnetic gas | | | | | (b) | O ₂ is paramagnetic |
| | (c) | O ₃ molecule is bent | | | | | (d) | O – O bond lengths in ozone are unequal |
| | | | | | | | | |

2. Read the passage given below and answer the following questions: $(1 \times 4 = 4)$ Marks

The carbonyl carbon atom is sp^2 -hybridised and forms three sigma (σ) bonds. The fourth valence electron of carbon remains in its p-orbital and forms a π -bond with oxygen by overlap with p-orbital of an oxygen. In addition, the oxygen atom also has two non bonding electron pairs. Thus, the carbonyl carbon and the three atoms attached to it lie in the same plane and the π -electron cloud is above and below this plane. The bond angles are approximately 120° as expected of a trigonal coplanar structure . The carbon-oxygen double bond is polarised due to higher electronegativity of oxygen relative to carbon. Hence, the carbonyl carbon is an electrophilic (Lewis acid), and carbonyl oxygen, a nucleophilic (Lewis base) centre. Carbonyl compounds have substantial dipole moments and are polar than ethers.

In these questions (I -iv) a statement of Assertion followed by a statement of Reason is given.

Choose the correct answer out of the following choices.

- (a) Both Assertion and Reason are correct statements and Reason is the correct explanation of the assertion.
- (b) Both Assertion and Reason are correct statements, but Reason is not the correct explanation of the assertion.
- (c) Assertion is correct, but reason is incorrect statement.
- (d) Assertion is incorrect statement but reason is correct statement.
- (i) **Assertion:** Benzaldehyde is less reactive than ethanal towards nucleophilic attack.

Reason: All the carbon atoms of benzaldehyde are sp² hybridized.

(ii) Assertion: alpha hydrogen of aldehydes and ketones are slightly acidic in nature.

Reason: Enolate ion is stabilized by the resonance.

(iii) Assertion: Ketones are not oxidized by tollen's reagent

Reason: Tollen's reagent is mild oxidizing agent.

(iv) **Assertion:** Benzaldehyde is not oxidized by iodoform

Reason: Absence of methyl ketone group.

Multiple Choice questions

Following questions (No. 3 -11) are multiple choice questions carry 1 mark each:

- Q.3 The crystal field splitting energy for octahedral () and tetrahedral complex is related as
 - (a) $\Delta_{\rm t} = 0.55 \, \Delta_{\rm o}$

(b) $\Delta_{\rm t} = 0.88 \, \Delta_{\rm o}$

(c) $\Delta_t = 0.44 \Delta_o$

- (d) $\Delta_t = 0.66 \Delta_o$
- Q.4 Glucose on oxidation with Br₂ water gives
 - (a) Gluconic acid

(b) Tartaric acid

(c) Saccharic acid

(d) Meso -oxalic acid

| | In aqueous solution an amino acid exist as | | | | | | | | |
|--|--|---|---|---------------------|--|------------------|--------------------------------|-----------|------------------|
| | (a) | Cation | | | (b) | Anion | | | |
| | (c) | Zwitter ion | | | (d) | Neutr | al molecule | | |
| Q.5 | Which | of the following has highest boiling point? | | | | | | | |
| | (a) | CH₃F | (b) | CH ₃ Cl | | (c) | CH₃Br | (d) | CH₃I |
| Q.6 | Reacti | on of Phenol with chloroform in the presence of dilute sodium hydroxide | | | | | | | |
| | Finally | y introduces which one of the following functional group? | | | | | | | |
| | (a) | -COOH | | | (b) | -СНО | | | |
| | (c) | -CH ₂ Cl | | | (d) | -CHCl | 2 | | |
| Q.7 | Which | nich of the following ions is the most stable in aqueous solution ? | | | | | | | |
| | (a) | Mn ²⁺ | (b) | Cr ³⁺ | | (c) | V ³⁺ | (d) | Ti ³⁺ |
| Q.8 | EDTA | is a | | | | | | | |
| | (a) | Bidentate l | igand | | | (b) | Tridentate | ligand | |
| | (c) | Tetradenta | te ligand | | | (d) | Hexadentat | te ligand | |
| | | | | OR | | | | | |
| How many moles of AgCl will be precipitated when an excess of AgNO $_3$ is added to a molar solution of $[Cr(H_2O)_5Cl]Cl_2$? | | | | | | | | | |
| (a) | 1 mol | 1 mol of AgCl | | | (b) | 2 moles of AgCl | | | |
| (c) | 3 mole | moles of AgCl | | | (d) | 4 moles of AgCl | | | |
| Q.9 | Order | Order of basic strength of amines in aqueous solution is | | | | | | | |
| | (a) | (a) $(C_2H_5)_2NH > (C_2H_5)_3N > C_2H_5NH_2 > NH_3$ | | | | | | | |
| | (b) | $(C_2H_5)_2NH$ | > C ₂ H ₅ N | IH ₂ > (| C ₂ H ₅) ₃ N | > | NH ₃ | | |
| | (c) | (C ₂ H ₅) ₃ N | > (C ₂ H ₅) ₂ N | IH > | C ₂ H ₅ NH | 2 > | NH ₃ | | |
| | (d) | (C ₂ H ₅) ₃ N | > (C ₂ H ₅) ₂ N | IH > | NH ₃ | > C ₂ | H ₅ NH ₂ | | |
| | OR | | | | | | | | |
| Aniline on heating with chloroform and ethanolic potassium hydroxide gives | | | | | | | | | |
| (a | (a) Phenylisocyanide (b) | | | | Benzo | zonitrile | | | |
| (c) | (c) N-methylaniline (d) Chlorobenzene | | | | | | | | |
| Q.10 | An ideal solution is formed when its components: | | | | | | | | |

(a)

have no volume change on mixing

- (b) have no enthalpy change on mixing
- (c) have both of the above characteristics
- (d) have high solubility.
- Q.11 Magnetic moment of 2.84 B.M. is given by :

(At. Nos. Ni = 28, Ti = 22, Cr = 24, Co = 27)

- (a) Cr³⁺
- (b) Co²⁺
- (c) Ni²⁺
- (d) Ti³⁺

OR

Four successive members of the first series of the transition metals are listed below . For which one of them the standard potential ($E^{\circ}M^{2+}/M$) value has a positive sign?

- (a) Cu(Z=29)
- (b) Fe (Z = 26) (c)
 - Co (Z = 27)
- (d) Ni (Z = 28)

Assertion - Reason

In the following questions (Q.No. 12 -16) a statement of Assertion followed by a statement of Reason is given .Choose the correct answer out of the following Choices

- (a) Both Assertion and Reason are correct statements and Reason is the correct explanation of the assertion.
- (b) Both Assertion and Reason are correct statements, but Reason is not the correct explanation of the assertion.
- (c) Assertion is correct, but reason is incorrect statement.
- (d) Assertion is incorrect statement but reason is correct statement.
- Q.12 Assertion: Hydrazones of acetaldehydes are not prepared in highly acidic medium

Reason: Hydrazines are basic in nature.

Q.13 **Assertion:** p-Nitrophenol is stronger acid than o-Nitrophenol.

Reason: Intramolecular H- bonding makes o-isomer weaker than p-isomer.

OR

Assertion: C-O-C bond angle in ether is higher than water

Reason: Greater repulsion between bulkier alkyl groups as compared to smaller H atoms

Q.14 Assertion: Aquatic species are more comfortable in cold water as compared to warm water

Reason: The solubility of oxygen in water decrease with increase in temperature.

Q.15 **Assertion:** Glucose does not give DNP test.

Reason: It form hemiacetal between the -CHO group & -OH group on the C₅ atom.

Q.16 **Assertion:** Electron gain enthalpy of Cl is more negative than fluorine.

Reason: Fluorine is more electronegative than chlorine.

SECTION B: Short Answer Type I Questions (2 Marks)

| JLCII | CIVE | . Short Answer Type I Questions (2 Marks) | | | | | | | | |
|-------|--|--|--|--|--|--|--|--|--|--|
| Q.17. | Give rea | asons | | | | | | | | |
| (i) | Why is methylchloride hydrolysed more easily than chlorobenzene ? | | | | | | | | | |
| (ii) | Why p-Nitrochlorobenzene is more reactive than Chlorobenzene towards nucleophilic substitution reactions ? | | | | | | | | | |
| | | OR | | | | | | | | |
| Carry | out the | following conversions in not more than 2 steps: | | | | | | | | |
| | (i) Ethanol to But-1-yne | | | | | | | | | |
| | (ii) | Propene to 1-lodopropane | | | | | | | | |
| Q.18 | Draw the structures of the following: | | | | | | | | | |
| | (i) | XeO ₃ (ii) HClO ₃ | | | | | | | | |
| Q.19 | A met | etal oxide has the formula $M_{0.96}O_{1.00}$. What fractions of the metal exist as M^{2+} and M^{3+} ions ? | | | | | | | | |
| Q.20 | | Define azeotropes. What type of azeotrope is formed by negative deviation from Raoult's law ? Give an example | | | | | | | | |
| Q.21 | An op | tically active compound having molecular formula C ₇ H ₁₅ Br reacts with aqueous | | | | | | | | |
| | KOH to give a racemic mixture of products. Write the mechanism involved in the reaction. | | | | | | | | | |
| Q.22 | Explai | Explain the following with example | | | | | | | | |
| | (i) | Kolbe's reaction | | | | | | | | |
| | (ii) | Williamson's ether synthesis | | | | | | | | |
| Q.23 | Write one difference in each of the following: (i) Coagulation and peptisation (ii) Lyophobic and lyophillic colloid. | | | | | | | | | |
| | OR Write the dispersed phase and dispersion medium of the following colloidal system: (a) Smoke (ii) Milk | | | | | | | | | |
| Q.24 | (i) | (i) Write the IUPAC name of the following complex: | | | | | | | | |
| | [Cr(NH ₃) ₂ Cl ₂ (en)]Cl | | | | | | | | | |
| | (ii) Write the formula for the following complex: | | | | | | | | | |
| | Pentaamminenitrito-Ocobalt(III) | | | | | | | | | |
| | | OR | | | | | | | | |
| | - | oin only magnetic moment of $[MnBr_4]^{2-}$ is 5.9 B.M. Predict the geometry of the complex atomic number of $Mn = 25$ | | | | | | | | |

Q.25 Explain what is observed, when

(ii) KCl, an electrolyte is added to hydrated ferric oxide sol. SECTION - C Q.26 Explain the following: ICl is more reactive than I₂ NaBiO₃ is stronger oxidizing agent Noble gases have very low boiling point. Q.27 Define the following as related to protiens. Denaturation (a) Peptide linkage (b) Primary structure (c) Q.28 The elements of 3d transition series are given as: Τi, ٧, Cr Mn Fe Co Zn Sc, Ni Cu Answer the following: Write the element which shows maximum number of oxidation states. Give reason. Which element has the highest melting point and why? Which element is a strong oxidizing agent in +3 oxidation state and why? OR Explain giving reasons: Which is stronger reducing agent Cr²⁺ or Fe²⁺? A compound of vanadium has a magnetic moment of 1.73 B.M. Work out the electronic configuration of vanadium ion in the compound. The transition metals generally form coloured compounds. An aromatic compound 'A' on treatment with aqueous ammonia and heating forms compound Q.29 'B' which on heating with Br_2 and KOH forms a compound 'C' of molecular formula C_6H_7N . Write the structures and IUPAC names of compounds A, B and C OR Account for the following: pKb of aniline is more than that of methylamine

SECTION D: Long Answer Type Questions (5 Marks)

length of its unit cell is 300 pm, identify the type of unit cell.

Gabriel phthalimide synthesis is preferred for synthesising primary amines.

An element has atomic mass 93 g mol⁻¹ and density 11.5 g cm⁻³. If the edge

Aniline does not undergo Friedel-Crafts reaction.

(i)

(a)

(b)

(c)

(i)

(ii)

(iii)

(i)

(ii)

(iii)

(i)


(ii)

(iii)

Q.30

an electric current is passed through a sol.

- Q.31 (a) Define the term Molar conductivity of a solution and explain how molar conductivity change with change in concentration of a solution for weak and strong electrolyte.
- (a) State Kohlrausch's law of independent migration of ions. Write an expression for the molar conductivity of formic acid at infinite dilution according to Kohlrausch's law.
- (b) The molar conductivity of 0.025 mol L=1 methanoic acid is 46.1 S cm2 mol=1. Calculate its degree of dissociation and dissociation constant. Given $\lambda 0$ (H+) = 349.6 S cm2 mol=1 and $\lambda 0$ (HCOO=) = 54.6 S cm2 mol=1
- Q.32 (a) Write the structures of A,B,C,D &E in the following reactions:



- (a) Write the structures of the main products when ethanal (CH₃CHO) reacts with the following Reagents.
 - (i) HCN (ii) dil. NaOH (iii) Zn Hg / conc.HCl
- (b) Arrange the following in increasing order of their reactivity towards nucleophilic addition reaction: Ethanal, Propanal, Propanone, Butanone

(5)

- (c) Give a chemical test to distinguish between following pair of compounds
 Propanal and Propanone
- Q.33 (a)A first order reaction takes 40 minutes for 30 % decomposition. Calculate $t_{1/2}$.
 - (b) The following results have been obtained during the kinetics of the reaction:

$$2A + B \longrightarrow C + D$$

| Experiment No. | [A] | [B] | Initial Rate of formation of D |
|----------------|-------|-------|---|
| 1 | 0.1 M | 0.1 M | 6.0x10 ⁻³ M min ⁻¹ |
| 2 | 0.3 M | 0.2 M | 7.2x10 ⁻² M min ⁻¹ |
| 3 | 0.3 M | 0.4 M | 2.88x10 ⁻¹ M min ⁻¹ |
| 4 | 0.4 M | 0.1 M | 2.4x10 ⁻² M min ⁻¹ |

Determine the Rate law and the rate constant for the reaction:

OR

(a) Rate constant k for first order reaction has been found to be $2.54 \times 10^{-3} \text{ s}^{-1}$. Calculate its three fourth life

