

**Class: XII Session: 2020-21**

**Subject: Mathematics**

**Practice Question Paper 5  
(Theory)**

**Time Allowed: 3 Hours**

**Maximum Marks: 80**

**General Instructions:**

1. This question paper contains two **parts A and B**. Each part is compulsory. Part A carries **24** marks and Part B Carries **56** marks
2. **Part-A** has Objective Type Questions and **Part-B** has Descriptive Type Questions.
3. Both Part A and Part B have choices.


**Part-A:**

1. It consists of two sections- **I and II**
2. Section **I** comprises of 16 very short answer type questions.
3. Section **II** contains **2** case studies. Each case study comprise of 4 case-based MCQs.

**Part –B:**

1. It consists of three sections- **III, IV and V**.
2. Section **III** comprises of 10 questions of **2 marks** each.
3. Section **IV** comprises of 7 question of **3 marks** each.
4. Section **V** comprises of 3 questions of **5 marks** each.
5. Internal choice is provided in **3** questions of Section –III, **2** questions of Section- IV and **3** questions of Section-V. You have to attempt only one of the alternatives in all such questions

Sr. No.	Part-A	Marks
	<b>Section I</b> <b>All questions are compulsory. In case of internal choices attempt any one.</b>	
1.	If A is a square matrix of 3 order and $ A =5$ , then find the value of $ 2A $ <p style="text-align: center;"><b>OR</b></p> If $A = \begin{bmatrix} 5 & x \\ y & 0 \end{bmatrix}$ and $A = A'$ , what will be the relation between x and y.	1
2.	An equivalence relation R in A is divided into equivalence classes A1, A2, A3. What is the value of $A1 \cup A2 \cup A3$ and $A1 \cap A2 \cap A3$ .	1
3.	If matrix $A = \begin{bmatrix} 0 & a & -3 \\ 2 & 0 & -1 \\ b & 1 & 0 \end{bmatrix}$ is skew symmetric, find the value of a and b. <p style="text-align: center;"><b>OR</b></p> If $A = \begin{bmatrix} \alpha & \beta \\ \gamma & -\alpha \end{bmatrix}$ is such that $A^2 = I$ . Find the relation between $\alpha, \beta, \gamma$ .	1
4.	What is the property of relation if each element of A is related to itself?	1
5.	Check if relation R in the set R of real numbers defined as $R = \{(a,b) : a < b\}$ is symmetric.	1
6.	Suppose P and Q are two different matrices of order $3 \times n$ and $n \times p$ , then what will be the order of $P \times Q$ ?	1
7.	Evaluate $\int x^2 e^{x^3} dx$ <p style="text-align: center;"><b>OR</b></p> Find the value of $\int_0^{\pi/2} \cos x e^{\sin x} dx$	1
8.	A card is picked at random from a pack of 52 cards. Given that picked card is queen, find the probability to be spade.	1
9.	A die is thrown once. Let A be the event that number obtained is greater than 3. Let B be the event that number obtained is less than 5. Find the value of $P(A \cup B)$ .	1

10.	Find the vector equation of line which passes through point (3,4,5) and is parallel to vector $2\hat{i}+2\hat{j}-3\hat{k}$ . <b>OR</b> Find the distance of a point P (a,b,c) from x axis.	1
11.	Find the order and degree of differential equation $x^2 \left[ \frac{d^2y}{dx^2} \right] = \left[ 1 + \left( \frac{dy}{dx} \right)^2 \right]^4$	1
12.	A line makes an angle $\alpha, \beta, \gamma$ with x axis, y axis and z axis. Find the value of $\cos 2\alpha + \cos 2\beta + \cos 2\gamma$ .	1
13.	If line $\frac{x-1}{2} = \frac{y+3}{1} = \frac{z-5}{-1}$ is parallel to plane $px+3y-z+5=0$ . Find the value of p.	1
14.	Evaluate $\int_{-\pi/2}^{\pi/2} x^2 \sin x dx$ <b>OR</b> $\int_2^8 \frac{\sqrt{10-x}}{\sqrt{x} + \sqrt{10-x}} dx$	1
15.	Find the area bounded by curve $y=x^2$ , the x axis and lines $x=-1$ and $x=1$ . <b>OR</b> Find the area region bounded by curve $x=y^2$ , y axis, line $y=3$ and $y=4$ .	1
16.	For what value of n will the following line be a homogenous differential equations $\frac{dy}{dx} = \frac{x^3 - y^n}{x^2y + xy^2}$ <b>OR</b> Find the integrating factor of differential equation $\frac{dy}{dx} (x \log x) + y = 2 \log x$	1
	<b>SECTION – II</b> <b>Case study based MCQ's ( There are 2 quetions of case studies with 5 MCQ's each, all are compulsory)</b>	
17.	A rectangle of perimeter 36 cm is rolled out to form cylinder of volume as large as possible. Based on this information, answer the following: 	

	<p>(i) If <math>x</math> and <math>y</math> represent the length and breadth of rectangular region, then relation between variable is:</p> <p>a) <math>X=18-y</math>  b) <math>Y=18-x</math>  c) <math>2y=18-2x</math>  d) <math>2x=36-y</math></p>	1
	<p>(ii) Volume of formed cylinder <math>V</math> expressed as function of <math>x</math> is:</p> <p>a) <math>V(x) = \pi x^3(x - 18)</math>  b) <math>V(x) = \pi(18x^2 - 18)</math>  c) <math>V(x) = \pi(18x^2 - x^3)</math>  d) <math>V(x) = \pi(18x^3 - x^2)</math></p>	1
	<p>(iii) The dimensions of rectangle for maximum volume should be:</p> <p>a) <math>X=12, y=6</math>  b) <math>X=10, y=8</math>  c) <math>X=8, y=2</math>  d) <math>X=11, y=7</math></p>	1
	<p>(iv) Maximum volume of cylinder:</p> <p>a) <math>846\pi \text{ cm}^3</math>  b) <math>864\pi \text{ cm}^3</math>  c) <math>684\pi \text{ cm}^3</math>  d) <math>866\pi \text{ cm}^3</math></p>	1
	<p>(v) The value of <math>\frac{d^2v}{dx^2}</math> at maximum point is</p> <p>(a) <math>-36\pi</math>  (b) <math>-63\pi</math>  (c) <math>36\pi</math>  (d) <math>63\pi</math></p>	1
18.	<p>There are 3 urns containing 2 white and 3 black balls, 3 white and 2 black balls and 4 white and 1 black balls respectively. There is an equal probability of each urn being chosen. A ball is drawn at random from the chosen urn and it is found to be white. There are 3 urns <math>U_1, U_2, U_3</math></p> <p><math>E_1</math> – an event where ball is chosen from <math>U_1</math>  <math>E_2</math> - an event where ball is chosen from <math>U_2</math>  <math>E_3</math> - an event where ball is chosen from <math>U_3</math>  <math>E</math> - an event where white ball is drawn</p> <p>Based on this information, answer the following:</p>	


	i) What is the value of $P(U_1)$ ? a) $2/3$ b) $3/3$ c) $1/3$ d) $3/2$	1
	ii) Calculate $P(E_2)$ a) $2/3$ b) $1/3$ c) $2/5$ d) $4/5$	1
	iii) The value of $P(E/E_1)$ and $P(E/E_3)$ will be? a) $4/5, 2/5$ b) $2/5, 3/5$ c) $3/5, 4/5$ d) $2/5, 4/5$	1
	iv) Find the probability that ball drawn was from second urn? a) $1/3$ b) $4/5$ c) $3/15$ d) $3/5$	1
	v) What is the value of $P(E_2).P(E/E_2)$ ?	1
	<b>PART B</b> <b>SECTION III</b>	
19.	Express $\tan^{-1}\left(\frac{\cos x}{1-\sin x}\right) : \frac{-3\pi}{2} < x < \frac{\pi}{2}$ in simplest form	2
20.	Find a matrix A such that $2A-3B+5C=0$ where $B = \begin{bmatrix} -2 & 2 & 0 \\ 3 & 1 & 4 \end{bmatrix}$ And $C = \begin{bmatrix} 2 & 0 & -2 \\ 7 & 1 & 6 \end{bmatrix}$  OR Solve for x and y $x \begin{bmatrix} 2 \\ 1 \end{bmatrix} + y \begin{bmatrix} 3 \\ 5 \end{bmatrix} + \begin{bmatrix} -8 \\ 11 \end{bmatrix} = 0$	2
21.	If function f is defined as: $f(x) = f(x) = \begin{cases} \frac{x^2-9}{x-3}, & x \neq 3 \\ k, & x = 3 \end{cases}$ is continuous at $x=3$ . Find k	2
22.	Find the equation of tangent and normal to the curve $16x^2 + 9y^2 = 145$ at (2,3)	2

23.	Evaluate $\int \frac{\cos 2x + 2 \sin^2 x}{\cos^2 x} dx$ OR $\int_0^a \frac{dx}{1+4x^2} = \frac{\pi}{8}$ Find the value of a	2
24.	Find the area bounded by a parabola $y^2 = x$ and straight line $2y=x$	2
25.	Find the particular solution of the differential equation $\frac{dy}{dx} + 2y \tan x = \sin x$ at $y=0$ and $x= \pi/3$	2
26.	Using vector show P(2,-1,3), Q(3,-5,1) and R(-1,11,9) are collinear	2
27.	Find the vector equation of plane that passes through the point (1,0,0) and contains the line $\vec{r} - \lambda \hat{j}$	2
28.	Given E, F are events such that $P(E)=0.8$ , $P(F)=0.7$ , $P(E \cap F)=0.6$ . Find $P(\bar{E}/\bar{F})$ OR If $P(A \cap B) = \frac{7}{10}$ , $P(B) = \frac{17}{20}$ find $P\left(\frac{A}{B}\right)$	2
<b>SECTION IV</b> <b>7 questions of 3 marks each</b>		
29.	Show that function f in $A = \mathbb{R} - \{2/3\}$ defined as function $f(x) = \frac{4x+3}{6x-4}$ is one-one and onto.	3
30.	Prove that curve $x=y^2$ and $xy=k$ cut at right angles if $8k^2 = 1$ OR Differentiate $\tan^{-1} \sqrt{\frac{1-\cos x}{1+\cos x}}$ , $-\frac{\pi}{4} < x < \frac{\pi}{4}$	3
31.	Using integration, find the area of region in the first quadrant enclosed by x axis, the line $y=x$ and circle $x^2 + y^2 = 32$	3
32.	Find the interval in which the function $f(x) = \frac{x^4}{4} - x^3 - 5x^2 + 24x + 2$ is: a) Strictly increasing b) Strictly decreasing	3

33.	<p>Solve <math>\int \frac{2 \cos x + 1}{(1 - \sin x)(1 + \sin^2 x)} dx</math></p> <p>OR</p> <p>If <math>\int \frac{dx}{(x+2)(x^2+1)} = a \log 1 + x^2  + b \tan^{-1} x + \frac{1}{5} \log x + 2  + C</math> Find the values of a and b</p>	3
34.	Solve the differential equation $x dy - y dx = \sqrt{x^2 + y^2} dx$	3
35.	If $\log(x^2 + y^2) = 2 \tan^{-1} \left(\frac{y}{x}\right)$ , show that $\frac{dy}{dx} = \frac{x+y}{x-y}$	3
<p><b>SECTION V</b></p> <p><b>3 questions of 5 marks each</b></p>		
36	<p>Evaluate the product AB where <math>A = \begin{bmatrix} 1 &amp; -1 &amp; 0 \\ 2 &amp; 3 &amp; 4 \\ 0 &amp; 1 &amp; 2 \end{bmatrix}</math> <math>B = \begin{bmatrix} 2 &amp; 2 &amp; -4 \\ -4 &amp; 2 &amp; -4 \\ 2 &amp; -1 &amp; 5 \end{bmatrix}</math>.</p> <p>Hence, solve the system of linear equation:</p> <p><math>x - y = 3</math>  <math>2x + 3y + 4z = 17</math>  <math>y + 2z = 7</math></p> <p style="text-align: center;">OR</p> <p>Solve the following system of equations</p> $\begin{bmatrix} 3 & 0 & 3 \\ 2 & 1 & 0 \\ 4 & 0 & 2 \end{bmatrix} \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} 8 \\ 1 \\ 4 \end{bmatrix} + \begin{bmatrix} 2y \\ z \\ 3y \end{bmatrix}$	5
37	<p>Maximize <math>Z = 15x + 10y</math></p> <p>Subject to</p> <p><math>3x + 2y \leq 80</math></p> <p><math>2x + 3y \leq 70</math></p> <p><math>x, y \geq 0</math></p> <p style="text-align: center;">OR</p> <p>Find the coordinates of foot of perpendicular drawn from origin to the plane <math>3y + 4z - 6 = 0</math></p>	5
38	Find the Cartesian equation of plane passing through points $(2, 2, -1)$ , $(3, 4, 2)$ , $(7, 0, 6)$ . Also find the vector equation of a plane passing through $(4, 3, 1)$ and parallel to the plane obtained above.	5

OR

The corner points of the feasible region determined by the system of linear constraints are as shown below:



Answer each of the following:

- (i) Let  $Z = 3x - 4y$  be the objective function. Find the maximum and minimum value of  $Z$  and also the corresponding points at which the maximum and minimum value occurs.
- (ii) Let  $Z = px + qy$ , where  $p, q > 0$  be the objective function. Find the condition on  $p$  and  $q$  so that the maximum value of  $Z$  occurs at  $B(4,10)$  and  $C(6,8)$ . Also mention the number of optimal solutions in this case.



