

**DIRECTORATE OF EDUCATION**  
**Govt. of NCT, Delhi**

**SUPPORT MATERIAL**  
**( 2022-2023)**

**Class : XII**

**CHEMISTRY**

Under the Guidance of

**Shri Ashok Kumar**  
Secretary (Education)

**Shri Himanshu Gupta**  
Director (Education)

**Dr. Rita Sharma**  
Addl. DE (School & Exam.)

**Coordinators**

**Mr. Sanjay Subhas Kumar**  
DDE (Exam)

**Mrs. Sunita Dua**  
OSD (Exam)

**Mr. Raj Kumar**  
OSD (Exam)

**Mr. Krishan Kumar**  
OSD (Exam)



**Production Team**  
Anil Kumar Sharma

---

Published at Delhi Bureau of Text Books, 25/2, Institutional Area, Pankha Road, New Delhi-58 by **Rajesh Kumar**, Secretary, Delhi Bureau of Text Books and Printed at: Supreme Offset Press, 133, Udhog Kendra Ext.-1, Greater Noida, U.P.

**ASHOK KUMAR  
IAS**



सचिव ( शिक्षा )  
राष्ट्रीय राजधानी क्षेत्र  
दिल्ली सरकार  
पुराना सचिवालय, दिल्ली-110054  
दूरभाष : 23890187 टेलीफैक्स : 23890119

Secretary (Education)  
Government of National Capital Territory of Delhi  
Old Secretariat, Delhi-110054  
Phone : 23890187 Telefax : 23890119  
e-mail : secyedu@nic.in

### **MESSAGE**

Remembering the words of John Dewey, "Education is not preparation for life, education is life itself, I highly commend the sincere efforts of the officials and subject experts from Directorate of Education involved in the development of Support Material for classes IX to XII for the session 2022-23.

The Support Material is a comprehensive, yet concise learning support tool to strengthen the subject competencies of the students. I am sure that this will help our students in performing to the best of their abilities.

I am sure that the Heads of School and teachers will motivate the students to utilise this material and the students will make optimum use of this Support Material to enrich themselves.

I would like to congratulate the team of the Examination Branch along with all the Subject Experts for their incessant and diligent efforts in making this material so useful for students.

I extend my Best Wishes to all the students for success in their future endeavours.

**(Ashok Kumar)**

**HIMANSHU GUPTA, IAS**  
Director, Education & Sports



Directorate of Education  
Govt. of NCT of Delhi  
Room No. 12, Civil Lines  
Near Vidhan Sabha,  
Delhi-110054  
Ph.: 011-23890172  
E-mail: diredu@nic.in

## MESSAGE

**“A good education is a foundation for a better future.”**

**- Elizabeth Warren**

Believing in this quote, Directorate of Education, GNCT of Delhi tries to fulfill its objective of providing quality education to all its students.

Keeping this aim in mind, every year support material is developed for the students of classes IX to XII. Our expert faculty members undertake the responsibility to review and update the Support Material incorporating the latest changes made by CBSE. This helps the students become familiar with the new approaches and methods, enabling them to become good at problem solving and critical thinking. This year too, I am positive that it will help our students to excel in academics.

The support material is the outcome of persistent and sincere efforts of our dedicated team of subject experts from the Directorate of Education. This Support Material has been especially prepared for the students. I believe its thoughtful and intelligent use will definitely lead to learning enhancement.

Lastly, I would like to applaud the entire team for their valuable contribution in making this Support Material so beneficial and practical for our students.

Best wishes to all the students for a bright future.

**(HIMANSHU GUPTA)**

**Dr. RITA SHARMA**  
Additional Director of Education  
(School/Exam)



**Govt. of NCT of Delhi**  
Directorate of Education  
Old Secretariat, Delhi-110054  
Ph. : 23890185

D.O. No. PS/Addl.DE/Sch/2022/131  
Dated: 01 सितम्बर, 2022

### संदेश

शिक्षा निदेशालय, दिल्ली सरकार का महत्वपूर्ण लक्ष्य अपने विद्यार्थियों का सर्वांगीण विकास करना है। इस उद्देश्य को ध्यान में रखते हुए शिक्षा निदेशालय ने अपने विद्यार्थियों को उच्च कोटि के शैक्षणिक मानकों के अनुरूप विद्यार्थियों के स्तरानुकूल सहायक सामग्री कराने का प्रयास किया है। कोरोना काल के कठिनतम समय में भी शिक्षण अधिगम की प्रक्रिया को निर्बाध रूप से संचालित करने के लिए संबंधित समस्त अकादमि समूहों और क्रियान्वित करने वाले शिक्षकों को हार्दिक बधाई देती हूँ।

प्रत्येक वर्ष की भाँति इस वर्ष भी कक्षा 9वीं से कक्षा 12वीं तक की सहायक सामग्रियों में सी.बी.एस.ई के नवीनतम दिशा-निर्देशों के अनुसार पाठ्यक्रम में आवश्यक संशोधन किए गए हैं। साथ ही साथ मूल्यांकन से संबंधित आवश्यक निर्देश भी दिए गए हैं। इन सहायक सामग्रियों में कठिन से कठिन सामग्री को भी सरलतम रूप में प्रस्तुत किया गया है ताकि शिक्षा निदेशालय के विद्यार्थियों को इसका भरपूर लाभ मिल सके।

मुझे आशा है कि इन सहायक सामग्रियों के गहन और निरंतर अध्ययन के फलस्वरूप विद्यार्थियों में गुणात्मक शैक्षणिक संवर्धन का विस्तार उनके प्रदर्शन में भी परिलक्षित होगा। इस उत्कृष्ट सहायक सामग्री को तैयार करने में शामिल सभी अधिकारियों तथा शिक्षकों को हार्दिक बधाई देती हूँ तथा सभी विद्यार्थियों को उनके उज्ज्वल भविष्य की शुभकामनाएं देती हूँ।

**रीता शर्मा**  
(रीता शर्मा)



**DIRECTORATE OF EDUCATION**  
**Govt. of NCT, Delhi**

**SUPPORT MATERIAL**  
**( 2022-2023)**

**CHEMISTRY**

**Class : XII**

**NOT FOR SALE**

---

**PUBLISHED BY : DELHI BUREAU OF TEXTBOOKS**



# भारत का संविधान

## भाग 4क

### नागरिकों के मूल कर्तव्य

#### अनुच्छेद 51 क

**मूल कर्तव्य** - भारत के प्रत्येक नागरिक का यह कर्तव्य होगा कि वह -

- (क) संविधान का पालन करे और उसके आदर्शों, संस्थाओं, राष्ट्रध्वज और राष्ट्रगान का आदर करे;
- (ख) स्वतंत्रता के लिए हमारे राष्ट्रीय आंदोलन को प्रेरित करने वाले उच्च आदर्शों को हृदय में संजोए रखे और उनका पालन करे;
- (ग) भारत की संप्रभुता, एकता और अखंडता की रक्षा करे और उसे अक्षुण्ण बनाए रखे;
- (घ) देश की रक्षा करे और आह्वान किए जाने पर राष्ट्र की सेवा करे;
- (ङ) भारत के सभी लोगों में समरसता और समान भ्रातृत्व की भावना का निर्माण करे जो धर्म, भाषा और प्रदेश या वर्ग पर आधारित सभी भेदभावों से परे हो, ऐसी प्रथाओं का त्याग करे जो महिलाओं के सम्मान के विरुद्ध हों;
- (च) हमारी सामासिक संस्कृति की गौरवशाली परंपरा का महत्त्व समझे और उसका परिरक्षण करे;
- (छ) प्राकृतिक पर्यावरण की, जिसके अंतर्गत वन, झील, नदी और वन्य जीव हैं, रक्षा करे और उसका संवर्धन करे तथा प्राणिमात्र के प्रति दयाभाव रखे;
- (ज) वैज्ञानिक दृष्टिकोण, मानववाद और ज्ञानार्जन तथा सुधार की भावना का विकास करे;
- (झ) सार्वजनिक संपत्ति को सुरक्षित रखे और हिंसा से दूर रहे;
- (ञ) व्यक्तिगत और सामूहिक गतिविधियों के सभी क्षेत्रों में उत्कर्ष की ओर बढ़ने का सतत् प्रयास करे, जिससे राष्ट्र निरंतर बढ़ते हुए प्रयत्न और उपलब्धि की नई ऊँचाइयों को छू सके; और
- (ट) यदि माता-पिता या संरक्षक हैं, छह वर्ष से चौदह वर्ष तक की आयु वाले अपने, यथास्थिति, बालक या प्रतिपाल्य को शिक्षा के अवसर प्रदान करे।





# Constitution of India

## Part IV A (Article 51 A)

### Fundamental Duties


It shall be the duty of every citizen of India —

- (a) to abide by the Constitution and respect its ideals and institutions, the National Flag and the National Anthem;
- (b) to cherish and follow the noble ideals which inspired our national struggle for freedom;
- (c) to uphold and protect the sovereignty, unity and integrity of India;
- (d) to defend the country and render national service when called upon to do so;
- (e) to promote harmony and the spirit of common brotherhood amongst all the people of India transcending religious, linguistic and regional or sectional diversities; to renounce practices derogatory to the dignity of women;
- (f) to value and preserve the rich heritage of our composite culture;
- (g) to protect and improve the natural environment including forests, lakes, rivers, wildlife and to have compassion for living creatures;
- (h) to develop the scientific temper, humanism and the spirit of inquiry and reform;
- (i) to safeguard public property and to abjure violence;
- (j) to strive towards excellence in all spheres of individual and collective activity so that the nation constantly rises to higher levels of endeavour and achievement;
- \*(k) who is a parent or guardian, to provide opportunities for education to his child or, as the case may be, ward between the age of six and fourteen years.

---

**Note:** The Article 51A containing Fundamental Duties was inserted by the Constitution (42nd Amendment) Act, 1976 (with effect from 3 January 1977).

\*(k) was inserted by the Constitution (86th Amendment) Act, 2002 (with effect from 1 April 2010).



# भारत का संविधान

## उद्देशिका

हम, भारत के लोग, भारत को एक <sup>1</sup>[संपूर्ण प्रभुत्व-संपन्न समाजवादी पंथनिरपेक्ष लोकतंत्रात्मक गणराज्य] बनाने के लिए, तथा उसके समस्त नागरिकों को :

सामाजिक, आर्थिक और राजनैतिक न्याय,

विचार, अभिव्यक्ति, विश्वास, धर्म

और उपासना की स्वतंत्रता,

प्रतिष्ठा और अवसर की समता

प्राप्त कराने के लिए,

तथा उन सब में

व्यक्ति की गरिमा और <sup>2</sup>[राष्ट्र की एकता

और अखंडता] सुनिश्चित करने वाली बंधुता

बढ़ाने के लिए

दृढसंकल्प होकर अपनी इस संविधान सभा में आज तारीख 26 नवंबर, 1949 ई. को एतद्वारा इस संविधान को अंगीकृत, अधिनियमित और आत्मार्पित करते हैं।

1. संविधान (बयालीसवां संशोधन) अधिनियम, 1976 की धारा 2 द्वारा (3.1.1977 से) “प्रभुत्व-संपन्न लोकतंत्रात्मक गणराज्य” के स्थान पर प्रतिस्थापित।
2. संविधान (बयालीसवां संशोधन) अधिनियम, 1976 की धारा 2 द्वारा (3.1.1977 से) “राष्ट्र की एकता” के स्थान पर प्रतिस्थापित।

# **THE CONSTITUTION OF INDIA**

## **PREAMBLE**

**WE, THE PEOPLE OF INDIA**, having solemnly resolved to constitute India into a <sup>1</sup>**[SOVEREIGN SOCIALIST SECULAR DEMOCRATIC REPUBLIC]** and to secure to all its citizens :

**JUSTICE**, social, economic and political;

**LIBERTY** of thought, expression, belief, faith and worship;

**EQUALITY** of status and of opportunity; and to promote among them all

**FRATERNITY** assuring the dignity of the individual and the <sup>2</sup>[unity and integrity of the Nation];

**IN OUR CONSTITUENT ASSEMBLY** this twenty-sixth day of November, 1949 do **HEREBY ADOPT, ENACT AND GIVE TO OURSELVES THIS CONSTITUTION.**

1. Subs. by the Constitution (Forty-second Amendment) Act, 1976, Sec.2, for "Sovereign Democratic Republic" (w.e.f. 3.1.1977)
2. Subs. by the Constitution (Forty-second Amendment) Act, 1976, Sec.2, for "Unity of the Nation" (w.e.f. 3.1.1977)

**CHEMISTRY**  
**CLASS-XII**  
**2022-2023**

**LIST OF MEMBERS WHO REVIEWED AND  
REVISED SUPPORT MATERIAL OF CHEMISTRY**

S. No.	Name	Designation
1	<b>Dr. Hans Raj Modi</b> (Group Leader) (ID : 20025021)	OSD, NIOS HQ, Directorate of Education
2	<b>Mr. Mukesh Kumar Kaushik</b> (Member) (ID : 20092607)	Lecturer RPVV, Narela (ID : 1310409)
3	<b>Mr. Harpreet Singh</b> (Member) (ID : 20092637)	Lecturer RPVV, Kishan Ganj (ID : 1208092)
4	<b>Ms. Akshma</b> (Member) (ID : 20171208)	Lecturer SOE, Sector-22 Dwarka (ID : 1821282)
5	<b>Mohd Shahwez Siddiqui</b> (Member) (ID : 20192038)	Lecturer RPVV, Link Road Karol Bagh (ID : 2128031)



## CONTENTS

S.No.	Units	Page No
1	Solutions	1
2	Electrochemistry	33
3	Chemical Kinetics	69
4	d- and f-Block Elements	103
5	Coordination Compounds	126
6	Haloalkanes and Haloarenes	155
7	Alcohols, Phenols and Ethers	184
8	Aldehydes, Ketones and Carboxylic Acids	221
9	Amines	256
10	Biomolecules	290
11	CBSE Sample Paper	316
12	Marking Scheme	326
13	Unsolved Sample Paper	334

**SYLLABUS FOR SESSION 2022-23**  
**CLASS XII (THEORY)**

**Time: 3 Hours**

**70 Marks**

UNIT No.	UNIT	No. of Periods	Marks
2	Solutions	15	7
3	Electrochemistry	18	9
4	Chemical Kinetics	15	7
8	d -and f -Block Elements	18	7
9	Coordination Compounds	18	7
10	Haloalkanes and Haloarenes	15	6
11	Alcohols, Phenols and Ethers	14	6
12	Aldehydes, Ketones and Carboxylic Acids	15	8
13	Amines	14	6
14	Biomolecules	18	7
	<b>Total</b>	160	<b>70</b>

**Unit II Solutions**

**15 Periods**

Types of solutions, expression of concentration of solutions of solids in liquids, solubility of gases in liquids, solid solutions, Raoult's law, colligative properties - relative lowering of vapour pressure, elevation of boiling point, depression of freezing point, osmotic pressure, determination of molecular masses using colligative properties, abnormal molecular mass, Van't Hoff factor.

**Unit III Electrochemistry**

**18 Periods**

Redox reactions, EMF of a cell, standard electrode potential, Nernst equation and its application to chemical cells, Relation between Gibbs energy change and EMF of a cell, conductance in electrolytic solutions, specific and molar conductivity, variations of conductivity with concentration, Kohlrausch's Law, electrolysis and law of electrolysis (elementary idea), dry cell-electrolytic cells and Galvanic cells, lead accumulator, fuel cells, **corrosion**.



**Unit IV Chemical Kinetics****15 Periods**

Rate of a reaction (Average and instantaneous), factors affecting rate of reaction: concentration, temperature, catalyst; order and molecularity of a reaction, rate law and specific rate constant, integrated rate equations and half-life (only for zero and first order reactions), concept of collision theory (elementary idea, no mathematical treatment), activation energy, Arrhenius equation.

**Unit VIII d and f Block Elements****18 Periods**

General introduction, electronic configuration, occurrence and characteristics of transition metals, general trends in properties of the first-row transition metals — metallic character, ionization enthalpy, oxidation states, ionic radii, colour, catalytic property, magnetic properties, interstitial compounds, alloy formation, preparation and properties of  $K_2Cr_2O_7$  and  $KMnO_4$ .

**Lanthanoids** - Electronic configuration, oxidation states, chemical reactivity and lanthanoid contraction and its consequences.

**Actinoids** - Electronic configuration, oxidation states and comparison with lanthanoids.

**Unit IX Coordination Compounds****18 Periods**

Coordination compounds - Introduction, ligands, coordination number, colour, magnetic properties and shapes, IUPAC nomenclature of mononuclear coordination compounds. Bonding, Werner's theory, VBT, and CFT; structure and stereoisomerism, the importance of coordination compounds (in qualitative analysis, extraction of metals and biological system).

**Unit X Haloalkanes and Haloarenes****15 Periods**

**Haloalkanes:** Nomenclature, nature of C—X bond, physical and chemical properties, optical rotation mechanism of substitution reactions.

**Haloarenes:** Nature of C—X bond, substitution reactions (Directive influence of halogen in monosubstituted compounds only). Uses and environmental effects of - dichloromethane, trichloromethane, tetrachloromethane, iodoform, freons, **DDT**.

**Unit X Alcohols, Phenols and Ethers****14 Periods**

**Alcohols:** Nomenclature, methods of preparation, physical and chemical properties (of primary alcohols only), identification of primary, secondary and tertiary alcohols, mechanism of dehydration, uses with special reference to methanol and ethanol.

**Phenols:** Nomenclature, methods of preparation, physical and chemical properties, acidic nature of phenol, electrophilic substitution reactions, uses of phenols.

**Ethers:** Nomenclature, methods of preparation, physical and chemical properties, uses.



**Unit XII Aldehydes, Ketones and Carboxylic Acids** **15 Periods**

**Aldehydes and Ketones:** Nomenclature, nature of carbonyl group, methods of preparation, physical and chemical properties, mechanism of nucleophilic addition, reactivity of alpha hydrogen in aldehydes, uses.

**Carboxylic Acids:** Nomenclature, acidic nature, methods of preparation, physical and chemical properties; uses.

**Unit XIII Amines** **14 Periods**

Nomenclature, classification, structure, methods of preparation, physical and chemical properties, uses, identification of primary, secondary **and** tertiary amines.

**Diazonium salts:** Preparation, chemical reactions and importance in synthetic organic chemistry.

**Unit XIV Biomolecules** **18 Periods**

**Carbohydrates** - Classification (aldoses and ketoses), monosaccharides (glucose and fructose), D-L configuration oligosaccharides (sucrose, lactose, maltose), polysaccharides (starch, cellulose, glycogen); Importance of carbohydrates,

**Proteins** -Elementary idea of - amino acids, peptide bond, polypeptides, proteins, structure of proteins - primary, secondary, tertiary structure and quaternary structures (qualitative idea only), denaturation of proteins; enzymes. Hormones - Elementary idea excluding structure.

**Vitamins** - Classification and functions.

**Nucleic Acids: DNA and RNA.**

### QUESTION PAPER DESIGN CLASSES-XII (2022-23)

S.No.	Domains	Marks	%
1	<b>Remembering and Understanding:</b> Exhibit memory of previously learned material by recalling fact, terms, basic concepts and answers. Demonstrate understanding of facts and <b>idea by organizing, comparing, translating, interpreting, giving</b> descriptions and stating main idea.	28	40
2	<b>Applying:</b> Solve problems to new situations by applying <b>acquired knowledge</b> , facts, techniques and rules in a different way.	21	30
3	<b>Analyzing, Evaluating and Creating:</b> Examine and break information into parts by identifying motives or causes. Make inferences and find evidence to support generalizations. Present and defend opinions by making judgments about information, the validity of ideas or quality of work based on a set of criteria.  Compile information together in a <b>different way by combining elements in a new pattern</b> or proposing alternative solutions.	21	30

### QUESTION WISE BREAK-UP

Type of Question	Marks per Question	Total No. of Question	Total Marks
MCQ	01	18	18
VSA	02	07	14
SA	03	05	15
CASE STUDY	04	02	08
LA	05	03	15
<b>TOTAL</b>		<b>35</b>	<b>70</b>


**Choice : There will be no overall choice in the question paper.  
However, 33% internal choice be given in all the sections.**

## UNIT 2

## Solutions

### Points to Remember

- **SOLUTION:** A homogeneous mixture of two or more chemically non-reacting substances, whose composition can be varied within certain limits.
- A binary solution has two constituents one solute and one solvent.
- Solvent is the component present in largest amount and solute in smaller amount (in terms of moles).
- The solutions may be gaseous, liquid or solid depending upon the physical state of solvent.



**HENRY'S LAW:** The partial pressure of the gas in vapour phase ( $p$ ) is proportional to the mole fraction of the gas ( $\chi$ ) in the solution"

$p = K_H \chi$ ,  $K_H$  is Henry's law constant.

Higher the value of  $K_H$  at a given pressure, the lower is the solubility of the gas in the liquid.

### Applications of Henry's Law

- **Carbonated beverages:** To increase the solubility of  $\text{CO}_2$  in soft drinks and soda water, the bottle is sealed under high pressure.
- **In deep sea diving.** To avoid bends, toxic effects of high concentration of nitrogen in the blood the tanks used by scuba divers are filled with air diluted with He.
- **For climbers or people living at high altitude.** Concentration of  $\text{O}_2$  in the blood and tissues is so low that they feel weak and are unable to think properly, a disease called anoxia.

### RAOULT'S LAW

#### FOR A SOLUTION OF VOLATILE LIQUIDS:

The partial vapour pressure of each component of the solution is directly proportional to its mole fraction present in solution. If A and B are the two volatile components of solution then

$$p_A = P_A^0 \chi_A$$

$$p_B = P_B^0 \chi_B$$

Where  $p_A$  and  $p_B$  are partial vapour pressure of component 'B' respectively in solution.  $P_A^0$  and  $P_B^0$  are vapour pressure of pure components 'A' and 'B' respectively.

#### FOR A SOLUTION CONTAINING NON-VOLATILE SOLUTE:

The vapour pressure of the solution is directly proportional to the mole fraction of the solvent.

$$P_A = \chi_A$$

$$P_A = P_A^0 \chi_A$$

**Effect of adding non-volatile solute on vapour pressure of a liquid.** The vapour pressure of a liquid decrease if some non-volatile solute is dissolved in it because some molecules of the solvent on the surface are replaced by the molecules of the non-volatile solute.

- **Raoult's law becomes a special case Henry's law** in which  $K_H$  becomes equal to  $P_A^0$ , i.e., vapour pressure of pure solvent.

### TYPES OF LIQUID-LIQUID SOLUTIONS ON THE BASIS OF RAOULT'S LAW

(Let A and B be the two liquids in solution.)

#### IDEAL SOLUTIONS


- Solutions which strictly obey Raoult's law over the entire range of concentration
- The interactions between solute and solvent are similar to those in pure components.

$$P_A = P_B^0 \chi_A, P_B = P_B^0 \chi_B$$

$$\Delta H_{\text{mix}} = 0$$

$$\Delta V_{\text{mix}} = 0$$

- Benzene + toluene, hexane + heptane, bromoethane + chloroethane



Graph for an ideal solution

#### NON-IDEAL SOLUTIONS

- Solutions which do not obey Raoult's law over the entire range of concentration
- The interactions between solute and solvent are different from those of pure components.

$$P_A = P_B^0 \chi_A, P_B = P_B^0 \chi_B$$

$$\Delta H_{\text{mix}} \neq 0$$

$$\Delta V_{\text{mix}} \neq 0$$

- Two types (i) Solutions showing positive deviations from Raoult's law.  
(ii) Solutions showing negative deviations from Raoult's law.

## TYPES OF NON-IDEAL SOLUTIONS

### NON-IDEAL SOLUTIONS SHOWING POSITIVE DEVIATION FROM RAOULT'S LAW


- solute - solvent interactions are weaker than solute - solute and solvent - solvent interaction

$$p_A > p_A^0 \chi_A ; p_B > p_B^0 \chi_B$$

$$\Delta_{\text{mix}} H > 0$$

$$\Delta_{\text{mix}} V > 0$$

e.g. Acetone and ethanol, Water and ethanol, Acetone and benzene.



### NON-IDEAL SOLUTIONS SHOWING NEGATIVE DEVIATION FROM RAOULT'S LAW


- solute - solvent interactions are stronger than solute - solute and solvent - solvent interaction

$$p_A < p_A^0 \chi_A ; p_B < p_B^0 \chi_B$$

$$\Delta_{\text{mix}} H < 0$$

$$\Delta_{\text{mix}} V < 0$$

e.g. Acetone and aniline, Water and nitric acid, Water and HCl



## AZEOTROPES

Liquid mixture, having the same composition, in liquid and vapour phase and boiling like a pure liquid is called a constant boiling mixture or an azeotropic mixture or an azeotrope.

### MINIMUM BOILING AZEOTROPE

Minimum boiling azeotropes form when solutions exhibit positive deviation from Raoult's law. e.g. 95% ethanol - water mixture.

### MAXIMUM BOILING AZEOTROPE

Maximum boiling azeotropes form when solutions exhibit negative deviation from Raoult's law. e.g. 68% nitric acid - water mixture.

## COLLIGATIVE PROPERTIES

Physical properties of dilute solutions that depend upon the number of solute particles present in the solution irrespective of their nature.

**RELATIVE LOWERING IN VAPOUR PRESSURE**

$$\chi_B = \frac{P_A^0 - P_A}{P_A^0}$$

$P_A^0$  = Vapour pressure of solvent,  $P_A$  = Vapour pressure of solution

Where  $P_A^0 - P_A / P_A^0$  is relative lowering in vapour pressure.  $\chi_B$  = mole fraction of solute

$$\chi_B = \frac{n_B}{n_A + n_B}$$

For dilute solution,  $n_B \ll n_A$ , hence  $n_A$  is neglected in the denominator.

$$\frac{P_A^0 - P_A}{P_A^0} = \frac{n_B}{n_A}$$

$$\frac{P_A^0 - P_A}{P_A^0} = \frac{w_B}{M_B} \times \frac{M_A}{w_A}$$

$w_B$  = mass of solute,  $M_B$  = molar mass of solute

$w_A$  = mass of solvent,  $M_B$  = molar mass of solvent

**ELEVATION IN BOILING POINT ( $\Delta T_b$ )**


$$\Delta T_b \propto m, \Delta T_b = k_b m; m = \text{molality}$$

$k_b$  = molal elevation constant / Ebullioscopic constant. It is the elevation in boiling point when the molality of solution is unity. SI unit :  $K \text{ kg mol}^{-1}$

Elevation in boiling point and Molar mass of solute  $M_B = k_b \cdot 1000 w_B / \Delta T_b W_A$

$M_B$  = Molar mass of solute,  $w_B$  = mass of solute,

$w_A$  = mass of solvent (in gram)

**COLLIGATIVE PROPERTIES****DEPRESSION IN FREEZING POINT ( $\Delta T_f$ )**

$$\Delta T_f \propto m, \Delta T_f = K_f m$$

$K_f$  = molal depression constant / Cryoscopic constant. It is the depression in freezing point when the molality of solution is unity. SI unit :  $K \text{ kg mol}^{-1}$


**Depression in freezing point and Molar mass of solute**

$$M_B = k_f \cdot 1000 w_B / \Delta T_f W_A$$

$M_B$  = Molar mass of solute,  $w_B$  = mass of solute,

$W_A$  = mass of solvent (in gram)

**APPLICATION** : Ethylene glycol is used as antifreeze (causes depression in freezing point) in automobiles.

**OSMOTIC PRESSURE ( $\pi$ )**

The excess pressure that must be applied to a solution side to prevent osmosis i.e. to stop the passage of solvent molecules into it through semi-permeable membrane is called osmotic pressure.

$$\pi \propto C, \pi \propto T, \pi = CRT,$$

$C$  = Molarity of solution,  $C = n_B / V$ ,  $V$  = volume of solution (L),  $n_B$  = no. of moles of solute

$$\pi V = n_B RT$$

$$\pi = w_B RT / M_B V$$

$$R = 0.0821 \text{ L atm K}^{-1} \text{ mol}^{-1}; T = \text{Temperature in Kelvin}$$

**ISOTONIC SOLUTIONS**

Two solutions having same osmotic pressure and same concentration are called isotonic solutions.

**Hypertonic solution** have higher osmotic pressure and **Hypotonic solution** have lower osmotic pressure than the other solution. 0.91% of sodium chloride is isotonic with fluid present inside human red blood cells.

**REVERSE OSMOSIS**- If a pressure higher than osmotic pressure is applied on the solution the solvent will flow from the solution into the pure solvent through the semi permeable embrance. It is used in the desalination of sea water.

## OBJECTIVE TYPE QUESTIONS

### I MULTIPLE CHOICE QUESTIONS

1. The molarity of 98%  $\text{H}_2\text{SO}_4$  (density = 1.8 g/mL) by weight is:
 

(a) 6 M	(b) 18 M
(c) 10 M	(d) 4 M
2. Which of the following does not show positive deviation from Raoult's law?
 

(a) benzene + chloroform	(b) benzene + acetone
(c) benzene + ethanol	(d) benzene + $\text{CCl}_4$
3. Which solution will have least vapour pressure?
 

(a) 1 M glucose	(b) 2 M glucose
(c) 3 M glucose	(d) 4 M glucose
4. Which condition is not satisfied by an ideal solution?
 

(a) $\Delta H_{\text{mix}} = 0$	(b) $\Delta V_{\text{mix}} = 0$
(c) $\Delta P_{\text{mix}} = 0$	(d) $\Delta S_{\text{mix}} = 0$
5. Azeotrope mixture are:
 

(a) mixture of two solids	(b) those will boil at different temperature
(c) those which can be fractionally distilled	(d) constant boiling mixtures
6. Which is temperature independent term?
 

(a) w/W%	(b) v/V%
(c) w/V%	(d) Molarity
7. Solute when dissolve in water
 


(a) increases the vapour pressure of water	(b) increases the boiling point of water
(c) decrease the freezing point of water	(d) All of the above
8. The plant cell will shrink when placed in:
 

(a) water	(b) a hypotonic solution
(c) a hypertonic solution	(d) an isotonic solution



9. Two aqueous solutions  $S_1$  and  $S_2$  are separated by a semi-permeable membrane.  $S_2$  has lower vapour pressure than  $S_1$  of a non-volatile solute, Then
- more solvent will flow from  $S_1$  to  $S_2$
  - more solvent will flow from  $S_2$  to  $S_1$
  - solvent from  $S_1$  and  $S_2$  will flow at equal rates
  - no flow will take place
10. Temperature dependent concentration term is :
- M
  - m
  - $x$
  - All of these
11. Which of the following solutions would have the highest osmotic pressure:
- M/10 NaCl
  - M/10 Urea
  - M/10  $\text{BaCl}_2$
  - M/10 Glucose
12. 0.5 M aqueous solution of glucose is isotonic with:
- 0.5 M KCl solution
  - 0.5 M  $\text{CaCl}_2$  solution
  - 0.5 M Urea solution
  - 1 M solution of sucrose
13. Which of the following is true for Henry's constant ?
- It decreases with temperature
  - It increases with temperature
  - Independent on temperature
  - It do not depend on nature of gases.
14. Which one is the best colligative property for determination of molecular mass of polymer?
- osmotic pressure
  - elevation in boiling point
  - depression in freezing point
  - osmosis
15. An azeotropic solution of two liquids has boiling point lower than either of them when it
- shows negative deviation from Raoult's Law
  - shows no deviation from Raoult's Law
  - shows positive deviation from Raoult's Law
  - is saturated
16. Henry's law constant  $K_H$  of  $\text{CO}_2$  in water at  $25^\circ\text{C}$  is  $3 \times 10^{-2} \text{ mol/L atm}^{-1}$ . Calculate the mass of  $\text{CO}_2$  present in 100 L of soft drink bottled with a partial pressure of  $\text{CO}_2$  of 4 atm at the same temperature.
- 5.28 g
  - 12.0 g
  - 428 g
  - 528 g

17. If osmotic pressure of 1 M urea is  $\pi$ , what will be the osmotic pressure for 2 M urea?
- (a)  $\pi$  (b)  $0.1\pi$   
(c)  $2\pi$  (d)  $0.2\pi$
18. The most likely an ideal solution is:
- (a)  $\text{NaCl-H}_2\text{O}$  (b)  $\text{C}_2\text{H}_5\text{OH-C}_6\text{H}_6$   
(c)  $\text{C}_7\text{H}_{16}\text{-H}_2\text{O}$  (d)  $\text{C}_7\text{H}_{16}\text{-C}_8\text{H}_{18}$
19.  $\Delta H_{\text{mix}}$  for solution of  $\text{CHCl}_3$  and  $\text{CH}_3\text{COCH}_3$  is.
- (a) +ve (b) 0  
(c) -ve (d) None of these
20. The solutions A, B, C and D are respectively 0.1 M glucose, 0.05 M NaCl, 0.05 M  $\text{BaCl}_2$  and 0.1 M  $\text{AlCl}_3$ . which one of the following pairs is isotonic?
- (a) A & C (b) B & C  
(c) C & D (d) A & B
21. Which one of the following pairs will form an ideal solution?
- (a) Chloroform and acetone (b) Ethanol and acetone  
(c) n-hexane and n-heptane (d) Phenol and aniline
22. An azeotropic solution of two liquids has a boiling point lower than either of the two when it ?
- (a) shows a positive deviation from Raoult's law  
(b) shows a negative deviation from Raoult's law.  
(c) shows no deviation from Raoult's law.  
(d) is saturated.
23. Which of the following formula represents Raoult's law for a solution containing non-volatile solute?
- (a)  $P_{\text{solute}} = P^0_{\text{solute}} \cdot X_{\text{solute}}$   
(b)  $P = K_{\text{H},x}$   
(c)  $P_{\text{Total}} = P_{\text{solvent}}$   
(d)  $P_{\text{solute}} - P^0_{\text{solvent}} \cdot x_{\text{solvent}}$
24. On mixing 20mL of acetone with 30 mL of chloroform, the total volume of the solution is
- (a) <50mL (b) =50mL  
(c) >50mL (d) =10mL

25. Elevation of boiling point is inversely proportional to  
 (a) molal elevation constant ( $K_b$ ) (b) molality ( $m$ )  
 (c) molar mass of solute ( $M$ ) (d) weight of solute ( $w$ )
26. An unknown gas 'X' is dissolved in water at 2.5 bar pressure and has mole fraction 0.04 in solution. The mole fraction of 'X' gas when the pressure of gas is doubled at the same temperature is  
 (a) 0.08 (b) 0.04  
 (c) 0.02 (d) 0.92
27. The boiling point of a 0.2 m solution of a non-electrolyte in water is ( $K_b$  for water =  $0.52 \text{ kg mol}^{-1}$ )  
 (a)  $100^\circ\text{C}$  (b)  $100.52^\circ\text{C}$   
 (c)  $100.104^\circ\text{C}$  (d)  $100.26^\circ\text{C}$
28. In the following diagram point, 'X' represents
- 
- (a) Boiling point of solution (b) Freezing point of solvent  
 (c) Boiling point of solvent (d) Freezing point of solution

## II FILL IN THE BLANKS

- The sum of mole fractions of all the components in a three component system is equal to.....
- A Solution which distils without change in composition is called .....
- Desalination of sea water is based on the phenomenon of.....
- Relative lowering in vapour pressure is equal to the mole fraction of .....
- The evaporation of aqueous solution of glucose causes its molarity to .....
- The boiling point of sea water at 1 atm pressure is that of distilled water.
- The ratio of observed value of colligative property to the calculated value of colligative property is called .....
- The most suitable colligative property to measure molecular mass of polymers is.....
- People taking a lot of salt develop swelling or puffiness of their tissues. This disease is called.....
- If observed molar mass of a solute is more than calculated molar mass, then the solute undergoes..... in solution.

### III ASSERTION REASON TYPE QUESTIONS

- (a) Both assertion and reason are correct statements, and reason is the correct explanation of the assertion.
  - (b) Both assertion and reason are correct statements, but reason is not the correct explanation of the assertion.
  - (c) Assertion is correct, but reason is wrong statement
  - (d) Assertion is wrong, but reason is correct statement
1. **Assertion:** Molality is a better method to express concentration than molarity.  
**Reason:** Molality is defined in terms of mass of solvent and not volume of solution .
  2. **Assertion:** Soda bottles are sealed under high pressure.  
**Reason:** High pressure increases the solubility of carbon dioxide gas in solution .
  3. **Assertion:** Non-ideal solutions always form azeotropes.  
**Reason:** Boiling point of an azeotrope may be lower or higher than boiling points of both components.
  4. **Assertion:** Benzene and hexane form an ideal solution.  
**Reason :** Both benzene and hexane are hydrocarbons.
  5. **Assertion:** 1 molar NaCl solution has higher boiling point than one molar urea.  
**Reason:** NaCl dissociates into ions in solution .
  6. **Assertion:** Two solutions having same osmotic pressures will also have same vapour pressures.  
**Reason:** Lowering of vapour pressure is not a colligative property .
  7. **Assertion :** Helium is mixed with nitrogen and oxygen in diving cylinders  
**Reason:** Helium has comparatively low solubility in blood.
  8. **Assertion:** NaCl or  $\text{CaCl}_2$  is used to clear snow on roads in the hills.  
**Reason:** The salts depress the freezing point of water.
  9. **Assertion:** Molar mass of acetic acid in benzene calculated using colligative property is almost double the actual value.  
**Reason:** Acetic acid dimerises in solution.
  10. **Assertion:** Vapour pressure of a solution is more than that of the pure solvent.  
**Reason:** The solute particles occupy certain area of the surface of the solution which reduces the amount of vapour.

**IV ONE WORD ANSWER TYPE QUESTIONS**

1. Which of the following is a dimension less quantity : molarity, molality or mole fraction?
2. Liquid 'Y' has higher vapour pressure than liquid 'X' . Which of them will have higher boiling point?
3.  $N_2$  and  $O_2$  gases have  $K_H$  values 76.48 kbar and 34.86 kbar respectively at 293 K temperature. Which one of these will have more solubility in water?
4. Name for  $k_b$  is \_\_\_\_\_.
5. Mention the unit of ebullioscopic constant (molal boiling point elevation constant).
6. What type of deviation from Raoult's law is exhibited the solution forming minimum boiling azeotrope?
7. For reverse osmosis to take place external pressure applied must be lesser than or greater than osmotic pressure?
8. Name the law which can explain the solubility of gases in liquids at different pressures.
9. Out of molarity and molality which is preferred for expressing the concentration of solution?
10. A decrease in temperature is observed on mixing ethanol and acetone. What type of deviation from Raoult's law is this?
11. What is the sum of the mole fractions of all the components in a three component system?
12.  $10\text{ cm}^3$  of a liquid A was mixed with  $10\text{ cm}^3$  of liquid B. The volume of the resulting solution was found to be  $19.9\text{ cm}^3$ . What do you conclude?
13. Name the disease caused by low concentration of oxygen in the blood and tissues of people living at high altitude.
14. Mention a large scale use of reverse osmosis.
15. Under which condition van't Hoff factor is less than one.

**VERY SHORT ANSWER TYPE QUESTIONS (1 Mark Questions)****Q.1. What is van't Hoff factor ?****Ans.** It is the ratio of normal molecular mass to observed molecular mass. It is denoted as  $i$ .

$$i = \frac{\text{normal molecular mass}}{\text{observed molecular mass}}$$

$$i = \frac{\text{no. of particles after association or dissociation}}{\text{no. of particles before dissociation or association}}$$

**Q.2. What is the van't Hoff factor in  $K_4[Fe(CN)_6]$  and  $BaCl_2$  ?****Ans.** 5 and 3**Q.3. Why the molecular mass becomes abnormal ?****Ans.** Due to association or dissociation of solute in given solvent.**Q.4. What role does the molecular interaction play in the solution of alcohol and water ?****Ans.** Positive deviation from ideal behaviour.**Q.5. What is van't Hoff factor ? How is it related with :****(a) degree of dissociation****(b) degree of association**

$$\text{Ans. (a) } \alpha = \frac{i - 1}{n - 1}$$

$$\text{(b) } \alpha = \frac{i - 1}{1/n - 1}$$

**Q.6. Why NaCl is used to clear snow from roads ?****Ans.** It lowers freezing point of water.**Q.7. Why the boiling point of solution is higher than pure liquid ?****Ans.** Due to lowering in vapour pressure.**Q.8. Henry law constant for two gases are 21.5 and 49.5 atm, which gas is more soluble ?****Ans.**  $K_H$  is inversely proportional to solubility.**Q.9. Define azeotrope. Give an example of maximum boiling azeotrope.****Hint:** Refer "Points to remember"**Q.10. Calculate the volume of 75% of  $H_2SO_4$  by weight ( $d = 1.8 \text{ gm/mL}$ ) required to prepare 1 L of 0.2 M solution.**

$$\text{Hint: } M_1 = \frac{\text{Mass \%} \times d \times 10}{98}$$

$$M_1 V_1 = M_2 V_2$$

$$= 14.5 \text{ mL}$$

**Q.11. Why water cannot be completely separated from aqueous solution of ethyl alcohol ?****Ans.** Due to formation of azeotrope at (95.4%).**Q.12. Why anhydrous salts like NaCl or  $CaCl_2$  are used to clear snow from roads on hills ?****Hint :** They depress freezing point of water.

**Q.13. What is the effect on boiling and freezing point of a solution on addition of NaCl?**

*Hint :* Boiling point increases and freezing point decreases.

**Q.14. Why osmotic pressure is considered as colligative property ?**

*Hint :* It depends upon number of moles of solute present in solution.

**Q.15. Liquid A and B on mixing produce a warm solution. Which type of deviation does this solution show ?**

*Hint :* — ve deviations

**Q.16. Give an example of a compound in which hydrogen bonding results in the formation of a dimer.**

*Hint :* Carboxylic acids or other example

**Q.17. What role does the molecular interaction play in solution containing chloroform and acetone ?**

*Hint :* H-bonding formed, results in negative deviation from Raoult's law.

**Q.18. What is meant by 5%  $\text{Na}_2\text{CO}_3$  solution (w/W)?**

**Ans.** 5% w/W means 5g  $\text{Na}_2\text{CO}_3$  dissolves in 100 g solution.

**Q.19. What will be the mole fraction of  $\text{C}_2\text{H}_5\text{OH}$  in aqueous solution of  $\text{C}_2\text{H}_5\text{OH}$  when solution contain equal number of moles of water and  $\text{C}_2\text{H}_5\text{OH}$ ?**

**Ans.** Solution is equimolar, it means mole fraction of each component is 0.5.

**Q.20. If at the same temperature, hydrogen is more soluble in water than helium, which of them will have a higher value of  $K_H$ ?**

**Ans.** As  $\text{H}_2$  is more soluble than helium, so  $\text{H}_2$  will have lower value of  $K_H$  than that of helium.

**Q.21. State the formula relating to the pressure of a gas with its mole fraction in a liquid solution in contact with it**

*Hint:*  $p = K_H \cdot x$

**Q.22. If  $K_f$  for water is  $1.86 \text{ K kg mol}^{-1}$ , what is the boiling point of 0.01 molal aqueous solution of a substance which undergoes no association or dissociation?**

**Ans.**  $\Delta T_b = i K_b \cdot m$

$$\begin{aligned}\Delta T_b &= 1.1.86 \times 0.01 = 0.0186 \\ &= 100.0186^\circ\text{C}\end{aligned}$$

**Q.23. Why does sodium chloride solution freeze at a lower temperature than water?**

*Hint:* NaCl being non-volatile solute, decreases the vapour pressure and therefore decreases the freezing point.

**Q.24. Out of 0.1 molal solution of glucose and NaCl, which one will have a higher boiling point and why?**

**Ans.** 0.1 m NaCl solution will have higher boiling point because it dissociates in the solution. As a result, number of moles of the solute in solvent is higher in case of NaCl than glucose.

**Q.25. Ionic compounds are soluble in water but they are insoluble in organic solvents.**

**Give reason.**

*Hint:* "Like dissolves like"

### SHORT ANSWER TYPE QUESTIONS (2 or 3 Marks Questions)

**Q.1. State Henry's I law. What is the significance of  $K_H$  ?**

**Ans. Henry's Law:** It states that "the partial pressure of the gas in vapour phase (p) is directly proportional to the mole fraction of the gas (x) in the solution", and is expressed as :  $p=K_H \cdot x$  where,  $K_H$  is the Henry's Law constant

**Significance of  $K_H$  :** Higher the value of Henry's law constant  $K_H$ , the lower is the solubility of the gas in the liquid.

**Q.2. How is that measurement of osmotic pressure is more widely used for determining molar masses of macromolecules than the elevation in boiling point or depression in freezing point of their solutions?**

**Ans :** The osmotic pressure method has the advantage over elevation in boiling point or depression in freezing point for determining molar masses of macromolecules because

1. Osmotic pressure is measured at the room temperature and the molarity of solution is used instead of molality.
2. Compared to other colligative properties, its magnitude is large even for very dilute solutions.

**Q.3. Equal moles of liquid P and Q are mixed. What is the ratio of their moles in the vapour phase? Given that  $P_p^\circ = 2 \times P_q^\circ$ .**

Hint: Since equal moles of P and Q are mixed

$$\Rightarrow \text{Mole fraction of P} = \text{Mole fraction of Q} = x = 1/2$$

$$P_p = P_p^\circ \times 1/2 = 2 \times P_q^\circ \times 1/2 = P_q^\circ$$

$$P_q = P_q^\circ \times 1/2 = P_q^\circ/2$$

- In vapour phase, let the total pressure be P

$$\Rightarrow y_1 = P_p / P = P_q^\circ / P$$

$$\Rightarrow y_2 = P_q / P = (P_q^\circ/2) / P$$

$$\Rightarrow y_1/y_2 = 2/1$$

- Ratio of moles of P and Q in vapour phase =  $y_1 : y_2 = 2 : 1$

$$\Rightarrow \mathbf{P:Q=2:1}$$

**Q.4. On mixing liquid X and Y, volume of the resulting solution decreases. What type of deviation from Raoult's law is shown by the resulting solution ? What change in temperature would you observe after mixing liquids X and Y ?**

Hint: Negative; Increase in temp.

**Q.5. Explain the significance of Henry's constant ( $K_H$ ). At the same temperature, hydrogen is more soluble in water than helium. Which of them will have higher value of  $K_H$  and why ?**

Hint: Significance of  $K_H$  Higher the value of Henry's law constant  $K_H$ , the lower is the solubility of the gas in the liquid ; He has higher value of  $K_H$



**Q.6. How many grams of KCl should be added to 1 kg of water to lower its freezing point to  $-8.0^{\circ}\text{C}$  ? ( $K_f = 1.86 \text{ K kg/mol}$ )**

**Ans.** Since KCl dissociate in water completely,  $i = 2$ .

$$\begin{aligned}\Delta T_f &= iK_f \cdot m \\ m &= \Delta T_f / i K_f \\ &= \frac{8}{2 \times 1.86} \\ m &= 2.15 \text{ mol/kg} \\ \text{Grams of KCl} &= 2.15 \times 74. = 160.2 \text{ g/kg}\end{aligned}$$

**Q.7. Why is freezing point depression of 0.1 M sodium chloride solution nearly twice that of 0.1 M glucose solution?**

**Hint:** Colligative properties  $\propto$  number of particles.

**NaCl is a strong electrolyte** and gives two particles on dissociation, but glucose being non-electrolyte does not dissociate and remains as a single particle

**Q.8. a) Why is an increase in temperature observed on mixing chloroform and acetone?**

**b) Why does sodium chloride solution freeze at a lower temperature than water?**

**Ans: a)** The bonds between chloroform molecules and molecules of acetone are dipole-dipole interactions but on mixing, the chloroform and acetone molecules, they start forming hydrogen bonds which are stronger bonds resulting in the release of energy. This gives rise to an increase in temperature.

**b)** When a non-volatile solute is dissolved in a solvent, the vapour pressure decreases. As a result, the solvent freezes at a lower temperature.

**Q.9. Define reverse osmosis. Write its one use.**

**Hint:** If the pressure applied on the solution is greater than the osmotic pressure then the solvent molecules start to move from solution into solvent through a semipermeable membrane this process called the reverse osmosis. ; Desalination of water.

**Q.10. Why does an azeotropic mixture distill without any change in composition ?**

**Hint :** It has same composition of components in liquid and vapour phase.

**Q.11. Under what condition Van't Hoff factor is :**

**(a) equal to 1 ? (b) less than 1 ? (c) more than 1 ?**

**Hint:** (a) When the solute neither associates nor dissociates in solution,  $i$  is equal to 1.

(b) When the solute undergoes association in solution,  $i$  is less than 1.

(c) When the solute undergoes dissociation in solution,  $i$  is more than 1.

**Q.12.** An aqueous solution of 2% non-volatile exerts a pressure of 1.004 Bar at the normal boiling point of the solvent. What is the molar mass of the solute ?

Hint:

$$\frac{P_A^0 - P_A}{P_A^0} = \frac{W_B \times m_A}{m_B \times W_A}$$

$$\frac{1.013 - 1.004}{1.013} = \frac{2 \times 18}{m_B \times 98}$$

$$m_B = 41.35 \text{ gm/mol}$$

**Q.13.** Why is it advised to add ethylene glycol to water in a car radiator in hill station ?

Hint: Anti-freeze.

**Q.14.** Calculate the molarity of pure water ( $d = 1 \text{ g mL}^{-1}$ ).

Ans. Density of water =  $1 \text{ g mL}^{-1}$

Mass of 1000 mL of water =  $V \times d$

$$= 1000 \text{ mL} \times 1 \text{ gm}$$

$$= 1000 \text{ g}$$

$$\text{Moles of water} = \frac{1000}{18} = 55.55 \text{ mol}$$

Now, mole of  $\text{H}_2\text{O}$  present in 1000 mL or 1 L of water.

So, molarity = 55.55M

**Q.15.** Define Henry's law. Give their two application.

Hint: Refer "Points to remember"

**Q.16.** The dissolution of ammonium chloride in water is endothermic process. What is the effect of temperature on its solubility ?

Ans. Since dissolution of  $\text{NH}_4\text{Cl}$  in water is endothermic process, its solubility increases with rise in temperature (*i.e.*, Le-Chatelier process).

**Q.17.** Two liquids A and B boil at  $145^\circ\text{C}$  and  $190^\circ\text{C}$  respectively. Which of them has higher vapour pressure at  $80^\circ\text{C}$  ?

Ans. Lower the boiling point more volatile is the respective compound. Therefore, liquid A will have higher vapour pressure at  $80^\circ\text{C}$ .

**Q.18.** Why is liquid ammonia bottle first cooled in ice before opening it ?

Ans. At room temperature, the vapour pressure of liquid ammonia is very high. On cooling vapour pressure decreases, therefore the liquid ammonia will not splash out.


**Q.19** Determine the amount of  $\text{CaCl}_2$  dissolved in 2.5L at  $27^\circ\text{C}$  such that its osmotic pressure is 0.75 atm at  $27^\circ\text{C}$ . ( $i$  for  $\text{CaCl}_2 = 2.47$ )

Ans. For  $\text{CaCl}_2$ ,

$$\begin{aligned} i &= 2.47 \\ \pi &= iCRT \\ &= i \frac{n_B}{V} \times RT \\ 0.75 &= \frac{2.47 \times n_B \times 0.082 \times 300}{2.5} \\ n &= \frac{0.75 \times 2.5}{2.47 \times 0.082 \times 300} \\ n_B &= 0.0308 \text{ mol} \\ \text{Amount of } \text{CaCl}_2 &= 0.0308 \text{ mol} \times 111 \text{ g mol}^{-1} \\ &= 3.418 \text{ g} \end{aligned}$$

**Q.20.** Determine the osmotic pressure of a solution prepared by dissolving 25 mg of  $\text{K}_2\text{SO}_4$  in 2 litre of water at  $25^\circ\text{C}$  assuming that it is completely dissociated.

Ans. If  $\text{K}_2\text{SO}_4$  is completely dissociated,



$$i = 3$$

$$\text{Mol mass of } \text{K}_2\text{SO}_4 = 2 \times 39 + 32 + 4 \times 16 = 174 \text{ g mol}^{-1}$$

$$\begin{aligned} \pi &= iCRT \\ &= \frac{W_B \times RT}{M_B \times V} \\ &= \frac{3 \times 25 \times 10^{-3} \times 0.082 \times 298}{174 \times 2.0} \\ &= 5.27 \times 10^{-3} \text{ atm} \end{aligned}$$

**Q.21.** If the solubility product of  $\text{CuS}$  is  $6 \times 10^{-16}$ , calculate the maximum molarity of  $\text{CuS}$  in aqueous solution.

Ans.  $K_{sp}$  of  $\text{CuS} = 6 \times 10^{-16}$

If  $S$  is the solubility, then



$$[\text{Cu}^{2+}] = S, [\text{S}^{2-}] = S$$

$$K_{sp} = [\text{Cu}^{2+}][\text{S}^{2-}] = S \times S = S^2$$

$$\begin{aligned} \text{Solubility } S &= \sqrt{K_{sp}} = \sqrt{6 \times 10^{-16}} \\ &= 2.45 \times 10^{-8} \text{ M} \end{aligned}$$

$$\text{Highest molarity} = 2.45 \times 10^{-8} \text{ M}$$

**Q.22 Suggest the most important type of intermolecular attractive interaction in the following pairs:**

- (a) n-hexane and n-octane (b)  $I_2$  and  $CCl_4$   
 (c)  $NaClO_4$  and water

**Ans.** (a) Vander Waals interaction  
 (b) Vander Waals interaction  
 (c) Ion-dipole interaction

**23. The vapour pressure of water is 12.3 Kpa at 300K. Calculate vapour pressure of 1 molal solution of a non-volatile solute in it.**

**Ans.** Mole fraction of solute =

$$\frac{1}{1 + \frac{1000}{18}} = 0.0177$$

$$\frac{P^0 - P_A}{P^0} = 0.0177$$

$$\frac{12.3 - P_A}{12.3} = 0.0177$$

$$P_A = 12.08 \text{ Kpa}$$

**Q.24. 6.90M solution of KOH in water contains 30% by mass of KOH. Calculate the density of the KOH solution. (Molar mass of KOH = 56 g mol<sup>-1</sup>)**

**Ans.** Mass of KOH = 30 g

$$M = \frac{n_B}{V(\text{mL})} \times 1000$$

$$= \frac{W_B}{M_B \times V(\text{mL})} \times 1000 = \frac{30}{56 \times V} \times 1000$$

$$6.90 = \frac{30 \times 1000}{56 \times V}$$

$$V = \frac{30 \times 1000}{56 \times 6.90} = 81.43 \text{ mL}$$

$$D = \frac{M}{V}$$

$$= \frac{1000}{81.43} = 1.28 \text{ g mL}^{-1}$$

**Q.25.** An anti-freeze solution is prepared from 222.6 g of ethylene glycol  $C_2H_4(OH)_2$  and 200 g of water. Calculate the molality of the solution. If the density of this solution be  $1.072 \text{ g L}^{-1}$ , what will be the molarity of the solution ?

**Ans.**  $M_B$  of  $C_2H_4(OH)_2 = 62 \text{ g mol}^{-1}$

$$\text{Molality} = \frac{n_B}{W_A} \times 1000 = \frac{W_B}{M_B \times W_B} \times 1000 = \frac{222.6 \times 1000}{56 \times 200}$$

$$= 17.95 \text{ m}$$

$$\text{Density} = \frac{\text{Mass}}{\text{Volume}}$$

$$\text{So, Volume} = \frac{\text{Mass}}{\text{Density}} = \frac{422.6}{1.072} = 394.22 \text{ ml}$$

$$M = \frac{n_B}{V} \times 1000$$

$$= \frac{222.6}{394.22 \times 62} \times 1000 = 9.11 \text{ M}$$

**Q.26.** What would be the molar mass of compound if 6.21 g of it is dissolved in 24.0 g of  $CHCl_3$  from a solution that has a boiling point of  $68.04^\circ\text{C}$ . The boiling point of pure chloroform is  $61.7^\circ\text{C}$  and the boiling point elevation constant

$K_b$  for chloroform is  $3.63^\circ\text{C/m}$ .

**Ans.** Elevation in boiling point  $\Delta T_b = 68.04 - 61.7 = 6.31^\circ\text{C}$

Mass of substance  $W_B = 6.21 \text{ g}$

Mass of  $CHCl_3$   $W_A = 24.0 \text{ g}$

$K_B = 3.63^\circ\text{C/m}$

$$M_B = \frac{K_b \times W_B \times 1000}{\Delta T_b \times W_A} = \frac{3.63 \times 6.21 \times 1000}{6.34 \times 24}$$

$$= 148.15 \text{ g mol}^{-1}$$

**Q.27** A solution of glycerol ( $C_3H_8O_3$ ) in water was prepared by dissolving some glycerol in 500 g of water. This solution has a boiling point of  $100.42^\circ\text{C}$  while pure water boils at  $100^\circ\text{C}$ . What mass of glycerol was dissolved to make the solution ? ( $K_b = 0.512 \text{ K kg mol}^{-1}$ )

**Ans.** Given  $w_1 = 500\text{g}$   
 Boiling point of solution ( $T_b$ ) =  $100.42^\circ\text{C}$   
 $K_b(\text{H}_2\text{O}) = 0.512 \text{ K kg mol}^{-1}$   
 $M_2(C_3H_8O_3) = 3 \times 12 + 8 \times 1 + 3 \times 16 = 92 \text{ g mol}^{-1}$   
 $\Delta T_b = T_b - T_b^0 = 373.42 - 373 \text{ K} = 0.42 \text{ K}$

As we know 
$$\Delta T_b = \frac{K_b \cdot w_2 \times 1000}{M_2 \times w_1} \Rightarrow w_2 = \frac{T_b \times M_2 \times w_1}{K_b \times 100}$$

$$w_2 = \frac{0.42 \text{ K} \times 92 \text{ g mol}^{-1} \times 500 \text{ g}}{0.512 \text{ K kg mol}^{-1} \times 1000 \text{ g kg}^{-1}} = 37.73 \text{ g}$$

$= 17.95 \text{ m}$

**Q.28.** 18 g of glucose ( $C_6H_{12}O_6$ ) (molar mass =  $180 \text{ g mol}^{-1}$ ) is dissolved in 1 kg of water in a saucepan. At what temperature will this solution boil ? for water =  $0.52 \text{ K kg pure water} = 373.1 \text{ K}$ )

**Ans.** According to question,

$$M_1 = 18 \text{ g mol}^{-1}$$

$$M_1 = 18 \text{ g mol}^{-1}$$

$$w_1 = 1 \text{ kg} = 1000 \text{ g}$$

$$w_2 = 18 \text{ g } K_b = 0.52 \text{ K kg mol}^{-1}$$

$$\text{We know that } \Delta T_b = K_b \cdot \frac{w_2}{M_2} \times \frac{1000}{w_1}$$

$$\text{or } \Delta T_b = 0.52 \text{ K kg mol}^{-1} \times \frac{18 \text{ g}}{180 \text{ g mol}^{-1}} \times \frac{1000 \text{ kg}^{-1}}{1000 \text{ g}} = 0.052 \text{ K}$$

Since water boils at  $373.15 \text{ K}$  at  $1.013 \text{ bar}$  pressure, therefore, the boiling point of solution will be  $373.15 + 0.052 = 373.202 \text{ K}$

**LONG ANSWER TYPE QUESTIONS (5 Marks )**

**Q.1 (a) Define Raoult's law of binary solution containing non-volatile solute in it.**

**(b) On dissolving 3.24 g of sulphur in 40 g of benzene, boiling point of solution was higher than that of benzene by 0.81 K ( $K_b = 2.53 \text{ K kg mol}^{-1}$ ). What is molecular formula of sulphur ? (Atomic mass  $s = 32 \text{ g mol}^{-1}$ )**

Ans.(a) At a given temperature, the vapour pressure of a solution containing non-volatile solute is directly proportional to the mole fraction of the solvent.

$$\begin{aligned} \text{(b)} \quad M_B &= \frac{K_b \cdot w_2 \times 1000}{\Delta T_b \times W_A} = \frac{2.53 \times 3.24 \times 10^{-3}}{0.81 \times 40} \\ &= 253 \text{ g mol}^{-1} \end{aligned}$$

Let the molecular formula of sulphur =  $S_x$

Atomic mass of sulphur = 32

Molecular mass =  $32 \times x$

$$32x = 253$$

$$x = 7.91 \approx 8$$

Molecular formula of sulphur =  $S_8$

**Q.1(a) Outer shells of two eggs are removed. One of the egg is placed in pure water and the other is placed in saturated solution of NaCl. What will be observed and why ?**

**(b) A solution prepared by dissolving 8.95 mg of a gene fragment in 35.0 ml of water has an osmotic pressure of 0.335 ton at 25°C. Assuming the gene fragment is a non-electrolyte, determine the molar mass.**

Ans. (a) In pure water the egg swells and in saturated solution of NaCl it will shrink.

$$\begin{aligned} \text{(b)} \quad \text{Mass of gene fragment} &= 8.95 \text{ mg} \\ &= 8.95 \times 10^{-3} \text{ g} \\ \text{Volume of water} &= 35.0 \text{ ml} = 35 \times 10^{-3} \text{ L} \\ \pi &= 0.335 \text{ ton} = 0.335/760 \text{ atm} \\ \text{Temp} &= 25 + 273 = 298 \text{ K} \\ \pi &= \frac{W_B RT}{M_B \times V} \\ \frac{0.335}{760} &= \frac{8.95 \times 10^{-3} \times 0.0821 \times 298}{M_B \times 35 \times 10^{-3}} \\ M_B &= 141933 \text{ g mol}^{-1} \end{aligned}$$

**Q.3 (a) Define van't Hoff factor.**

**(b) Calculate the freezing point depression expected for 0.0711M aqueous solution of  $\text{Na}_2\text{SO}_4$ . If this solution actually freezes at  $-0.320^\circ\text{C}$ , what would be the value of van't Hoff factor ? ( $K_f = 1.86^\circ\text{C mol}^{-1}$ )**

Ans.(a) **van't Hoff factor** : It is the ratio of the normal molar mass to the observed molar mass of the solute.

$$(b) \quad \Delta T_f = K_f \times M$$

$$\Delta T_f = 1.86 \times 0.0711 = 0.132$$

$$\text{Observed freezing point} = 0 - (-0.320) = 0.320^\circ\text{C}$$

$$\begin{aligned} i &= \frac{\text{Observed Freezing point}}{\text{Calculate freezing point}} \\ &= \frac{0.320}{0.132} = 2.42 \end{aligned}$$

**Q.4. (a) What is the value of  $i$  when solute is associated and dissociated ?**

**(b) Calculate the freezing point of an aqueous solution containing 10.50 g of  $\text{MgBr}_2$  in 200 g of water. (Molar mass of  $\text{MgBr}_2 = 184$ ,  $K_f = 1.86^\circ\text{C kg mol}^{-1}$ )**

Ans. (a)  $i < 1$  when solute is associated and  
 $i > 1$  when solute is dissociated.

$$\begin{aligned} (b) \quad m &= \frac{n_g \times 1000}{W_A(g)} \\ &= \frac{W_b \times 1000}{W_B \times W_A} = \frac{10.50 \times 1000}{184 \times 200} = 0.2853\text{M} \end{aligned}$$



$$i = 3$$

$$\begin{aligned} \Delta T_f &= i \times K_f \times M \\ &= 3 \times 1.86 \times 0.2855 \\ &= 1.59 \end{aligned}$$

$$\text{Freezing point} = 0 - 1.59^\circ\text{C} = -1.59^\circ\text{C}$$



**Q.5 (a)** What is the value of  $i$  for  $\text{Al}_2(\text{SO}_4)_3$  when it is completely dissociated?

**(b)** Calculate the boiling point of a solution prepared by adding 15.00 g of NaCl to 250 g of water. ( $K_b = 0.512 \text{ K kg mol}^{-1}$  and molar mass of NaCl =  $58.44 \text{ g mol}^{-1}$ )

Ans.(a)  $\text{Al}_2(\text{SO}_4)_3 \rightarrow 2\text{Al}^{3+} + 3\text{SO}_4^{2-}$

$$i=5$$

$$\Delta T_b = \frac{iK_b \times 1000 \times W_B}{W_A \times M_B}$$

b)  $\text{NaCl} \rightarrow \text{Na}^+ + \text{Cl}^-$

$$i = 2$$

$$\Delta T_b = \frac{2 \times 0.512 \times 1000 \times 15}{250 \times 58.44}$$

$$= 1.05$$

Boiling Point of solution =  $100 + 1.05 = 101.05^\circ\text{C}$

**Q.6(a)** Calculate the freezing point of solution when 1.9 g of  $\text{MgCl}_2$  ( $M=95 \text{ g mol}^{-1}$ ) was dissolved in 50g of water, assuming  $\text{MgCl}_2$  undergoes complete ionization. ( $K_f$  for water =  $1.86 \text{ K kg mol}^{-1}$ ).

**(b) (i)** Out of 1 M glucose and 2 M glucose, which one has a higher boiling point and why?

**ii)** What happens when the external pressure applied becomes more than the osmotic pressure of solution?

Ans. (a)

$$\Delta T_f = \frac{K_f \times W_b \times 1000}{M_b \times W_a}$$

$$\Delta T_f = 3 \times (1.86 \times 1.9 / 95 \times 50) \times 1000 = 2.23 \text{ K}$$

$$T_f - \Delta T_f = 276.15 - 2.23 / 273 - 2.23$$

$$T_f = 270.92 \text{ K or } 270.77 \text{ K}$$

(b) (i) 2 M glucose; More number of particles/less vapour pressure

(ii) Reverse osmosis

**Q.7. (a) When 2.56 g of sulphur was dissolved in 100 g of CS<sub>2</sub>, the freezing point lowered by 0.383 K. Calculate the formula of sulphur S<sub>x</sub> [K<sub>f</sub> for CS<sub>2</sub> = 3.83 K Kg mol<sup>-1</sup>. Atomic mass of sulphur = 32 g mol<sup>-1</sup>]**

**(b) ) Blood cells are isotonic with 0.9% sodium chloride solution. What happens if we place blood cells in a solution containing.**

**(1) 1.2% sodium chloride solution?**

**(ii) 0.4% sodium chloride solution?**

**Ans. (a)**

$$\Delta T_f = \frac{K_f \times w_b \times 1000}{M_b \times w_a}$$

$$0.383 = (3.83 \times 2.56 / M \times 100 \times 1000)$$

$$M = 256$$

$$S_x \chi = 256$$

$$32x\chi = 256$$

$$\chi = 8$$

(b) (i) Shrinks (ii) Swells

**Q.8. (a) How will you determine the molecular mass from the relative lowering of vapour pressure?**

**(b) ) At 298 K, the vapour pressure of water is 23.75 mm Hg. Calculate the vapour pressure at the same temperature over 5% aqueous solution of urea NH<sub>2</sub>CONH<sub>2</sub>.**

**Hint: (i) Refer "Points To Remember"**

**(ii) According to Raoult's law,**

Substituting the values, we have 
$$\frac{P^0 - P_s}{P^0} = \frac{w_2}{M_2} \times \frac{M_1}{w_1}$$

$$\frac{23.75 - P_s}{23.75} = \frac{5 \times 18}{60 \times 95}$$

$$\Rightarrow 23.75 - P_s = \frac{5 \times 18}{60 \times 95} \times 23.75 = 0.375$$

$$\therefore P_s = 23.75 - 0.375 = 23.375 \text{ mm}$$

**Q9. (a) List three points of differences between ideal solution and non-ideal solution.**

**(b) Calculate the boiling point elevation for a solution prepared by adding 10 g of  $\text{CaCl}_2$  to 200 g of water. ( $K_b$  for water =  $0.512 \text{ K kg mol}^{-1}$ , Molar mass of  $\text{CaCl}_2 = 111 \text{ g mol}^{-1}$ )**

$$\Delta T_f = i K_b m = i K_b \cdot \frac{w_2}{M_2} \times \frac{1000}{w_1}$$

For  $\text{CaCl}_2; i = 2$   
 $K_b = 0.512 \text{ K kg mol}^{-1}$

$$\Delta T_b = 2 \times 0.512 \times \frac{10}{110} \times \frac{1000}{200}$$

$$\therefore \Delta T_b = 0.461 \text{ K}$$

**Q10. (i) Give reason for the following:**

**(a) Cold drink bottles are sealed at high pressure  $\text{CO}_2$**

**(b) Aquatic species are more comfortable in cold water than in warm water.**

**(ii) Calculate the amount of  $\text{KCl}$  which must be added to 100 g of water so that water freezes at  $-2.0^\circ\text{C}$ . Assume that  $\text{KCl}$  undergoes complete dissociation. [Given,  $K_f$  for water  $1.86 \text{ K/m}$ ]**

Ans. (a)  $\text{CO}_2$  is less soluble in water, so according to Henry's law to increase its solubility bottles are sealed at high pressure of  $\text{CO}_2$ .

(b) In warm water  $K_H$  value of  $\text{CO}_2$  is greater than in cold water, consequently solubility of  $\text{CO}_2$  in water increases with decrease of temperature. Hence aquatic species are more comfortable in cold water rather than in warm water.

(ii)  $\text{KCl} \rightarrow \text{K}^+ + \text{Cl}^-$

$$\text{van't Hoff factor (i)} = \frac{2}{1} = 2$$

According to formula  $\Delta T_f = i K_f m \Rightarrow 2 = 2 \times 1.86 \times m$

$$\therefore m = \frac{1}{1.86} = 0.538 \text{ m}$$

$$\text{Amount of KCl dissolved in 100g} = \frac{0.538}{1000} \times 100 \times 74.5 = 4.008 \text{ g}$$

[  $\therefore$  Molar mass of  $\text{KCl} = 74.5$  ]

### CASE STUDY BASED QUESTION

1. **Read the passage given below and answer the questions that follow:**

Dissolution of solids in water can be accompanied by absorption or evolution of heat i.e. dissolution process may be exothermic or endothermic in nature but dissolution of gases in water is an exothermic process. Dissolution of a substance in water is either due to ion dipole interaction or by hydrogen bond formation. Dissolution of gases in water is highly affected by pressure. The quantitative relationship between the solubility of gas in liquid and pressure is given by Henry's law in the form of mathematical relationship  $p = K_H \chi_B$ .

(A) **Dissolution of glucose in water can be explained by:**

- |                         |                            |
|-------------------------|----------------------------|
| (a) Hydrogen bond       | (b) ion-ion interaction    |
| (c) vander Waals' force | (d) ion-dipole interaction |

(B) **Solubility of KCl in water increases with the rise in temperature. This means that enthalpy of dissolution of KCl in water:**

- |        |                   |
|--------|-------------------|
| (a) =0 | (b) <0            |
| (c) >0 | (d) unpredictable |

(C) **The value of  $K_H$  for  $N_2$  gas in water at 298K is 86.76k bar, the value of  $K_H$  for  $N_2$  in water at 303K in kbar is :**

- |            |                   |
|------------|-------------------|
| (a) 86.76  | (b) >86.76        |
| (c) <86.76 | (d) unpredictable |

2. **Read the passage given below and answer the questions that follow:**

Many biological processes depend on osmosis, which is a spontaneous process by which the solvent molecules pass through a semi permeable membrane from a solution of higher concentration to a solution of higher concentration. The name osmosis is derived from the Greek word 'osmosis' which means 'to push'. It is also important to know that the semipermeable membrane selectivity allows certain

molecules in the solution to pass through it but not others. Two solutions having same osmotic pressure at a given temperature are called isotonic solutions. When such solutions are separated by a semipermeable membrane, solvent flow between one to the other one in either direction is same, i.e. the net solvent flow between the two isotonic solution is zero.

In the following questions a statement of assertion followed by a statement or reason is given. Choose the correct answer out of the following choices.

- a) Assertion and reason both are correct statements and reason is correct explanation for assertion.
  - b) Assertion and reason both are correct statements reason is not correct explanation for assertion.
  - c) Assertion is correct statement but reason is wrong statement.
  - d) Assertion is wrong statement but reason is correct statement.
- (A) **ASSERTION:** Among all the colligative properties, osmotic pressure measurement provides better method for determination of the molecular mass of the solute.  
**REASON:** Osmotic pressure measurement cannot be carried at room temperature.
- (B) **ASSERTION:** The osmotic pressure of 0.1 M urea solution is less than 0.1 M NaCl solution.  
**REASON :** Osmotic pressure is not a colligative property.
- (C) **ASSERTION :** The molecular mass of polymers cannot be calculated using the boiling point or freezing point method.  
**REASON :** The boiling point method for determining the molecular mass is used for compounds stable at high temperature.
- (D) **ASSERTION :** The elevation in boiling point for two isotonic solutions is same.  
**REASON :** The boiling point depends upon concentration of solute.

3. **Read the passage given below and answer the questions that follow:**

The colligative property of a solution is a property that depends only on the number of solute particles present, not on their identity. An ideal solution is a solution in which all components obey Raoult's law (i.e.,  $P_A = x_A P_A^0$ ) throughout the composition range. The vapour pressure of a binary volatile mixture is  $P = P_B^0 + (P_A^0 - P_B^0)x_A$ . The composition of the vapour is given by  $Y_A^v = x_A P_A^0 / [P_B^0 + (P_A^0 - P_B^0)x_A]$  and  $Y_B = 1 - Y_A$ . The total vapour pressure of a mixture is  $P = P_A^0 P_B^0 / [P_A^0 + (P_B^0 - P_A^0)Y_A]$ . Azeotrope is a mixture that boils without change in composition. In colligative properties, the elevation of boiling point is given by  $\Delta T = k_b m$  and the depression of freezing point by  $\Delta T_f = k_f m$ . During dissociation of ionic electrolytes, the van't Hoff factor equals,  $i = 1 + (n-1)\alpha$ .

During association of electrolytes,  $i = 1 - \beta + \beta/n$

Here  $\alpha$  and  $\beta$  are the degrees of dissociation and association, respectively, of electrolytes.

(A) **The vapour-phase compositions in two binary liquid mixtures follow:**

- |                  |                  |
|------------------|------------------|
| (a) Boyle's law  | (b) Dalton's law |
| (c) Raoult's law | (d) Henry's law  |

(B) **The mole fraction of a solute is 0.4. The relative lowering of vapour pressure is :**

- |         |         |
|---------|---------|
| (a) 60% | (b) 80% |
| (c) 40% | (d) 20% |

(C) **Which is not a colligative property?**

- |                                  |                      |
|----------------------------------|----------------------|
| (a) Elevation in boiling point   | (b) Boiling point    |
| (c) Depression in freezing point | (d) Osmotic pressure |

(D) **The most accurate method for the measurement of molar mass is:**

- |                      |                  |
|----------------------|------------------|
| (a) osmotic pressure | (b) ebullioscopy |
| (c) cryoscopy        | (d) Raoult's law |

## ANSWERS

### I MULTIPLE CHOICE QUESTIONS

- |         |         |         |         |         |         |         |
|---------|---------|---------|---------|---------|---------|---------|
| 1. (b)  | 2. (a)  | 3. (d)  | 4. (d)  | 5. (d)  | 6. (a)  | 7. (d)  |
| 8. (c)  | 9. (a)  | 10. (a) | 11. (c) | 12. (c) | 13. (b) | 14. (a) |
| 15. (c) | 16. (d) | 17. (c) | 18. (d) | 19. (c) | 20. (d) | 21. (c) |
| 22. (a) | 23. (c) | 24. (a) | 25. (c) | 26. (a) | 27. (c) | 28. (d) |

**II FILL IN THE BLANKS**

- |                      |                       |                    |
|----------------------|-----------------------|--------------------|
| 1. One               | 2. Azeotropic mixture | 3. Reverse osmosis |
| 4. Solute            | 5. Increase           | 6. Greater than    |
| 7. van't Hoff factor | 8. Osmotic pressure   | 9. Edema           |
| 10 Association       |                       |                    |

**III ASSERTION REASON TYPE QUESTIONS**

- |        |        |        |         |        |        |
|--------|--------|--------|---------|--------|--------|
| 1. (a) | 2. (a) | 3. (d) | 4. (c)  | 5. (a) | 6. (d) |
| 7. (a) | 8. (a) | 9. (a) | 10. (d) |        |        |

**IV ONE WORD ANSWER TYPE QUESTIONS**

- |                             |                               |                        |
|-----------------------------|-------------------------------|------------------------|
| 1. Mole fraction            | 2. X                          | 3. $O_2$               |
| 4. Molal elevation constant | 5. $K \text{ kg mol}^{-1}$    | 6. Positive deviation  |
| 7. Greater                  | 8. Henry's law                | 9. Molality            |
| 10. Positive                | 11. One                       | 12. Negative deviation |
| 13. Anoxia                  | 14. Desalination of sea water |                        |
| 15. Association             |                               |                        |

**CASE STUDY TYPE QUESTIONS:****PASSAGE: 1** (A) a (B) c (C) b**PASSAGE: 2** (A) c (B) c (C) a (D) c**PASSAGE : 3** (A) c (B) c (C) b (D) a


**UNIT TEST - 1**  
**CHAPTER-2**  
**SOLUTIONS**

**TIME ALLOWED: 1 HR.**

**M.M. 20**

1. 250g fluoride is present in 1000 kg toothpaste sample, concentration of fluoride in ppm is 1
  - (a) 250 ppm
  - (b) 25 ppm
  - (c) 2500 ppm
  - (d) 4 ppm
2. At a given temperature, the osmotic pressure of a concentrated solution of a substance 1
  - (a) is higher than that of a dilute solution.
  - (b) is lower than that of a dilute solution
  - (c) is same as that of a dilute solution
  - (d) cannot be compared with osmotic pressure of dilute solution.
3. The value of Henry's law constant  $K_H$  is: 1
  - (a) Greater for gases with higher solubility
  - (b) greater for gases with lower solubility
  - (c) constant for all gases
  - (d) not related with the solubility of gases
4. What type of deviation from Raoult's law is shown by the liquid mixture forming minimum boiling azeotrope? 1
5. Justify that relative lowering in vapour pressure is a colligative property. 1
6. Draw the graph between vapour pressure and temperature and explain the elevation in boiling point of a solvent in solution. 2
7.  $\text{CCl}_4$  and water are immiscible whereas ethanol and water are miscible in all proportions. Explain. 2
8. The graphical representation of vapour pressures of two component system as a function of composition is given below. On the basis of graph answer the questions that follow: 3





- (i) Are the A-B interactions weaker, stronger or of the same magnitude as A-A and B-B?
  - (ii) Name the type of deviation from Raoult's law exhibited by this solution.
  - (iii) Predict the sign of  $\Delta_{\text{mix}} H$  for this system.
  - (iv) Predict the sign of  $\Delta_{\text{mix}} V$  for this solution.
  - (v) Give one example of such a solution.
  - (vi) What type of azeotrope will this system form?
9. A solution containing 1.9g per 100 mL of KCl (molar mass=74.5  $\text{gmol}^{-1}$ ) is isotonic with a solution containing 3g per 100 mL of urea (molar mass=60 $\text{gmol}^{-1}$ ). Calculate the degree of dissociation of KCl solution. Assume that both the solutions have same temperature. 3
10. (i) Boiling point is not a colligative property but elevation in boiling point is a colligative property. Comment. 5
- (ii) What happens when we place the red blood cell in distilled water?
  - (iii) State Raoult's law for a solution containing non-volatile solute.
  - (iv) Define Cryoscopic constant.

## UNIT TEST -2

### CHAPTER-2

### SOLUTION

**TIME ALLOWED: 1 HR.**

**M.M. 20**

1. Which of the following is a dimensionless quantity: molarity, molality or mole fraction? (1)
2.  $N_2$  and  $O_2$  gases have  $K_H$  values 76.48 kbar and 34.86 kbar respectively at 293 K temperature. Which one of these will have more solubility in water? (1)
3. Liquid 'Y' has higher vapour pressure than liquid 'X'. Which of them will have higher boiling point? (1)
4. Mention the unit of ebullioscopic constant (molal elevation constant). (1)
5. What is the maximum value of Van't Hoff factor ( $i$ ) for  $Na_2SO_4 \cdot 10H_2O$ ? (1)
6. Define the term osmosis and osmotic pressure. What is the advantage of using osmotic pressure as compared to other colligative properties for the determination of molar masses of solutes in solutions? (2)
7. Account for the following: (2)
  - (a) Aquatic species are more comfortable in winter than summer.
  - (b) Solution of acetone and  $CHCl_3$  is not an ideal solution.
8. Define isotonic solutions. A 5% solution of cane sugar (Molar mass =  $342 \text{ g mol}^{-1}$ ) is isotonic with 0.887% solution of urea. Find the molar mass of urea. (3)
9. A solution containing 8 g of a substance in 100 g of diethyl ether boils at  $36.86^\circ\text{C}$ , whereas pure ether boils at  $35.6^\circ\text{C}$ . Determine the molar mass of the solute. [For diethyl ether  $K_b = 2.02 \text{ K kg mol}^{-1}$ ] (3)
10. (a) How will you determine the molecular mass from the relative lowering of vapour pressure? (b) At 298 K, the vapour pressure of water is 23.75 mm Hg. Calculate the vapour pressure at the same temperature over 5% aqueous solution of urea  $NH_2CONH_2$ . (5)


## UNIT 3

## Electrochemistry

### Points to Remember

**Galvanic cells:** A galvanic cell is a device in which chemical energy is converted into electrical energy. e.g. Daniell cell.

- Daniell cell consists of two beakers containing  $\text{CuSO}_4$  and  $\text{ZnSO}_4$  solutions. A zinc rod is dipped into  $\text{ZnSO}_4$  while a copper rod is dipped into  $\text{CuSO}_4$  solution. In this cell zinc reacts with copper (II) ions and produces metallic copper and zinc (II) ion according to the reaction :



Electrons flow from anode to cathode in the external circuit. The solutions of two beakers are connected by salt bridge.

#### Functions of salt bridge :

- (i) It allows the flow of current by completing the electrical circuit.
- (ii) It maintains electrical neutrality of the cell.

#### Electrode Potential

The potential difference that develops between the metal and its electrolyte is known as electrode potential.

(a) **Oxidation potential:** The tendency of an electrode to lose electrons or to get oxidised is called oxidation potential. For example,  $\text{M(s)} \longrightarrow \text{M}^{n+}(\text{aq}) + n\text{e}^-$

(b) **Reduction potential:** The tendency of an electrode to gain electrons or to get reduced is called reduction potential. For example,  $\text{M}^{n+}(\text{aq}) + n\text{e}^- \longrightarrow \text{M(s)}$

According to latest IUPAC convention, the half reactions are always written as reduction half reaction and their potentials are represented by reduction potentials

$$E_{\text{ox}}^{\circ} = - E_{\text{red}}^{\circ}$$

#### Cell potential of a cell

The potential difference between the two electrodes of galvanic cell is called the cell potential and is measured in volts.

$$E_{\text{cell}}^{\circ} = E_{\text{cathode}}^{\circ} - E_{\text{anode}}^{\circ}$$

- Half cell potential or electrode potential of  $\text{M}^{n+}/\text{M}$  cannot be measured directly because a half cell whether oxidation or reduction half cell cannot work on its own. We cannot determine the absolute electrode potential of an electrode. To solve this problem, a reference electrode standard hydrogen electrode (SHE) or normal hydrogen electrode (NHE) is used, its standard electrode potential (oxidation as well as reduction) is arbitrarily taken as zero.

**Electrochemical series or e.m.f. series**

Arrangement of different electrodes in the order (increasing or decreasing) of their standard electrode potentials.

**Applications of the electrochemical series:**

1. **To compare the relative oxidising and reducing powers:** substances with higher reduction potentials are stronger oxidising agents.
2. **Calculation of standard emf of electrochemical cell ( $E^\circ_{\text{cell}}$ ):**  

$$E^\circ_{\text{cell}} = E^\circ_{\text{Cathode}} - E^\circ_{\text{anode}}$$
3. **Comparison of the reactivity of metals:** A metal with smaller reduction potential has a strong tendency to get oxidised and can displace metals having greater reduction potential from the aqueous solution of their salts.
4. **To predict whether a metal displace hydrogen from acids :** Metals having a negative reduction potential value can displace hydrogen from acid.
5. **To predict the spontaneity of a redox reaction:**  $E_{\text{cell}}$  should be positive. If emf comes out to be negative, the direct reaction as given cannot take place, the reverse reaction may take place.

**Effect of opposing potential on the cell reaction**

Consider a Daniell cell:  $\text{Zn}_{(\text{s})} | \text{Zn}^{2+}_{(\text{aq})} || \text{Cu}^{2+}_{(\text{aq})} | \text{Cu}$

- When Subscript  $E_{\text{ext}} < 1.1\text{V}$ , Electrons flow from Zn rod to Cu rod hence current flows from Cu to Zn.
- When  $E_{\text{ext}} > 1.1\text{ V}$ , flow of current in this case occurs from zinc electrode to copper electrode, the cell now **electrolytic cell**.

**NERNST EQUATION FOR SINGLE ELECTRODE**

For the electrode reaction  $\text{M}^{n+}(\text{aq}) + n\text{e}^- \rightarrow \text{M}(\text{s})$

$$E_{\text{M}^{n+}/\text{M}} = E^\circ_{\text{M}^{n+}/\text{M}} - \frac{RT}{nF} \ln \frac{[\text{M}_{(\text{s})}]}{[\text{M}^{n+}_{(\text{aq})}]}$$

Concentration of pure solid [M] should be taken as unity

$$E_{\text{M}^{n+}/\text{M}} = E^\circ_{\text{M}^{n+}/\text{M}} - \frac{2.303 RT}{nF} \log \frac{1}{[\text{M}^{n+}]} \quad (\text{since } \log_e = 2.303 \log_{10})$$

At 298 K

$$E_{\text{M}^{n+}/\text{M}} = E^\circ_{\text{M}^{n+}/\text{M}} - \frac{0.059}{n} \log \frac{1}{[\text{M}^{n+}]}$$

**Calculation of cell potential using Nernst Equation**

$$E_{\text{cell}} = E_{\text{cell}}^{\circ} - \frac{2.303RT}{nF} \log_{10} \frac{[\text{PRODUCTS}]}{[\text{REACTANTS}]} \quad \text{At 298K}$$

$$E_{\text{cell}} = E_{\text{cell}}^{\circ} - \frac{0.059}{n} \log_{10} \frac{[\text{PRODUCTS}]}{[\text{REACTANTS}]}$$

**EQUILIBRIUM CONSTANT (K<sub>c</sub>) FROM NERNST EQUATION**

$$E_{\text{cell}}^{\circ} = \frac{2.303RT}{nF} \log_{10} K_c$$

$$E_{\text{cell}}^{\circ} = \frac{0.059}{n} \log_{10} K_c \quad \text{at 298 K}$$

**GIBB'S ENERGY CHANGE AND CELL POTENTIAL**

$$\Delta_r G = -nF E_{\text{cell}}$$

$$\Delta_r G^{\circ} = -nF E_{\text{cell}}^{\circ}$$

**⇒ For cell reaction to be spontaneous, ΔG° must be negative, the value of E°<sub>cell</sub> must be positive.**

**FREE ENERGY CHANGE AND EQUILIBRIUM CONSTANT**

$$\Delta_r G^{\circ} = -2.303 RT \log_{10} K_c$$

**CONDUCTORS:** Substances that allow the flow of electric current through them are called electrical conductors .

**Metallic / electronic Conductor**

- Flow of electricity due to movement of electrons
- No chemical change as there is no transfer of matter.
- Faraday's law is not followed
- Conduction decreases with temperature because kernels start vibrating faster which interfere in the flow of electrons.

**Electrolytic conductor**

- Flow of electricity due to movement of ions
- Ions are oxidised or reduced at the electrodes, hence involve transfer of matter .
- Faraday's law is followed
- Conduction increase with temperature because dissociation increases and viscosity decreases

**FACTOR AFFECTING ELECTROLYTIC CONDUCTANCE**

1. **Interionic interactions:** Greater the interionic interactions lesser is the mobility of the ions, hence lesser will be conductance.
2. **Solvation of ions:** More the solvation of the ions, the lesser will be the electrical conductivity.
3. **Viscosity of the solvent:** Higher the viscosity of the solvent, lesser is the mobility of ions.
4. **Temperature:** As the temperature of the electrolytic solution is increased, the kinetic energy of the ions in the solution increases, hence their mobility increases. This results in the increase of electrical conductance of the electrolytic solution.
5. **Effect of concentration of solution:** More the concentration of electrolytic solution smaller will be its electrical conductivity.
  - Weak electrolyte ionise to a lesser extent in concentrated solution, on dilution ionisation increases which causes increase in conductivity.
  - Strong electrolyte ionise completely. On dilution interionic attraction decreases, so mobility of ions increases, consequently conductance increases.

**Electrolytic conduction**

**Resistance (R) :** A measure of obstruction in the flow of current. Unit: ohm ( $\Omega$ )

$$R \propto \frac{l}{A} \quad R = \rho \frac{l}{A}$$

$\rho$ , constant of proportionality, known as specific resistance or resistivity.

**Resistivity or specific resistance ( $\rho$ ) :**

$$\rho = R \frac{A}{l}$$

Resistivity may be defined as the resistance offered by the conductor of 1 m length with area of cross section equal to  $1 \text{ m}^2$

**Unit :** ohm. m or  $\Omega \cdot \text{m}$

**Conductance (G):** Conductance is a measure of the ease with which current flows through the conductor. It is reciprocal of electrical resistance.

$$G = 1/R$$

**Units:**  $\text{ohm}^{-1}$  or  $\Omega^{-1}$  i.e., Siemen (S),  $1 \text{ S} = 1 \Omega^{-1}$

**Specific conductance or conductivity (k):** conductivity is the reciprocal of resistivity.


$$k = 1/\rho = \frac{1}{R} \cdot \frac{l}{a}, \quad l/a \text{ is known as cell constant (G*)}$$

$$k = C \cdot G^* \text{ i.e. } \boxed{\text{Conductivity} = \text{Conductance} \times \text{cell constant}}$$

Units of  $k = \text{ohm}^{-1} \text{cm}^{-1}$  or SI units are  $\text{ohm}^{-1} \text{m}^{-1}$  or  $\text{S m}^{-1}$

- **Molar conductivity ( $\Lambda_m$ )** - of a solution is the conductance of all the ions produced from one mole of the electrolyte dissolved in a given volume of the solution when the electrodes are one cm apart and the area of the electrodes is so large that the whole of the solution is contained between them.  
 $\boxed{\Lambda_m = k \times 1000 / \text{Molarity}}$ , Unit:  $\text{ohm}^{-1} \text{cm}^2 \text{mol}^{-1}$  ( $\text{S cm}^2 \text{mol}^{-1}$ ), SI unit =  $\text{S m}^2 \text{mol}^{-1}$
- **Effect of dilution.** Conductance increases (because total no. of ions increase), conductivity decreases (because no. of ions per unit volume decreases), molar conductivity increase with dilution.
- **Variation of molar conductivity with concentration.** For a strong electrolyte, it is given by **Debye - Huckel - Onsager equation**:  $\Lambda_m = \Lambda_m^0 - A\sqrt{c}$  where  $A$  is a constant depending upon the nature of the solvent and temperature.  $\Lambda_m^0$  is **limiting molar conductivity** and it is defined as the molar conductivity of electrolyte when concentration approaches zero i.e. at infinite dilution.

**Graphical representation of the variation of  $\Lambda_m$  vs  $\sqrt{c}$**



It can be seen that if we plot  $\Lambda_m$  against  $c^{1/2}$ , we obtain a straight line with intercept equal to  $\Lambda_m^0$  and slope equal to  $-A'$ .

**Reasons for increase of  $\Lambda_m$  with dilution.**

- (Molar conductivity of a strong electrolyte increases with dilution because interionic attractions decrease with dilution. Small deviations at higher concentration are due to large interionic attractions.) (Molar conductivity of a weak electrolyte increases with dilution because dissociation increases with dilution.)

**Inability to determine limiting molar conductivity experimentally for a weak electrolyte.** Molar conductivity at infinite dilution for a strong electrolyte can be found by extrapolation to zero concentration but that of weak electrolyte cannot be thus found.

**KOHLRAUSCH LAW OF INDEPENDENT MIGRATION OF IONS**

The law states that limiting molar conductivity of an electrolyte can be represented as the sum of the individual contributions of the anion and cation of the electrolyte.

For e.g.  $\text{Al}_2(\text{SO}_4)_3$   $\Lambda_m^0[\text{Al}_2(\text{SO}_4)_3] = 2\lambda_m^0(\text{Al}^{3+}) + 3\lambda_m^0(\text{SO}_4^{2-})$








**Applications of Kohlrausch's Law**

- In calculation of limiting molar conductivity of weak electrolyte
- In calculation of degree of dissociation, i.e.,  $\alpha = \Lambda_m / \Lambda_m^0$
- In calculation of dissociation constant ( $K_c$ ) by using value of  $\alpha$ ,  $K_c = c\alpha^2 / 1 - \alpha$


**ELECTROLYTIC CELLS** : The device in which conversion of electrical energy into chemical energy is done is known as electrolytic cell.

**PRODUCTS OF ELECTROLYSIS:** If an electrolytic solution consists of more than two ions then during electrolysis all the ions are not discharged simultaneously but certain ions are liberated at the electrode in preference to the others. This is based on the principle of preferential discharged theory which states that the ion which requires least energy is discharged first.



**Batteries and Fuel Cells :****Batteries are classified as two types:****Primary:** use oxidation-reduction reactions that cannot be reversed very easily**Secondary :** reactions of these batteries can be reversed (rechargeable batteries)**Primary Cells :****(i) Dry cell :****(ii) Mercury Cell :****Secondary Cells :****(i) Lead storage battery**



On recharging cell reaction is reversed.

**(ii) Nickel — Cadmium cell.**

Cell reaction is reversed on recharging.

**FUEL CELLS :**It is an electrochemical device used to convert free energy of a chemical reaction to electrical energy. Fuel used are in gaseous state ( $\text{H}_2$ ,  $\text{CH}_4$ ,  $\text{CO}$  etc)e.g.  **$\text{H}_2$ - $\text{O}_2$  fuel cell.**

Reactions are


**Advantages**

- 1) Efficient than any conventional source.
- 2) No pollution.
- 3) Electrodes are not affected


**CORROSION :** It involves the slow destruction of a metal as a result of its reaction with moisture and gases present in atmosphere. More reactive metals corrode more easily. Corrosion of Iron is called rusting.

**Mechanism of Rusting****Electrochemical theory of rusting :**


Impure surface of iron act as an electrochemical cell. Pure Iron act as anode and impure iron as cathode. Carbonic acid act as the electrolyte (It provide  $H^+$  ion).



Setting up of Electrochemical cell on Iron surface



Cathode is further oxidised by atmospheric oxygen to form rust.  $Fe_2O_3 \cdot xH_2O$

**Prevention of corrosion :**

1) **Barrier protection** — By coating with a suitable material — paint, oil, grease etc

2) **Sacrificial protection** — Coating with a more reactive metal. The process of coating the surface of iron with Zinc is called Galvanization. More reactive metal act as anode.

3) **Alloying** with metals that form oxide coats.

4) **Antirust solutions** — Alkaline phosphate or chromate solutions are applied on iron surface to form a heat resistant iron phosphate or chromate coating which prevent corrosion.

4) **Cathodic protection** - Here metal to be protected is set as cathode by attaching a more reactive metal to it. Now the more reactive metal undergo oxidation. for ex. Zn, Al or Mg can be used for cathodic protection.

### OBJECTIVE TYPE QUESTIONS

#### I. MULTIPLE CHOICE QUESTIONS

1. The potential of a hydrogen electrode at pH=10 is :
 

(a) 0.591 V	(b) 0.00 V
(c) -0.591 V	(d) -0.059 V
2. How many coulomb are required for the oxidation of 1 mol of  $\text{H}_2\text{O}_2$  to  $\text{O}_2$ ?
 

(a) $9.65 \times 10^4 \text{ C}$	(b) 93000 C
(c) $1.93 \times 10^5 \text{ C}$	(d) $19.3 \times 10^2 \text{ C}$
3. KCl is used in salt bridge because:
 

(a) It forms a good jelly with agar-agar	(b) It is a strong electrolyte
(c) It is a good conductor of electricity	(d) Migration factor of $\text{K}^+$ and $\text{Cl}^-$ ions are almost equal
4. For a spontaneous reaction the  $\Delta G$ , equilibrium constant (K) and  $E^\circ_{\text{cell}}$  will be respectively.
 

(a) -ve, < 1, - ve	(b) -ve, > 1, - ve
(c) -ve, > 1, + ve	(d) +ve, > 1, - ve
5. If a salt bridge is removed between the half cells, the voltage:
 

(a) drops to zero	(b) does not change
(c) increase gradually	(d) increases rapidly
6. The process in which chemical change occurs on passing electricity is termed :
 

(a) Ionisation	(b) neutralisation
(c) electrolysis	(d) hydrolysis
7. The charge required for the reduction of 1 mol of  $\text{MnO}_4^-$  to  $\text{MnO}_2$  is:
 

(a) 1F	(b) 3F
(c) 5F	(d) 4F
8. The value of  $\Lambda_m^\circ$  for  $\text{NH}_4\text{Cl}$ ,  $\text{NaOH}$  and  $\text{NaCl}$  are 129.8, 248.1 and 126.4  $\text{Ohm}^{-1}\text{cm}^2\text{mol}^{-1}$  respectively. Calculate  $\Delta_m^\circ$  for  $\text{NH}_4\text{OH}$  solution.
 

(a) $215.5 \text{ Ohm}^{-1}\text{cm}^2\text{mol}^{-1}$	(b) $251.5 \text{ Ohm}^{-1}\text{cm}^2\text{mol}^{-1}$
(c) $244.7 \text{ Ohm}^{-1}\text{cm}^2\text{mol}^{-1}$	(d) $351.5 \text{ Ohm}^{-1}\text{cm}^2\text{mol}^{-1}$

9. In a Galvanic cell the electrical work done is equal to :
  - (a) Free energy change
  - (b) mechanical work done
  - (c) thermodynamic work done
  - (d) all of the above
10. Zn cannot displace following ions from their aqueous solution :
  - (a)  $\text{Al}^{3+}$
  - (b)  $\text{Cu}^{2+}$
  - (c)  $\text{Fe}^{2+}$
  - (d)  $\text{Na}^+$
11. Electrical work done is equal to :
  - (a)  $-nFE_{\text{cell}}^0$
  - (b)  $nFE_{\text{cell}}^0$
  - (c)  $nE_{\text{cell}}^0$
  - (d) None of these
12. Which are not the following decrease with increase in concentration?
  - (a) Conductance
  - (b) Molar conductance
  - (c) Conductivity
  - (d) All of the above
13. The standard electrode potential values of three metallic cations, X, Y, Z are 0.52, -3.03 and -1.18V, respectively. The order of reducing power of the corresponding metals is
  - (a)  $Y > Z > X$
  - (b)  $X > Y > Z$
  - (c)  $Z > Y > X$
  - (d)  $Z > X > Y$
14. How is electrical conductance of a conductor related with length and area of cross section of the conductor?
  - (a)  $G = k.l.a^{-1}$
  - (b)  $G = l.a.k^{-1}$
  - (c)  $G = k.a.l^{-1}$
  - (d)  $G = k.l.a^{-2}$
15. What will happen during the electrolysis of aqueous solution of  $\text{CuSO}_4$  in the presence of Cu electrodes?
  - (a) Copper will deposit at cathode.
  - (b) Copper will dissolve at anode.
  - (c) Oxygen will be released at anode.
  - (d) Copper will deposit at anode.
16. The cell constant of a conductivity cell \_\_\_\_\_.
  - (a) changes with change of electrolyte.
  - (b) changes with change of concentration of electrolyte.
  - (c) changes with temperature of electrolyte.
  - (d) remains constant for a cell.
17. An electrochemical cell can behave like an electrolytic cell when
  - (a)  $E_{\text{cell}} = 0$
  - (b)  $E_{\text{cell}} > E_{\text{ext}}$
  - (c)  $E_{\text{ext}} > E_{\text{cell}}$
  - (d)  $E_{\text{cell}} = E_{\text{ext}}$
18. Which of the following statement is not correct about an inert electrode in a cell?
  - (a) It does not participate in the cell reaction.
  - (b) It provides surface either for oxidation or for reduction reaction.
  - (c) It provides surface for conduction of electrons.
  - (d) It provides surface for redox reaction.

19. The difference between the electrode potentials of two electrodes when no current is drawn through the cell is called
- Cell potential
  - Cell emf
  - Potential difference
  - Cell voltage
20. The positive value of the standard electrode potential of  $\text{Cu}^{2+}/\text{Cu}$  indicates that :
- this redox couple is a stronger reducing agent than the  $\text{H}^+/\text{H}_2$  couple.
  - this redox couple is a stronger oxidising agent than  $\text{H}^+/\text{H}_2$ .
  - Cu can displace  $\text{H}_2$  from acid.
  - Cu cannot displace  $\text{H}_2$  from acid.

## II FILL IN THE BLANKS:

- The conductance of a solution placed between two opposite faces of a centimetre cube is called.....
- Strong electrolytes give almost a linear plot of  $\Lambda_m$  versus.....
- In a galvanic cell, the electrons flow from ..... to.....through connecting wire.
- The unit of resistivity is .....
- For the spontaneous cell reaction,  $E^0$  should be.....
- Conductivity of an electrolytic solution.....with increase in dilution.
- The unit of cell constant is.....
- An aqueous solution of copper nitrate.....be stored in iron vessel.
- Protection of iron by coating with zinc is called .....
- To deposit 2 mol of Ca from  $\text{CaCl}_2$ .....electricity is required.

## III ASSERTION REASON TYPE QUESTIONS

- Both assertion and reason are correct statements, and reason is the correct explanation of the assertion.
- Both assertion and reason are correct statements, but reason is not the correct explanation of the assertion.
- Assertion is correct, but reason is wrong statement.
- Assertion is wrong, but reason is correct statement.

1. **Assertion:** When aqueous sodium chloride solution is electrolysed, Oxygen gas is produced at the anode.  
**Reason :** It is due to the overpotential for oxidation of water to oxygen.
2. **Assertion:** Molar conductivity of an electrolyte increases with decrease in concentration.  
**Reason:** The mobility of ions decrease with increase in concentration.
3. **Assertion:** Reduction of 1 mole of  $\text{Cu}^{2+}$  ions require 2 faraday of charge.  
**Reason:** 1 Faraday is equal to the charge of 1 mole of electrons.
4. **Assertion:** Leclanche cell gives constant voltage throughout its life.  
**Reason:** The overall reaction of button cell does not involve any ion in solution whose concentration can change during its life time.
5. **Assertion:** Coating iron with zinc prevents rusting.  
**Reason:** The coating of zinc prevents moist air to come in contact with the metal.
6. **Assertion:** More negative the electrode potential greater is the power to act as oxidising agent.  
**Reason:** As the electrode potential becomes more negative there is greater tendency to undergo oxidation.
7. **Assertion:** Secondary cells are cells which can be recharged after use.  
**Reason:** The products are electrolysed back to the initial reactants during recharge of the cell.
8. **Assertion :** Kohlrausch law helps to find the molar conductivity of weak electrolyte at infinite dilution.  
**Reason :** Molar conductivity of a weak electrolyte at infinite dilution cannot be determined experimentally.
9. **Assertion :** Fluorine is the best oxidising agent.  
**Reason :** Fluorine has highest reduction potential.
10. **Assertion:**  $\Lambda_m$  for weak electrolytes shows a sharp increase when the electrolyte solution is diluted.  
**Reason:** For weak electrolytes degree of dissociation increases with dilution of solution.

**IV ONE WORD ANSWER TYPE QUESTIONS**

1. How much charge in Faraday is required for the reduction of  $1 \text{ mol Al}^{3+}$  to Al?
2. What is the effect of increase of temperature on ionic conductance?
3. What flows in the internal circuit of a Galvanic cell?
4. Name of the reference electrode in determining the standard electrode potential.
5. Can  $E_{\text{cell}}^0$  for a cell reaction ever be equal to zero?
6. Name the quantity which is reciprocal of resistivity.
7. How will pH of brine (aqueous NaCl solution) be affected when it is electrolysed?
8. What is the name given to the constant quantity of charge carried by one mole electrons?
9. Name of reference electrode used for determination of  $E^0$  instead of NHE/SHE.
10. What is the effect of presence of salt in water on the rate of rusting of iron?
11. Name the type of cell which was used in Apollo Space Programme for providing electrical power.
12. Under what condition is  $E_{\text{cell}} = 0$  or  $\Delta_r G = 0$ ?
13. How is equilibrium constant of a reaction related to standard cell potential?
14. Mention the direction of flow of electrons in the following cell:  
 $\text{Zn(s)} | \text{Zn}^{2+}(\text{aq}) || \text{Ag}^+ | \text{Ag(s)}$
15. A galvanic cell has electrical potential of 1.1 V. If an opposing potential of 1.1 V is applied to this cell. What will happen to the cell reaction and current flowing through the cell?

**VERY SHORT ANSWER TYPE QUESTIONS (1 Mark Questions)**

**Q. 1. Why is it not possible to measure single electrode potential ?**

**Ans.** Because the half cell containing single electrode cannot work independently, as charge cannot flow on its own in a single electrode.

**Q. 2. Name the factors on which emf of a cell depends.**

**Ans.** Emf of a cell depends on following factors :

- (a) Nature of reactants
- (b) Concentration of solution in two half cells
- (c) Temperature

**Q. 3. What is the effect of temperature on the electrical conductance of metal ?**

**Ans.** Temperature increases, electrical conductance decreases.

**Q. 4. What is the effect of temperature on the electrical conductance of electrolyte ?**

**Ans.** Temperature increases, electrical conductance increases.

**Q. 5. What is the relation between conductance and conductivity ?**

**Ans.**  $\Lambda_m^c = \frac{k}{C}$

**Q. 6. Reduction potentials of 4 metals A, B, C and D are  $-1.66$  V,  $+0.34$  V,  $+0.80$  V and  $-0.76$  V. What is the order of their reducing power and reactivity ?**

**Ans.**  $A > D > B > C$

**Q.7. Why does a dry cell become dead even if it has not been used for a long time?**

**Ans.**  $\text{NH}_4\text{Cl}$  is acidic in nature. It corrodes zinc container

**Q.8. Why Na cannot be obtained by the electrolysis of aqueous NaCl solution ?**

**Ans.** Due to low reduction potential,  $\text{Na}^+$  ions are not reduced at cathode. Instead,  $\text{H}^+$  are reduced and  $\text{H}_2$  is obtained.

**Q.9. What is the use of platinum foil in the hydrogen electrode ?**

**Ans.** It is used for the in and out flow of electrons.

**Q.10. Why  $\Lambda_m^\circ$  for  $\text{CH}_3\text{COOH}$  cannot be determined experimentally ?**

**Ans.** Molar conductivity of weak electrolytes keeps on increasing with dilution and does not become constant even at very large dilution.

**Q.11. Why is it necessary to use a salt bridge in a galvanic cell?**

**Ans.** To complete the inner circuit and to maintain electrical neutrality of the electrolytic solutions of the half cells.



**Q.12. Why does mercury cell gives a constant voltage throughout its life?**

Ans. This is because the overall cell reaction does not have any ionic concentration in it.

**Q.13. What is the role of  $\text{ZnCl}_2$  in a dry cell ?**

Ans.  $\text{ZnCl}_2$  combines with the  $\text{NH}_3$  produced to form a complex salt  $[\text{Zn}(\text{NH}_3)_2]\text{Cl}_2$ .

**Q.14. Why does the conductivity of a solution decrease with dilution ?**

Ans. Conductivity of a solution is dependent on the number of ions per unit volume. On dilution, the number of ions per unit volume decreases, hence the conductivity decreases.

**Q.15. Suggest two materials other than hydrogen that can be used as fuels in fuel cells.**

Ans. Methane and methanol.

**Q.16. How does the pH of Al-NaCl solution be affected when it is electrolysed ?**

Ans. When Al-NaCl solution is electrolysed,  $\text{H}_2$  is liberated at cathode,  $\text{Cl}_2$  at anode and NaOH is formed in the solution. Hence pH of solution increase.

**Q.17. Which reference electrode is used to measure the electrode potential of other electrodes?**

Ans. Standard hydrogen electrode (SHE) whose electrode potential is taken as zero.

**Q.18. Out of zinc and tin, which one protects iron better even after cracks and why ?**

Ans. Zinc protects better because oxidation potential of zinc is greater but that of tin is less than that of iron.

**Q.19. Define corrosion. What is the chemical formula of rust ?**

Ans. Corrosion is the slow eating away of the surface of the metal due to attack of atmospheric gases.  $\text{Fe}_2\text{O}_3 \cdot x\text{H}_2\text{O}$

**20. What is the electrolyte used in a dry cell?**

Ans. A paste of  $\text{NH}_4\text{Cl}$

**21. How much electricity is required in Coulomb for the oxidation of 1 mole of  $\text{FeO}$  to  $\text{Fe}_2\text{O}_3$  ?**

Ans.  $\text{Fe}^{2+} \rightarrow \text{Fe}^{3+} + \text{e}^{-1}$   
So,  $IF = I \times 96500\text{C} = 96500\text{C}$

**22. Two metals A and B have reduction potential values - 0.76 V and +0.34V respectively. Which of these will liberate  $\text{H}_2$  from dil.  $\text{H}_2\text{SO}_4$  ?**

Ans. Metal having higher oxidation potential will liberate  $\text{H}_2$  from  $\text{H}_2\text{SO}_4$ . Thus, A will liberate  $\text{H}_2$  from  $\text{H}_2\text{SO}_4$ .

**23. How does conc. of sulphuric acid change in lead storage battery when current is drawn from it ?**

Ans. Concentration of sulphuric acid decreases.

**24. Why is alternating current used for measuring resistance of an electrolytic solution ?**

Ans. The alternating current is used to prevent electrolysis so that the concentration of ions in the solution remains constant.

**Q.25.**  $E^0$  values of  $\text{MnO}_4^-$ ,  $\text{Ce}^{4+}$  and  $\text{Cl}_2$  are 1.507, 1.61 and 1.358 V respectively.

**Arrange these in order of increasing strength as oxidizing agent.**

Ans.  $\text{Cl}_2 < \text{MnO}_4^- < \text{Ce}^{4+}$

**26. Explain Kohlrausch's law of independent migration of ions.**

Ans. It states that at infinite dilution, molar conductivity of an electrolyte is equal to sum of contributions due to cation as well as anion.


$$\Lambda_{\text{m}(\text{Na}_2\text{SO}_4)}^\infty = 2\Lambda_{\text{m}(\text{Na}^+)}^\infty + \Lambda_{\text{m}(\text{SO}_4^{2-})}^\infty$$

**27. Give products of electrolysis of an aqueous solution of  $\text{AgNO}_3$  with silver electrode.**

Ans. At anode:  $\text{Ag}_{(\text{s})} \rightarrow \text{Ag}_{(\text{aq})}^+ + \text{e}^-$   
 At cathode:  $\text{Ag}_{(\text{aq})}^+ + \text{e}^- \rightarrow \text{Ag}_{(\text{s})}$

### SHORT ANSWER TYPE QUESTIONS (2 or 3 Marks)

1. How can you increase the reduction potential of an electrode for the reaction :




Ans. Nernst equation is :

$$E_{M^{n+}/M} = E_{M^{n+}/M}^{\ominus} - \frac{0.0591}{n} \log \frac{1}{[M^{n+}]} \text{ at } 298K$$

$E_{M^{n+}/M}$  can be increased by

- (a) Increase in concentration of  $M^{n+}$  ions in solution.
- (b) By increasing the temperature.

2. Calculate emf of the following cell at 298 K :



The Nernst equation for the cell is :

$$[\text{Given : } E_{\text{cell}}^{\ominus} = 3.17 \text{ V}]$$

Ans.

$$\begin{aligned} E &= E_{\text{cell}}^{\ominus} - \frac{0.059}{2} \log \frac{[Mg^{2+}]}{[Ag^{+}]^2} \\ &= 3.17 - \frac{0.059}{2} \log \frac{.130}{(.0001)^2} \\ &= 3.17 - 0.21 = 2.96V \end{aligned}$$

Q.3. Suggest a way to determine the  $\Delta_m^{\ominus}$  value of water

Ans.  $\Delta_m^{\ominus}(H_2O) = \Delta_m^{\ominus}(H^{+}) + \Delta_m^{\ominus}(OH^{-})$

It can be determined from the value of

$\Delta_m^{\ominus}(HCl)$ ,  $\Delta_m^{\ominus}(NaOH)$  -  $\Delta_m^{\ominus}(NaCl)$ . then,

$\Delta_m^{\ominus}(H_2O) = \Delta_m^{\ominus}(HCl) + \Delta_m^{\ominus}(NaOH) - \Delta_m^{\ominus}(NaCl)$

Q. 4. How much electricity in term of Faraday is required to produce 40 gram of Al from  $Al_2O_3$  ? (Atomic mass of Al = 27 g/mol)


Ans.  $Al^{3+} + 3e^{-} \rightarrow Al$

27 gram of Al require electricity = 3F


40 gram of Al require electricity =  $\frac{3F}{27} \times 40 = 4.44 \text{ F}$

**Q.5. Predict the product of electrolysis of an aqueous solution of  $\text{CuCl}_2$  with an inert electrode.**


**Ans.**  $\text{CuCl}_2(\text{s}) + \text{Aq} \rightarrow \text{Cu}^{2+} + 2\text{Cl}^-$



At cathode (Reduction):  $\text{Cu}^{2+}$  will be reduced in preference to  $\text{H}^+$  ions.



At anode (Oxidation):  $\text{Cl}^-$  will be reduced in preference to  $\text{OH}^-$  ions.



**Q.6. Calculate  $\Lambda_m^\circ$  for  $\text{CaCl}_2$  and  $\text{MgSO}_4$  from the following data :**

$\Lambda_m^\circ(\text{Ca}^{2+}) = 119.0, \text{Scm}^2 \text{mol}^{-1}$ ,  $\text{Scm}^2 \text{mol}^{-1}$ ,  $\text{Mg}^{2+} = 106.0$ ,  $\text{Cl}^- = 76.3$  and  $\text{SO}_4^{2-} = 160.0 \text{Scm}^2 \text{mol}^{-1}$

$$\Lambda_{m(\text{CaCl}_2)}^\circ = \Lambda_{m(\text{Ca}^{2+})}^\circ + 2\Lambda_{m(\text{Cl}^-)}^\circ = 119 + (2 \times 76.3) = 271.6 \text{ S cm}^2 \text{mol}^{-1}$$

$$\Lambda_{m(\text{MgSO}_4)}^\circ = \Lambda_{m(\text{Mg}^{2+})}^\circ + 2\Lambda_{m(\text{SO}_4^{2-})}^\circ = 106 + 160 = 266 \text{ S cm}^2 \text{mol}^{-1}$$

**Q. 7. Calculate the potential of hydrogen electrode in contact with a solution whose pH is 10.**

**Ans.**  $\text{H}^+ + \text{e}^- \rightarrow 1/2\text{H}_2$   $n = 1$

$$E = E^\ominus - \frac{0.0591}{n} \log \frac{1}{[\text{H}^+]}$$

$$E = 0 - \frac{0.0591}{1} \times \text{pH}$$

$$E = -0.0591 \times 10 \text{ v}$$

$$E = -0.591/\text{V}$$

**Q. 8. If a current of 0.5 amp flows through a metallic wire for 2 hours, how many electrons would flow through the wire ?**


**Ans.**  $q = i \times t = 0.5 \times 2 \times 60 \times 60 = 3600 \text{ C}$

96500 Coulombs are equal to  $6.022 \times 10^{23} e^-$

$$\text{So, } 3600 \text{ Coulombs} = \frac{6.022 \times 10^{23}}{96500} \times 3600 = 2.246 \times 10^{22} \text{ electrons}$$

**Q.9. Calculate the electrode potential of a copper wire dipped in 0.1M CuSO<sub>4</sub> solution at 25°C. The standard electrode potential of copper is 0.34 Volt.**

**Ans.** The electrode reaction written as reduction potential is



$$E_{\text{Cu}^{2+}/\text{Cu}} = E_{\text{Cu}^{2+}/\text{Cu}}^0 - \frac{0.0591}{2} \log \frac{1}{[\text{Cu}^{2+}]} = 0.34 - \frac{0.0591}{2} \log \frac{1}{0.1} = 0.3104 \text{ V}$$

**Q.10. The conductivity of a 0.20M solution of KCl at 298K is 0.0248 S cm<sup>-1</sup>. Calculate molar conductivity.**

$$\begin{aligned} \text{Ans.} \quad \text{Molar conductivity} &= \frac{k \times 1000}{M} = \frac{0.0248 \text{ S cm}^{-1} \times 1000 \text{ cm}^3 \text{ L}^{-1}}{0.2 \text{ mol L}^{-1}} \\ &= 124.0 \text{ S cm}^2 \text{ mol}^{-1} \end{aligned}$$

**Q.11. Define conductivity and molar conductivity for a solution of an electrolyte.**

**Ans.** Conductivity is defined as ease with which current flows through electrolyte. It is reciprocal of specific resistance. Molar conductivity is conductance of all the ions produced by one mole of electrolyte when electrodes are at unit distance apart and have sufficient area of cross-section to hold electrolyte.


**Q.12. The resistance of a conductivity cell containing 0.001M KCl solution at 298 K is 1500 Ω. What is the cell constant if conductivity of 0.001M KCl solution at 298 K is 0.146 × 10<sup>-3</sup> S cm<sup>-1</sup> ?**

$$\begin{aligned} \text{Ans.} \quad \text{Cell constant} &= k \times R \\ &= 0.146 \times 10^{-3} \times 1500 \\ &= 0.219 \text{ cm}^{-1} \end{aligned}$$

**Q.13. Indicate the reaction which take place at cathode and anode in fuel cell.**




**Ans.** At cathode:  $\text{O}_2(\text{g}) + 2\text{H}_2\text{O} + 4e^- \rightarrow 4\text{OH}^-(\text{aq})$   
 At anode:  $2\text{H}_2(\text{g}) + 4\text{OH}^-(\text{aq}) \rightarrow 4\text{H}_2\text{O} + 4e^-$   
 The overall reaction is:  $2\text{H}_2(\text{g}) + \text{O}_2(\text{g}) \rightarrow 2\text{H}_2\text{O}(\text{l})$

**Q.14. The standard reduction potential for the Zn<sup>2+</sup> (aq)/Zn (s) half cell is – 0.76V. Write the reactions occurring at the electrodes when coupled with standard hydrogen electrode (SHE).**




**Q.15. What type of a battery is lead storage cell ? Write the anode and cathode reaction and overall reaction occurring in a lead storage battery during discharging and recharging cell.**

**Ans.** It is a secondary cell.



**Q.16. Draw a graph between  $\Lambda_m^\circ$  and  $\sqrt{C}$  for strong and weak electrolyte.**




**Q.17. Predict the products of electrolysis in each of the following:**


(a) An aqueous solution of  $\text{AgNO}_3$  with platinum electrodes.

(b) An aqueous solution of  $\text{CuCl}_2$  with Pt electrodes.


**Ans.** At Anode (Oxidation)




At cathode (Reduction)




(b) At anode (Oxidation)



At cathode (Reduction)



**Q.18. Determine the values of equilibrium constant  $K_c$  and  $\Delta G^\theta$  for the following reaction :**



**Ans.**

$$\Delta G^\theta = -nFE_{\text{cell}}^\theta$$

$$n = 2, E_{\text{cell}}^\theta = 1.05 \text{ V}$$

$$F = 96500 \text{ C mol}^{-1}$$

$$\Delta G^\theta = -2 \times 1.05 \times 96500$$

$$= -202.650 \text{ kJ}$$

$$\Delta G^\theta = -RT \ln K_c$$


$$\ln K_c = -\frac{\Delta G^\theta}{RT} = \frac{-202.650 \times 10^3}{8.314 \times 298}$$

$$K_c = 3.32 \times 10^{35}$$

**Q.19. The  $K_{sp}$  for AgCl at 298 K is  $1.0 \times 10^{-10}$ . Calculate the electrode potential for**

**Ag<sup>+</sup>/Ag electrode immersed in 1.0M KCl solution. Given  $E_{\text{Ag}^+/\text{Ag}}^\theta = 0.80 \text{ V}$ .**

**Ans.**



$$K_{sp} = [\text{Ag}^+][\text{Cl}^-]$$

$$[\text{Cl}^-] = 1.0 \text{ M}$$



$$[\text{Ag}^+] = \frac{K_{sp}}{[\text{Cl}^-]} = \frac{1 \times 10^{-10}}{1} = 1 \times 10^{-10} \text{ M}$$

Now,  $\text{Ag}^+ + e^- \rightarrow \text{Ag (s)}$

$$E = E^\theta - \frac{0.059}{1} \log \frac{1}{[\text{Ag}^+]} = 0.80 - \frac{0.059}{1} \log \frac{1}{10^{-10}}$$

$$= 0.80 - 0.059 \times 10 = 0.21 \text{ V}$$

**Q.20. Estimate the minimum potential difference needed to reduce  $\text{Al}_2\text{O}_3$  at  $500^\circ\text{C}$ .  
The free energy change for the decomposition reaction :**



$$n = \frac{6 \times 2}{3} = 4e^-$$

$$\Delta G = -nFE$$

$$\Delta G = 960 \times 10^3 \text{ J, } n = 4, F = 96500 \text{ C mol}^{-1}$$

$$960 \times 10^3 = -4 \times 96500 \times E$$

$$E = -2.487 \text{ V}$$


Minimum potential difference needed to reduce  $\text{Al}_2\text{O}_3 = -2.487 \text{ V}$ .

**Q.21. Two electrolytic cells containing silver nitrate solution and copper sulphate solution are connected in series. A steady current of 2.5 amp was passed through them till 1.078 g of Ag were deposited . How long did the current flow ? What weight of copper will be deposited ? ( $\text{Ag} = 107.8 \text{ u}$ ,  $\text{Cu} = 63.5 \text{ u}$ )**

**Ans.**  $W = z \times i \times t$

$$t = \frac{w}{z \times i}$$

$$t = \frac{1.078 \times 1 \times 96500}{107.8 \times 2.5} = 386 \text{ Seconds}$$



$$w = \frac{63.5}{2 \times 96500} \times 2.5 \times 386 = 0.3175 \text{ gram}$$

**Q.22. A solution of  $\text{Ni}(\text{NO}_3)_2$  is electrolysed between platinum electrodes using a current of 5.0 amp for 20 minutes . What mass of the nickle will be deposited at the cathode ? ( $\text{Ni} = 58.7 \text{ u}$ )**


**Ans.**  $W = z \times i \times t$

$$z = \frac{58.7}{2 \times 96500}$$

$$w = 1.825 \text{ gram}$$



**Q.23.** The cell in which the following reaction occurs :



Calculate the standard Gibbs energy and the equilibrium constant of the cell reaction.

**Ans.**

$$n = 2$$

$$\Delta G^{\circ} = -nFE^{\circ}_{\text{cell}} = -2 \times 96500 \times 0.236 \text{ J} = -45.55 \text{ kJ/mol}$$

$$\Delta G^{\circ} = -2.303 RT \log K_c$$

$$\log K_c = \frac{\Delta G^{\circ}}{-2.303RT} = \frac{45.55 \times 10^3}{2.303 \times 8.314 \times 298} = 7.983$$

$$K_c = \text{antilog}(7.983) = 9.616 \times 10^7$$

**Q.24.** The molar conductivity of  $0.025 \text{ mol L}^{-1}$  methanoic acid is  $46.1 \text{ S cm}^2 \text{ mol}^{-1}$ .


Calculate its degree of dissociation and dissociation constant. Given  $\Lambda^{\circ}(\text{H}^{+}) = 349.6 \text{ S cm}^2 \text{ mol}^{-1}$ ,  $\Lambda^{\circ}(\text{HCOO}^{-}) = 54.6 \text{ S cm}^2 \text{ mol}^{-1}$ .

**Ans.**

$$\Lambda^{\circ}_m(\text{HCOOH}) = \Lambda^{\circ}_m(\text{H}^{+}) + \Lambda^{\circ}_m(\text{HCOO}^{-})$$

$$= 349.6 + 54.6 \text{ S cm}^2 \text{ mol}^{-1} = 404.2 \text{ S cm}^2 \text{ mol}^{-1}$$

$$\Lambda^{\circ}_m = 46.1 \text{ S cm}^2 \text{ mol}^{-1}$$



$$\alpha = \frac{\Lambda^c_m}{\Lambda^{\circ}_m} = \frac{46.1}{404.2} = 0.114$$



Initial conc.

$$C \text{ mol L}^{-1}$$

$$0$$

$$0$$

At equil.


$$C(1 - \alpha)$$

$$C\alpha$$

$$C\alpha$$

$$K_a = \frac{C\alpha^2}{1 - \alpha} = \frac{0.025 \times (0.114)^2}{1 - 0.114} = 3.67 \times 10^{-4}$$

**Q.25.** Calculate the standard cell potentials of galvanic cells in which the following reaction take place:



Also calculate  $\Delta G^\circ$  and equilibrium constant of the reaction.

**Ans.**  $E_{\text{cell}}^0 = E_{\text{cathode}}^0 - E_{\text{anode}}^0$   
 $= 0.40 - (-0.74) = 0.34\text{V}$   
 $\Delta G^\circ = -nFE_{\text{cell}}^0 = -6 \times 96500 \times 0.34 = -196860$   
 $= -196860 \text{ J mol}^{-1} = -196.86 \text{ kJ/mol}$   
 $-\Delta G^\circ = 2.303RT \log K_c$   
 $196860 = 2.303 \times 8.314 \times 298 \log K_c$   
 or  
 $\log K_c = 34.5014$   
 $K_c = \text{antilog } 34.5014 = 3.193 \times 10^{34}$

**Q.26.** Calculate the potential of the following cell



**Given :**  $E_{\text{Sn}^{4+}/\text{Sn}^{2+}}^0 = 0.13\text{V}, E_{\text{Zn}^{2+}/\text{Zn}}^0 = -0.76\text{V}$

Will the cell potential increase or decrease, if the concentration of  $\text{Sn}^{4+}$  is increased?


**Ans.** 
$$E_{\text{cell}} = E_{\text{cell}}^0 - \frac{0.0591}{n} \log \frac{[\text{Sn}^{2+}][\text{Zn}^{2+}]}{[\text{Sn}^{4+}][\text{Zn}]}$$

$$= 0.89 - \frac{0.0591}{2} \log \frac{0.5 \times 2}{1.5 \times 1} = 0.89 - \frac{0.0591}{2} \log \frac{1}{1.5} = 0.895 \text{ V}$$

On increasing the concentration of  $\text{Sn}^{4+}$ , EMF of the cell will increase.

**Q.27.**  $E^\circ (\text{Cu}^{2+}/\text{Cu})$  and  $E^\circ (\text{Ag}^+/\text{Ag})$  is  $+0.337 \text{ V}$  and  $+0.799 \text{ V}$  respectively. Make a cell whose EMF is +ve. If the concentration of  $\text{Cu}^{2+}$  is  $0.01\text{M}$  and  $E_{\text{cell}}$  at  $25^\circ\text{C}$  is zero, calculate the concentration of  $\text{Ag}^+$ .


**Ans.** Cu is more reactive than silver, so that the cell is as  $\text{Cu}|\text{Cu}^{2+} (0.01\text{M}) \parallel \text{Ag}^+(\text{C})|\text{Ag}$  or cell reaction



$$\begin{aligned} E_{\text{cell}} &= E_{\text{cell}}^{\circ} - \frac{0.0591}{n} \log \frac{[\text{Cu}^{2+}][\text{Ag}]^2}{[\text{Cu}][\text{Ag}^+]^2} \\ &= E_{\text{cell}}^{\circ} - \frac{0.0591}{n} \log \frac{(0.01) \times 1^2}{1 \times [\text{Ag}^+]^2} \end{aligned}$$

$$\text{Or } [\text{Ag}^+] = 1.47 \times 10^{-9} \text{ M}$$

**Q.28. Calculate the potential of the cell at 298 K :**



**Given  $E^{\circ}$  for  $\text{Cd}^{2+}/\text{Cd} = -0.403 \text{ V}$ ,  $R = 8.314 \text{ J}^{-1} \text{ mol}^{-1}$ ,  $F = 96500 \text{ C mol}^{-1}$ .**

**Ans.** The cell reaction is  $\text{Cd} + 2\text{H}^+ (0.2\text{M}) \rightarrow \text{Cd}^{2+} (0.1\text{M}) + \text{H}_2 (0.5 \text{ atm})$

$$E_{\text{cell}}^{\circ} = 0 - (-0.403) = +0.403 \text{ V}$$

$$\begin{aligned} E_{\text{cell}} &= 0.403 - \frac{2.303RT}{nF} \log \frac{[\text{Cd}^{2+}] \times P_{\text{H}_2}}{[\text{Cd}][\text{H}^+]^2} \\ &= 0.403 - \frac{2.303 \times 8.314 \times 298}{2 \times 96500} \log \frac{0.1 \times 0.5}{(0.2)^2} \end{aligned}$$

$$E_{\text{cell}} = 0.403 - 0.003 = 0.40 \text{ V}$$

**Q.29. The electrical resistance of a column of 0.05M NaOH solution of diameter 1 cm and length 50 cm is  $5.55 \times 10^3 \text{ ohm}$ . Calculate its resistivity, conductivity and molar conductivity.**

**Ans.** Diameter = 1 cm, radius = 0.5 cm

$$\text{Area} = \pi r^2 = 3.14 \times (0.5)^2 = 0.785 \text{ cm}^2$$

$$\rho = \frac{R \times A}{l} = \frac{5.55 \times 10^3 \times 0.785}{50} = 87.135 \text{ ohm cm}$$

$$\text{Conductivity (k)} = \frac{1}{\rho} = \frac{1}{87.135} = 0.01148 \text{ ohm}^{-1} \text{ cm}^{-1} = 0.01148 \text{ ohm cm}$$

$$\text{Molar conductivity } \Lambda_m^c = \frac{k \times 1000}{M} = \frac{0.01148 \times 1000}{0.05} = 29.6 \text{ S cm}^2 \text{ mol}^{-1}$$

## LONG ANSWER TYPE QUESTIONS (5 Marks)

**Q. 1.** Conductivity of 0.00241M acetic acid is  $7.896 \times 10^{-5} \text{ S cm}^{-1}$ . Calculate its molar conductivity and if  $\Lambda_m^\circ$  for acetic acid is  $390.5 \text{ S cm}^2 \text{ mol}^{-1}$ , what is its dissociation constant ?

**Ans.**

$$\Lambda_m = \frac{k \times 1000}{M}$$

$$= \frac{7.896 \times 10^{-5} \text{ S cm}^{-1} \times 1000 \text{ cm}^3 \text{ L}^{-1}}{0.00241 \text{ mol L}^{-1}} = 32.76 \text{ S cm}^2 \text{ mol}^{-1}$$

$$\alpha = \frac{\Lambda_m}{\Lambda_m^\circ} = \frac{32.76}{390.5} = 8.39 \times 10^{-2}$$

$$K_a = \frac{C\alpha^2}{1-\alpha} = \frac{0.00241 \times (8.39 \times 10^{-2})^3}{1 - 8.39 \times 10^{-2}} = 1.86 \times 10^{-5}$$

**Q.2.** Three electrolytic cells A, B, C containing solution of  $\text{ZnSO}_4$ ,  $\text{AgNO}_3$  and  $\text{CuSO}_4$  respectively all connected in series. A Steady current of 1.5 amperes was passed through then until 1.45g of silver deposited at the cathode of cell B How long did the current flow? What mass of copper and of zinc were deposited ?

**Ans.** 108 g of silver is deposited by  $\text{Ag}^+ + \text{e}^- \rightarrow \text{Ag}_{(s)}$  is 96500C.

$$1.45 \text{ g silver is deposited by} = \frac{96500 \times 1.45}{108}$$


$$= 1295.6 \text{ C}$$

$$Q = I \times t$$

$$1295.6 = 1.5 \times t$$


$$t = \frac{1295.6}{1.5} = 863 \text{ s}$$

In cell A, the electrode reaction is



2F of electricity deposit Zn = 65.3 g


In cell A, the electrode reaction is



2F of electricity deposit Zn = 65.3g

$$1295.6 \text{ of electricity deposit Zn} = \frac{65.3 \times 1295.6}{2 \times 96500} \\ = 0.438 \text{ g}$$

In cell C, the electrode reaction is



2F of electricity deposit Cu = 63.5g

$$1295.6 \text{ of electricity deposit Cu} = \frac{63.5 \times 1295.6}{2 \times 96500} \\ = 0.426 \text{ g}$$

**Q. 3. (a) State Kohlraush's law.**

**(b) Suggest a way to determine the  $\Lambda_m^\circ$  for  $\text{CH}_3\text{COOH}$ .**

**(c) The  $\Lambda_m^\circ$  for sodium acetate, HCl, NaCl are 91.0, 425.9 and 126.4  $\text{S cm}^2 \text{mol}^{-1}$  respectively at 298 K. Calculate  $\Lambda_m^\circ$  for  $\text{CH}_3\text{COOH}$ .**

**Ans. (a)** The molar conductivity at a infinite dilution for a given salt can be expressed as the sum of the individual contribution from the ions of electrolyte.

**(b)**  $\Lambda^\circ \text{CH}_3\text{COOH} = ?$

$$\begin{aligned} \lambda^\circ \text{CH}_3\text{COO}^- + \lambda^\circ \text{H}^+ &= \lambda^\circ \text{CH}_3\text{COO}^- + \lambda^\circ \text{Na}^+ + \lambda^\circ \text{H}^+ \\ &\quad + \lambda^\circ \text{Cl}^- - \lambda^\circ \text{Na}^+ - \lambda^\circ \text{Cl}^- \quad \dots(i) \end{aligned}$$

$$\Lambda_m^\circ \text{CH}_3\text{COOH} = \Lambda^\circ \text{CH}_3\text{COONa} + \Lambda^\circ \text{HCl} - \Lambda^\circ \text{NaCl}$$

$$\begin{aligned} \text{(c)} \quad \Lambda_m^\circ \text{CH}_3\text{COOH} &= \Lambda^\circ \text{CH}_3\text{COONa} + \Lambda^\circ \text{HCl} - \Lambda^\circ \text{NaCl} \\ &= 91.0 + 425.9 - 126.4 \\ &= 390.5 \text{ S cm}^2 \text{mol}^{-1} \end{aligned}$$


**Q.4. (a) Define weak and strong electrolytes**

**(b) The  $E^\circ$  values corresponding to the following two reduction electrode processes are:**

**(i)  $\text{Cu}^+/\text{Cu} = 0.52\text{V}$  (ii)  $\text{Cu}^{2+}/\text{Cu} = 0.16\text{V}$**

**Formulate the galvanic cell for their combination . Calculate the cell potential and  $\Delta G^\circ$  for the cell reaction.**

**Ans. (a)** Weak electrolyte: the substance which partially ionized in solution is known as weak electrolyte. Example:  $\text{NH}_4\text{OH}$   
Strong electrolyte: The substance which completely ionized in solution is known as strong electrolyte. Example:  $\text{NaCl}$ .



Q.5. Calculate emf and  $\Delta G^\circ$  for the following cell at 298K.



[Given :  $E^\circ \text{Mg}^{2+}/\text{Mg} = -2.36 \text{ V}$ ;  $E^\circ \text{Cu}^{2+}/\text{Cu} = 0.34 \text{ V}$ ;  $1 \text{ F} = 96500 \text{ C mol}^{-1}$ ]

Ans.  $E_{\text{cell}} = E^\circ_{\text{cell}} - \frac{0.059}{2} \log \frac{[\text{Mg}^{2+}]}{[\text{Cu}^{2+}]}$   
 $E_{\text{cell}} = 0.34 - (-2.36) - \frac{0.059}{2} \log \frac{10^{-3}}{10^{-4}}$   
 $= 2.70 - 0.02655 = 2.67\text{V}$   
 $\Delta G^\circ = -nF E^\circ_{\text{cell}} = -2 \times 96500 \times 2.71$   
 $= -5.23 \times 10^2 \text{ kJ mol}^{-1}$

Q.6. (a) Give the units of conductivity and molar conductivity

(b) Write down Nernst equation and calculate the emf of the following cell at 298 K:




Given:  $E^\circ (\text{Cu}^{2+}/\text{Cu}) = +0.34 \text{ V}$  and  $E^\circ (\text{Ag}^+/\text{Ag}) = +0.80 \text{ V}$ .

Hint: (a) Conductivity  $\text{Scm}^{-1} \text{ cm}$ , Molar conductivity  $\text{Scm}^{-2} \text{ mol}^{-1}$

(b)  $Q = \frac{[\text{Cu}^{2+}][\text{Ag}^+]^2}{[\text{Cu}][\text{Ag}^+]^2} = \frac{0.13 \times 10^{-8}}{1 \times (10^{-4})^2} = 0.13 \times 10^8$

$E = E^\circ - \frac{0.0591}{n} \log_{10} Q$   
 $= 0.46 - \frac{0.0591}{2} \log_{10}(0.13 \times 10^8) = 0.25\text{V}$

Q.7. (a) Calculate the emf of the following concentration cell:



(b) How can the reduction potential of an electrode be increased ?

Hint: (a)

$E = E^\circ - \frac{0.0591}{n} \log_{10} Q$   
 $= 0 - \frac{0.0591}{n} \log 0.1 = 0.0295 \text{ Volt}$



$E_{\text{M}^+/\text{M}} = E^\circ_{\text{M}^{n+}/\text{M}}$

$\frac{2.303 RT}{nF} \log \frac{1}{[\text{M}^{n+}]}$

or

$E_{\text{M}^+/\text{M}} = E^\circ_{\text{M}^{n+}/\text{M}}$

$\frac{2.303 RT}{nF} \log [\text{M}^{n+}]$

From the above relation it is clear that the reduction potential can be increased either by increasing temperature or by increasing the concentration of metal ion.

- Q.8.** (i) The conductivity of 0.02M solution of NaCl is  $2.6 \times 10^{-2} \text{ S cm}^{-1}$ .  
What is its molar conductivity ?  
(ii) Give reasons:  
(a) Rusting of iron pipe can be prevented by joining it with a piece of magnesium.  
(b) Dry cell become dead after a long time, even if it has not been used?

**Ans.**

$$K = 2.6 \times 10^{-2} \text{ S cm}^{-1}$$

$$C = 0.02 \text{ M}$$

$$\begin{aligned} \Lambda_m &= \frac{k \times 1000}{C(M)} \\ &= \frac{2.6 \times 10^{-2} \times 1000}{0.02} \\ &= \frac{26 \times 100}{0.02 \times 100} = \frac{26 \times 10^2}{2} \\ &= 13 \times 10^2 \text{ S cm mol}^{-1} \end{aligned}$$

- (ii) (a) It is due to cathodic protection in which magnesium metal is oxidised in preference to iron and acts as the anode.  
(b) A dry cell becomes dead after a long time because the acidic  $\text{NH}_4\text{Cl}$  corrodes with the zinc container of dry cell.

- Q.9.** (i) Depict the galvanic cell in which the reaction  
 $\text{Zn(s)} + 2\text{Ag}^+ \rightarrow \text{Zn}^{2+} + 2\text{Ag(s)}$   
take place. Further show :  
(a) Which of the electrode is negatively charged?  
(b) The carriers of the current in the cell.  
(c) Individual reaction at each electrode?  
(ii) A solution of  $\text{CuSO}_4$  is electrolysed for 10 mins. With a current of 1.5 amperes. What is the mass of copper deposited at the cathode ?

- Ans.** (i)  $\text{Zn(s)} | \text{Zn}^{2+}(\text{aq}) || \text{Ag}^+(\text{aq}) | \text{Ag(s)}$   
(a) Zn electrode (anode)  
(b) Ions are carriers of the current in the cell.  
(c) At anode:  
 $\text{Zn(s)} \rightarrow \text{Zn}^{2+} + 2\text{e}^-$   
At cathode:  
 $\text{Ag}^+ + \text{e}^- \rightarrow \text{Ag(s)}$


- (ii)  $I = 1.5 \text{ Ampere}$   
Time:  $10 \times 60 \text{ s} = 600 \text{ s}$   
 $Q = I \times t$   
 $= 1.5 \times 600 = 900 \text{ C}$   
 $\text{Cu}^{2+} + 2\text{e}^- \rightarrow \text{Cu(s)}$

$$\begin{aligned} 2\text{F amount of electricity deposit copper} &= 63.5 \text{ g} \\ 900 \text{ C amount of electricity deposit copper} &= \frac{63.5 \times 900}{2 \times 96500} = 0.296 \text{ g} \end{aligned}$$

### CASE STUDY BASED QUESTIONS

1. **Read the passage given below and answer the questions that follow:**

The study of the conductivity of electrolyte solution is important for the development of electrochemical devices, for the characterisation of the dissociation equilibrium of weak electrolytes and for the understanding of charge transport by ions. The conductivity of electrolyte is measured for electrolyte solution with concentration in the range of  $10^{-3}$  to  $10^{-4}$  mol/L, as solution in this concentration range can be easily prepared. The variation in molar conductivity ( $\Lambda_m$ ) of strong electrolyte with concentration is given by equation  $\Lambda_m = \Lambda_m^0 - A\sqrt{C}$ . Where  $\Lambda_m^0$  is the molar conductivity at infinite dilution and  $C$  is the concentration for solution. Following graph shows the, variation of molar conductivity with concentration for both weak and strong electrolytes.





Limiting molar conductivity cannot be determined by extrapolation of  $\Lambda_m$  versus  $\sqrt{C}$  curve. Molar conductivity at infinite dilution can be calculated by sum of contributions of each ion.  $\Lambda_m^0 = \nu^+ \lambda_0^+ + \nu^- \lambda_0^-$

Where  $\lambda_0^+$  and  $\lambda_0^-$  are the limiting ionic conductivities of positive and negative ions respectively and  $\nu^+$  and  $\nu^-$  are their stoichiometric coefficients in the salt molecular formula.

**In the following questions a statement of assertion followed by a statement of reason is given. Choose the correct answer out of the following choices.**

- a) Assertion and reason both are correct statements and reason is correct explanation for assertion.
  - b) Assertion and reason both are correct statements but reason is not correct explanation for assertion.
  - c) Assertion is correct statement but reason is wrong statement.
  - d) Assertion is wrong statement but reason is correct statement.
- (A) **ASSERTION:** For  $\text{CH}_3\text{COOH}$   $\Lambda_m^0$  cannot be determined experimentally.  
**REASON:**  $\text{CH}_3\text{COOH}$  is a weak acid and Debye Huckel Onsager equation cannot be used. Extrapolation method cannot be employed.
- (B) **ASSERTION :** The ratio of conductivity to the observed conductance does not depend upon the concentration of the solution taken in the conductivity cell.  
**REASON :** Conductivity of solution decreases with dilution whereas observed conductance increases with dilution.
- (C) **ASSERTION :** Molar conductance of all electrolytes decrease with increasing concentration.  
**REASON :** Lesser no. of ions are available per unit volume of solution at higher concentration.
- (D) **ASSERTION :**  $0.1 \text{ M NH}_4\text{OH}$  at  $25^\circ\text{C}$  has lesser conductance than at  $50^\circ\text{C}$ .  
**REASON :** Conductance of a weak electrolyte decreases with increase in temperature.

2. **Read the passage given below and answer the questions that follow:**

In the Daniell cell, the copper electrode is the anode. The electrons leave the cell from the zinc and enter into the copper electrode. To complete the circuit a salt bridge (an inverted U-tube) is used. Salt bridge consists of a concentrated solution of agar-agar + KCl/ KNO<sub>3</sub>/NH<sub>4</sub>NO<sub>3</sub>. The mobility of cations and anions are the same. The maximum electrical work is given by  $W_{e, \max} = \Delta G$ . This  $\Delta G$  is related to emf as  $\Delta G = -nFE$ . The extent of reaction is measured by  $\Delta G$ . The emf of a cell is determined by the Nernst equation,

$$E = E^0 - 0.059 \log Q.$$

The Nernst equation is also used to calculate the emf of concentration cell.

$M/M^+_{(aq)} \parallel M^+_{(ar)}/M$ . When a given cell is at equilibrium,  $Q = K_{eq}$ .

To calculate the standard electrode potential of a half cell like Ag/Ag<sup>+</sup>, Cu/Cu<sup>2+</sup>, one has to complete it with SHE e.g.

Pt/H<sub>2</sub>(g) /H<sup>+</sup>(aq) its  $E^0 = 0$  (by convention). From the emf study, we can calculate  $E^0$ , pH, valency,  $K_{eq}$ ,  $K_s$ , thermodynamic parameters, etc.

(A) **An electrochemical cell stops working after some time because**

- (a) Electrode potential of both the electrodes becomes zero.
- (b) Electrode potential of both the electrodes becomes equal.
- (c) One of the electrode is eaten away.
- (d) The reaction starts proceeding in opposite direction.

(B) **Which of the following statements is correct for a galvanic cell?**

- (a) Reduction occurs at cathode. (b) Oxidation occurs at anode.
- (c) Electrons flow from anode to cathode. (d) All statements are correct.

(C) **What is correct when net cell reaction is spontaneous?**

- (a)  $E^0_{cell}$  is negative (b)  $E_{cell} > 0$
- (c)  $E_{cell} = E^0_{cell}$  (d)  $\Delta G < 0$


(D) **The function of salt bridge is to:**

- (a) allow ions to move from anode to cathode
- (b) allow solutions from one half cell to the other half cell
- (c) allow the current to flow through the cell and keep the solutions electrically neutral
- (d) keep the level of solutions same.

**3. Read the passage given below and answer the questions that follow:**

Oxidation-reduction reactions are commonly known as redox reactions. They involve transfer of electrons from one species to another. In a spontaneous reaction, energy is released which can be used to do useful work. The reaction is split into two half reactions. Two different containers are used and a wire is used to drive the electrons from one side to the other and a Voltaic/Galvanic cell is created. It is an electrochemical cell that uses spontaneous redox reactions to generate electricity. A salt bridge also connects to the half cells. The reading of the voltmeter gives the cell voltage or cell potential or electromotive force. If  $E^{\circ}_{\text{cell}}$  is positive the reaction is spontaneous and if it is negative the reaction is non-spontaneous and is referred to as electrolytic cell. Electrolysis refers to the decomposition of a substance by an electric current. One mole of electric charge when passed through a cell will discharge half a mole of a divalent metal ion such as  $\text{Cu}^{2+}$ . This was first formulated by Faraday in the form of laws of electrolysis.

The conductance of material is the property of materials due to which a material allows the flow of ions through itself and thus conducts electricity. Conductivity is represented by  $k$  and it depends upon nature and concentration of electrolyte, temperature etc. A more common term molar conductivity of a solution at a given concentration is conductance of the volume of solution containing one mole of electrolyte kept between two electrodes with the unit area of cross-section and distance of unit length. Limiting molar conductivity of weak electrolytes cannot be obtained graphically.



- (A) Is silver plate the anode or cathode?  
 (B) What will happen if the salt bridge is removed?  
 (C) When does electrochemical cell behaves like an electrolytic cell?  
 (D) (i) What will happen to the concentration of  $\text{Zn}^{2+}$  and  $\text{Ag}^{+}$  when  $E_{\text{cell}}=0$   
 (ii) Why does conductivity of a solution decreases with dilution?

**OR**

- (D) The molar conductivity of a 1.5 M solution of an electrolyte is found to be  $138.9 \text{ S cm}^2 \text{ mol}^{-1}$ . Calculate the conductivity of this solution.

## ANSWERS

### I MULTIPLE CHOICE QUESTIONS

1. d 2. c 3. d 4. c 5. a 6. c 7. b 8. b 9. a 10. a, d  
11. a 12. a, b 13. a 14. a 15. a, b 16. a 17. c 18. d 19. c 20. b, d

### II FILL IN THE BLANKS

1. Conductivity 2.  $c^{1/2}$  3. Anode to cathode  
4. ohm metre 5. positive 6. decreases  
7.  $m^{-1}$  8. cannot 9. galvanisation  
10. 4F

### III ASSERTION REASON TYPE QUESTIONS

1. a 2. a 3. a 4. d 5. c.  
6. d 7. a 8. a 9. a 10. a

### IV ONE WORD ANSWER TYPE QUESTIONS

1. 3F 2. Increases 3. ions  
4. Standard hydrogen electrode 5. No 6. Conductivity  
7. increases 8. Faraday constant  
9. Calomel electrode 10. increases 11. Fuel Cell  
12. At equilibrium 13.  $E^0_{\text{cell}} = \frac{0.0591}{n} \log K_c$   
14. From Zinc to Silver 15. Equilibrium state is attained

### CASE STUDY TYPE QUESTIONS

PASSAGE: 1: (A) a (B) b (C) c (D) c

PASSAGE: 2: (A) b (B) d (C) b, d (D) c

PASSAGE: 3: (A) Cathode (B) Voltage will drop to zero (C) When  $E_{\text{ext}} > E_{\text{cell}}$

(D) (i) Increasing concentration of  $Zn^{2+}$  and decreasing concentration of  $Ag^+$  reaches to an equilibrium (ii) Due to decrease in no. of ions per unit volume.

OR

$$\Lambda^0_m = K \times \frac{1000}{M}$$

$$138.9 = K \times \frac{1000}{1.5}$$

$$K = 0.208 \text{ ohm}^{-1} \text{ cm}^{-1}$$

**UNIT TEST-1**  
**CHAPTER-3**  
**ELECTROCHEMISTRY**

**TIME ALLOWED :1 HR.****M.M. 20**

1. What does the negative sign in the expression  $E^0_{\text{zn}^{2+}/\text{zn}} = -0.76\text{V}$  mean? 1
2. Write unit of molar conductivity. 1
3. Suggest a way to determine the  $\Lambda_m^0$  value of water. 1
4. Write the nernst equation of the following cell  
 $\text{Mg(s)}/\text{Mg}^{2+}(0.001\text{M}) \parallel \text{Cu}^{2+}(0.001\text{M})/\text{Cu(s)}$  1
5. Why is it not possible to measure single electrode potential? 1
6. Calculate emf of the following cell 2  
 $\text{Cd}/\text{Cd}^{2+}(0.10\text{M}) \parallel \text{H}^+(0.20\text{M})/\text{H}_2(0.5 \text{ atm})/\text{Pt}$   
 (Given  $E^0$  for  $\text{Cd}^{2+}/\text{Cd} = -0.403\text{V}$ )
7. Why on dilution  $\Lambda_m$  of  $\text{CH}_3\text{COOH}$  increases drastically while that of  $\text{CH}_3\text{COONa}$  increases gradually? 2
8. Conductivity of  $2.5 \times 10^{-4} \text{ M}$  methanoic acid ( $\text{HCOOH}$ ) is  $5.25 \times 10^{-5} \text{ Scm}^{-1}$ .  
 Calculate its molar conductivity and degree of dissociation. 3  
 Given:  $\lambda^0(\text{H}^+) = 349.5 \text{ Scm}^2\text{mol}^{-1}$  and  $\lambda^0(\text{HCOO}^-) = 50.5 \text{ Scm}^2\text{mol}^{-1}$ .
9. (i) The conductivity of an aqueous solution of  $\text{NaCl}$  in a cell is  $92 \text{ ohm}^{-1}$ , the  
 resistance offered by the cell is  $247.8 \text{ ohm}$ . Calculate the cell constant. 3  
 (ii) What is the effect of dilution on the conductivity of an electrolytic solution?
10. Calculate EMF and  $\Delta G$  for the following cell at  $298\text{K}$ :  
 $\text{Mg(s)}/\text{Mg}^{2+}(0.01\text{M}) \parallel \text{Ag}^+(0.0001\text{M})/\text{Ag(s)}$   
 Given :  $E^0_{\text{Mg}^{2+}/\text{Mg}} = -2.37\text{V}$ ,  $E^0_{\text{Ag}^+/\text{Ag}} = +0.80\text{V}$  5

## UNIT TEST-2

## CHAPTER-3

## ELECTROCHEMISTRY

TIME ALLOWED :1 HR.

M.M. 20

1. Express the relation between conductivity and molar conductivity of the solution. 1
2. Name any two metals which can be used for cathodic protection of iron. 1
3. Name a battery used in Apollo space programme. 1
4. Write the correct representation of the cell: 2
  2.  $\text{Cr}_{(s)} + 3\text{Cd}^{2+}_{(aq)} \longrightarrow \text{Cr}^{3+}_{(aq)} + 3\text{Cd}_{(s)}$
5. How many Faradays of charge are required to convert 1 mole of  $\text{Fe}^{2+}$  to Fe? 1
6. How does molar conductivity vary with dilution for 2
  - (i) weak electrolyte and for
  - (ii) strong electrolyte? Give reasons for these variations.
7. The conductivity of 0.2 M solution of KCl at 298 K is  $0.025 \text{ S cm}^{-1}$ . Calculate the molar conductivity. 2
8. Account for the following: 2
  - (i) Alkaline medium inhibits the rusting of iron.
  - (ii) Iron does not rust even if the zinc coating is broken in a galvanised iron pipe.
9. Calculate the emf for the given cell at  $25^\circ\text{C}$ . 3
 

$\text{Cr} \mid \text{Cr}^{3+}(0.1\text{M}) \parallel \text{Fe}^{2+}(0.01\text{M}) \mid \text{Fe}$

[Given:  $E^\circ \text{Cr}^{3+}/\text{Cr} = -0.74 \text{ V}$ ,  $E^\circ \text{Fe}^{2+}/\text{Fe} = -0.44 \text{ V}$ ]
- 10.(a) Write the cell reactions which occur in lead storage battery 3
  - (i) When the battery is in use and
  - (ii) When the battery is on charging.
  - (b) Mention two advantages of fuel cells.
11. Molar conductivities at infinite dilution for  $\text{NH}_4\text{Cl}$ ,  $\text{NaOH}$  and  $\text{NaCl}$  solutions at 298 K are respectively 129.8, 217.4 and  $108.9 \text{ S cm}^2 \text{ mol}^{-1}$  and the molar conductivity of a  $10^{-2} \text{ M}$  solution of  $\text{NH}_4\text{OH}$  is  $9.33 \text{ S cm}^2 \text{ mol}^{-1}$ . Calculate the degree of dissociation (a) of  $\text{NH}_4\text{OH}$  in the above mentioned solution. 3

## UNIT 4

## Chemical Kinetics

### Points to Remember

#### RATE OF REACTION:-

For a reaction  $R \rightarrow P$

Rate of reaction = change of conc. of R or P / Time interval

**Rate =  $-\Delta[R]/\Delta t = \Delta[P]/\Delta t$ , This is average rate of reaction.**

For expressing the rate of such a reaction where stoichiometric coefficients of reactants or products are not equal to one, rate of disappearance of any of the reactants or the rate of appearance of products is divided by their respective stoichiometric coefficients.

For e.g. for a chemical reaction:  $N_2(g) + 3H_2(g) \rightarrow 2NH_3(g)$

$$r_{\text{avg}} = -\Delta[N_2]/\Delta t = -1/3 (\Delta[H_2]/\Delta t) = 1/2 (\Delta[NH_3]/\Delta t)$$

#### Units of rate of a reaction:-

Concentration time<sup>-1</sup>

- if concentration is in mol L<sup>-1</sup> and time is in seconds then the units will be mol L<sup>-1</sup> s<sup>-1</sup>.
- In gaseous reaction, when the concentration of the gases is expressed in terms of their partial pressure, then the units will be atm s<sup>-1</sup>.

**INSTANTANEOUS RATE** is defined as the rate of change in concentration of any one of reactant or product at a particular instant of time.


$$\text{when } \Delta t \rightarrow 0; r_{\text{inst}} = -d[R]/dt = d[P]/dt$$

#### Factors affecting rate of a reaction:

- Nature of the reacting species:** Chemical reaction is a process in which new
- Concentration of reactants:** The rate of reaction increases with increase in concentration of reactants.
- Effect of temperature:** The rate of reaction is nearly double for every 10°C rise in temperature.
- Catalyst:** generally catalyst increases the rate of reaction.
- Effect of radiations:** The rates of some reactions are enhanced due to absorption of radiation. These reactions are called photochemical reactions.

**LAW OF MASS ACTION:**

The rate of a chemical reaction is directly proportional to the product of the molar concentrations of the reactants.



According to law of mass action

$$\text{Rate} \propto [A]^a [B]^b,$$

$$\text{Rate} = k[A]^a [B]^b$$

- **RATE CONSTANT OF A REACTION**

at a given temperature may be defined as rate of the reaction when the molar concentration of each of the reactants is unity.

- **CHARACTERISTICS OF RATE CONSTANT**

- Rate constant is a measure of the rate of the reaction.
- Larger the value of  $k$ , faster is the reaction.
- Different reactions have different values of  $k$ .
- For a particular reaction, the rate constant is independent of concentration.
- At a particular temperature, the value of  $k$  is constant. However, it changes with temperature.

- **RATE LAW** is the expression in which reaction rate is given in terms of molar concentration of reactants with each term raised to some power, which may or may not be same as the stoichiometric coefficient of the reacting species in a balanced chemical equation. It is determined experimentally.

- **ORDER OF A REACTION:** The sum of powers of the concentration of the reactants in the rate law expression is called the order of the chemical reaction.

For the rate law expression

$$\text{Rate} = k[A]^x [B]^y$$

$$\text{Order} = x + y$$

- **UNITS OF RATE CONSTANTS:** Units of rate constant are different from reaction of different order :  $(\text{mol L}^{-1})^{1-n} \text{ time}^{-1}$  where 'n' is order of reaction. For gas phase reaction unit of rate constant is  $(\text{atm or bar})^{1-n} \text{ s}^{-1}$

Reaction	Order ( $\alpha + \beta$ )	Units of rate constant
Zero order reaction	0	$\frac{\text{mol L}^{-1}}{\text{s}} \times \frac{1}{(\text{mol L}^{-1})^0} = \text{mol L}^{-1} \text{ s}^{-1}$
First order reaction	1	$\frac{\text{mol L}^{-1}}{\text{s}} \times \frac{1}{(\text{mol L}^{-1})^1} = \text{s}^{-1}$
Second order reaction	2	$\frac{\text{mol L}^{-1}}{\text{s}} \times \frac{1}{(\text{mol L}^{-1})^2} = \text{mol}^{-1} \text{ L}^1 \text{ s}^{-1}$




**MECHANISM AND RATE LAW:**

The reactions taking place in one step are called **elementary reactions**. When a sequence of elementary reactions, (called mechanism) gives us the products, the reactions are called **Complex reactions**. In complex reactions, the rate of the reaction is determined by the slowest step in the sequence. The slowest step is called **rate determining step** in the proposed mechanism.


**ZERO ORDER REACTION**

Integrated rate equation  $k = [R]_0 - [R]/t$

**INTEGRATED RATE EQUATION FOR FIRST ORDER REACTION:**

$$k = \frac{2.303}{t} \log \frac{[R]_0}{[R]}$$

where  $[R]_0$  is initial concentration of reactants and  $[R]$  is concentration at time  $t$ .



For a typical first order gas phase reaction:  $A(g) \rightarrow B(g) + C(g)$

$$k = \frac{2.303}{t} \log \frac{P_i}{(2P_t - P_i)}$$

Where  $p_i$  is the initial pressure of A and  $P_t$  the total pressure at time 't'.

- **HALF-LIFE PERIOD** ( $t_{1/2}$ ): The half-life of a reaction is the time in which the concentration of a reactant is reduced to one half of its initial concentration.

For zero order reaction :  $t_{1/2} = [R]_0 / 2k$

For first order reaction :  $t_{1/2} = 0.693/k$

**For zero order reaction  $t_{1/2} \propto [R]_0$ . For first order reaction  $t_{1/2}$  is independent of  $[R]_0$ .**

- **PSEUDO FIRST ORDER REACTIONS.** Chemical reactions which are not truly of the first order but under certain conditions become first order reactions are called e.g. A bimolecular reaction, in which one reactant is present in large excess and rate of reaction is independent of its concentration, the reaction follows first order kinetics.

For example, ester hydrolysis, where water is taken in excess.

$\text{CH}_3\text{COOCH}_2\text{CH}_3 + \text{H}_2\text{O} \rightarrow \text{CH}_3\text{COOH} + \text{CH}_3\text{CH}_2\text{OH}$  is a bimolecular but first order reaction.

- **Activation energy :** The minimum extra amount of energy absorbed by reactant molecules so that their energy becomes equal to the threshold energy is called activation energy.

Activation energy = Threshold energy - Kinetic energy

- **Temperature coefficient:** the ratio of rate constant at two temperatures having difference of 10°C is called temperature coefficient.

Temperature coefficient = Rate constant at  $T + 10^\circ\text{C}$  / Rate constant at  $T^\circ\text{C}$

- **Arrhenius Equation:**

$$K = Ae^{-E_a/RT}$$

where,  $K$  = Rate constant

$A$  = Arrhenius energy (Frequency factor or pre-exponential factor)

$E_a$  = Activation energy

$R$  = Rate constant

$T$  = Temperature

= Fraction of molecules having energy equal to or more than activation energy

$$\log K = \log A - \frac{E_a}{RT}$$

$$\log \frac{K_2}{K_1} = \frac{E_a}{2.303R} \left[ \frac{1}{T_1} - \frac{1}{T_2} \right]$$

$$\log \frac{K_2}{K_1} = \frac{E_a}{2.303 \times 8.314} \left[ \frac{T_2}{T_2} - \frac{T_1}{T_2} \right] \quad \text{Where } R = 8.314 \text{ JK}^{-1} \text{ mol}^{-1}$$

## OBJECTIVE TYPE QUESTIONS

### I MULTIPLE CHOICE QUESTIONS

1. **The initial concentration of the reactant is doubled, the time for half reaction is also doubled. Then the order of the reaction is**  
 (a) Zero      (b) one      (c) Fraction      (d) none
2. **Which of the following statements is correct?**  
 (a) The rate of a reaction decreases with passage of time as the concentration of reactants decreases  
 (b) The rate of a reaction is same at any time during the reaction  
 (c) The rate of a reaction is independent of temperature change  
 (d) The rate of a reaction decreases with increase in concentration of reactants(s)
3. **The rate constant of a reaction is  $5.8 \times 10^{-21} \text{ s}^{-1}$ . The order of the reaction is.**  
 (a) First order    (b) zero order    (c) Second order    (d) Third order
4. **A second order reaction between A and B is elementary reaction:  $A+B \rightarrow \text{Product}$  rate law expression of this reaction will be:**  
 (a)  $\text{Rate} = k[A][B]$     (b)  $\text{Rate} = k[A]^0[B]^2$     (c)  $\text{Rate} = k[A]^2[B]^0$     (d)  $\text{Rate} = k[A]^{3/2}[B]^{1/2}$
5. **Which of the following is pseudo first order reaction?**  
 (a)  $2\text{H}_2\text{O}_2 \rightarrow 2\text{H}_2\text{O} + \text{O}_2$   
 (b)  $2\text{O}_3 \rightarrow 3\text{O}_2$   
 (c)  $\text{CH}_3\text{COOC}_2\text{H}_5 + \text{NaOH} \rightarrow \text{CH}_3\text{COONa} + \text{C}_2\text{H}_5\text{OH}$   
 (d)  $\text{CH}_3\text{COOC}_2\text{H}_5 + \text{H}_2\text{O} \rightarrow \text{CH}_3\text{COOH} + \text{C}_2\text{H}_5\text{OH}$
6. **A large increase in the rate of reaction for rise in temperature is due to:**  
 (a) Increase in the number of collisions  
 (b) Increase in the number of activated molecules  
 (c) Lowering of activation energy  
 (d) Shortening of the mean free path.

**7. Radioactive decay is an example of:**

- (a) first order (b) zero order (c) second order (d) 0.5 order

**8. For a zero order reaction, the plot of concentration of reactant vs time is (intercept refers to concentration axis)**

- (a) linear with +ve slope and zero intercept  
(b) linear with -ve slope and zero intercept  
(c) linear with -ve slope and non-zero intercept  
(d) linear with positive slope and non-zero intercept

**9. The rate constant of nth order has units**

- (a)  $\text{litre}^{1-n} \text{mol}^n \text{s}^{-1}$  (b)  $\text{mol}^{1-n} \text{litre}^{1-n} \text{s}^{-1}$   
(c)  $\text{mol}^{1-n} \text{litre}^n \text{s}^{-1}$  (d)  $\text{mol}^{1-n} \text{litre}^{n-1} \text{s}^{-1}$

**10. A hypothetical reaction  $A_2 + B_2 \rightarrow 2AB$  follows the mechanism as given below:**



**The order of reaction is:**

- (a) 2 (b) 0 (c)  $1\frac{1}{2}$  (d) 1

**11. In a first order, the concentration of the reactant reduced to 1/4 in 60 minutes. What will be its half life?**

- (a) 120 min (b) 40 min  
(c) 30 min (d) 25 min

**12. For a complex reaction \_\_\_\_\_**

- (a) order of overall reaction is same as molecularity of the slowest step.  
(b) order of overall reaction is less than the molecularity of the slowest step.  
(c) order of overall reaction is greater than molecularity of the slowest step.  
(d) molecularity of the slowest step is never zero or non integer.

**13. Which of the reaction ends in infinite time?**

- (a) Zero order (b) First order  
(c) Second order (d) Third order

**14. Which one is correct for first order reaction.**

- (a)  $t_{75\%}/t_{50\%}=1.5$  (b)  $t_{75\%}/t_{50\%}=2$  (c)  $t_{99.9\%}/t_{50\%}=10$  (d)  $t_{99.9\%}/t_{50\%}=2$

**15. The rate constant of a zero order reaction is :**

- (a) Independent upon initial concentration
- (b) Inversely proportional to concentration
- (c) Does not depend upon concentration
- (d) None of these

**16. For the elementary reaction  $M \rightarrow N$ , the rate of disappearance of M increases by a factor of 8 upon doubling the concentration of M. The order of the reaction with respect to M is:**

- (a) 4      (b) 3      (c) 2      (d) 1

**17. When initial concentration of a reactant is doubled in a reaction, its half life period is not affected. The order of the reaction is.**

- (a) Second
- (b) more than zero but less than first
- (c) Zero
- (d) First

**18. Which of the following influences the rate of reaction?**

- (a) Temperature
- (b) Concentration
- (c) Light
- (d) All of these

**19. Higher order(>3) reactions are rare due to**

- (a) shifting of equilibrium towards reactants due to elastic collisions
- (b) loss of active species on collisions
- (c) low probability of simultaneous collision of all the reacting species
- (d) increase in entropy and activation energy as more molecules are involved.

**20. Half life period of a first order reaction is:**

- (a) directly proportional to the initial concentration of the reactant
- (b) half of the rate constant
- (c) same for all reactions
- (d) independent of initial concentration of reactants

**II FILL IN THE BLANKS**

1. Hydrolysis of ethyl acetate in an acidic solution is an example of ..... order reaction.
2. If the activation energy of the reaction is low, it proceeds at ..... rate.
3. In a multi step reaction, the ..... step determines the rate of reaction.
4. For a first order reaction, the half life period is equal to .....
5. The order and molecularity of a complex reaction.....be same.
6. The inversion of cane sugar is a ..... reaction though its molecularity is.....
7. The difference of energy between activated complex and that of the reactants is called.....
8. The unit of first order rate constant when concentration is Measured in terms of pressure and time in minutes is.....
9. A first order reaction has  $t_{1/2} = 6.93 \text{ min}$ . The rate constant is .....
10. Increase in temperature increases the number of .....

**III ASSERTION REASON TYPE QUESTIONS**

- (a) Both assertion and reason are correct statements, and reason is the correct explanation of the assertion.
  - (b) Both assertion and reason are correct statements, but reason is not the correct explanation of the assertion.
  - (c) Assertion is correct, but reason is wrong statement.
  - (d) Assertion is wrong, but reason is correct statement.
1. **Assertion:** Hydrolysis of methyl ethanoate is a pseudo first order reaction.  
**Reason:** Water is present in large excess and therefore its concentration remains constant throughout the reaction.

2. **Assertion:** The slowest elementary step in a complex reaction decides the rate of the reaction.  
**Reason:** The slowest elementary step always has the smallest molecularity.
3. **Assertion:** A catalyst increases the rate of a reaction.  
**Reason:** The catalyst increases the activation energy which in turn increases the rate of the reaction.
4. **Assertion:** Activation complex for the forward reaction will have lower energy than that for the backward reaction in an exothermic reaction.  
**Reason:** Reactants have greater energy than products for an exothermic reaction.
5. **Assertion:** Increase in temperature increases rate of reaction.  
**Reason:** More colliding molecules will have energy greater than threshold energy.
6. **Assertion:** Unit of rate constant is independent of order of reaction.  
**Reason:** The power of concentration terms in the rate equation keep changing with change in order.
7. **Assertion:** In zero order reaction, the conc. versus time graph is a straight line.  
**Reason:** The rate of change of concentration per unit time in zero order reaction remains constant.
8. **Assertion:** Half-life period is always independent of initial concentration.  
**Reason:** Half-life period is inversely proportional to rate constant.
9. **Assertion:** The rate of reaction is the rate of change of concentration of a reaction or a product.  
**Reason:** Rate of reaction remains constant during the course of reaction.
10. **Assertion:** Rate constants determined from Arrhenius equation are fairly accurate for simple as well as complex reactions.  
**Reason:** Reactant molecules undergo chemical change irrespective of their orientation during collision.

#### IV ONE WORD ANSWER TYPE QUESTIONS

1. For reactions of which order the units of rate constant and rate of reaction are same?
2. What is the difference in energy between the energy of activated complex and the average energy of reactants called?
3. A reaction is 50% complete in 2 hours and 75% complete in 4 hours. What is the order of reaction?
4. What is the effect of catalyst on activation energy of reaction?
5. For a reaction half-life is observed to be independent of the initial concentration of the reactants. What is the order of reaction?
6. What is the effect of catalyst on Gibbs energy change ( $\Delta G$ ) of a reaction?
7. For which type of reaction, order and molecularity have the same value?
8. Identify the order of reaction from the following unit of rate constant:  $\text{Lmol}^{-1}\text{s}^{-1}$
9. What is the effect of increase in surface area of reactants on rate of reaction?
10.  $E_1$  and  $E_2$  are the activation energies of the reactant and product respectively. If  $E_2 > E_1$ , predict the nature of reaction (Exothermic or Endothermic)?

11. The reaction,  $A + 2B \rightarrow C$  obeys the rate equation.  
 $\text{Rate} = K[A]^{1/2}[B]^{3/2}$   
 What is the order of a reaction?
12. Express the rate of the following reaction in terms of disappearance of hydrogen in the reaction.  
 $3H_2(g) + N_2(g) \rightarrow 2NH_3(g)$
13. For the reaction,  $A \rightarrow B$ , the rate of reaction becomes twenty seven times when the concentration of A is increased three times. What is the order of the reaction?
14. The decomposition reaction of ammonia gas on platinum surface has a rate constant  $= 2.5 \times 10^{-4} \text{ mol L}^{-1} \text{ s}^{-1}$ . What is the order of the reaction?
15. An endothermic reaction  $A \rightarrow B$  has an activation energy of 15 kcal/mole and the energy of the reaction is 5 kcal/mol. What is the activation energy for the reaction  $B \rightarrow A$ ?

#### VERY SHORT ANSWER TYPE QUESTIONS (1 MARK)

1. **The rate law for a reaction is  $\text{Rate} = K[A][B]^{3/2}$ . Can the reaction be an elementary process? Explain.**  
**Ans.** No, an elementary process would have a rate law with orders equal to its molecularities and therefore must be in integral form.
2. **For the reaction  $2H_2 + N_2 \rightarrow 2NH_3$ , how are the rate of reaction expression -  $d[H_2]/dt$  and  $d[NH_3]/dt$  inter-related?**  
**Ans.**  $-1/3 d[H_2]/dt = 1/2 d[NH_3]/dt$
3. **Identify the order of a reaction from the following rate constant :  $= 2.3 \times 10^{-5} \text{ L mol}^{-1} \text{ s}^{-1}$**   
**Ans.** Second order
4. **After five half-life periods for a first order reaction, what fraction of reactant remains?**  
**Ans.**  $1/32$
5. **What is the effect of adding catalyst on the free energy of a reaction?**  
**Ans.** No change in  $\Delta G$ .
6. **What value of k is predicted for the rate constant by Arrhenius equation is  $T \rightarrow \infty$ ? Is this value physically reasonable?**  
**Ans.** From the equation  $k = Ae^{-E_a/RT}$  if  $T \rightarrow \infty$   $k \rightarrow A$  so that  $E_a = 0$ . This is not feasible
7. **Determine the order of reaction?**  
**Step 1.**  $2NO + H_2 \rightarrow N_2 + H_2O_2$  (Slow)  
**Step 2.**  $H_2O_2 + H_2 \rightarrow 2H_2O$  (Fast)  
**Ans.**  $\text{Rate} = k[NO]^2[H_2]$   
 Order  $= 2 + 1$   
 $= 3$
8. **What is the order of reaction whose rate constant has the same units as the rate of reaction?**  
**Ans.** Zero order



**9. Why are reactions of higher order less in number ?**

**Ans.** Reaction takes place due to collide of molecules. The chances for a large number of molecules or ions to collide simultaneously are less. Hence, the reactions of higher order are less.

**10. What will be the effect of temperature on rate constant?**

**Ans.** Rate constant of a reaction is nearly doubled with rise in temperature by  $10^{\circ}\text{C}$ .

**11. State a condition under which a bimolecular reaction is kinetically first order reaction.**

**Ans.** A bimolecular reaction becomes first order reaction when one of the reactants is in excess.

**12. Why can't molecularity of any reaction be equal to zero?**

**Ans.** Molecularity of a reaction means the number of molecules of the reactants taking place in an elementary reaction. Since at least one molecule must be present, so that molecularity will be atleast one.

**13. The rate constant of a reaction is  $3 \times 10^2 \text{ min}^{-1}$ . What is its order of reaction ?(On the basis of units of rate constant)**

**Ans.** First order reaction.

**14. Three-fourth of a reaction is completed in 32 minutes. What is the half life period of this reaction?**

**Ans.** 16 minutes.

**15. What is meant by an elementary reaction?**

**Ans.** A reaction which takes place in one step is called an elementary reaction. For example:  $\text{H}_2 + \text{I}_2 \rightarrow 2\text{HI}$ .

**16. Give one example of a reaction where order and molecularity are equal?**

**Ans.**  $2\text{HI} \rightarrow \text{H}_2 + \text{I}_2$  (Order = Molecularity = 2)

**17. For a reaction  $\text{R} \rightarrow \text{P}$ , the rate becomes 2 time when the concentration of the reactant A is increased 4 times. What is the order of reaction ?**

**Ans.**  $r = k(a)^n$      $2r = k(4a)^n$      $2 = 4^n = 0.5$

**18. The rate constant of a zero order reaction in A is  $0.003 \text{ mol L}^{-1} \text{ sec}^{-1}$ . How long will it take for the initial concentration of A to fall from 0.10M to 0.075 M?**

**Ans.**  $t = [\text{R}]_0 - [\text{R}] / k = 0.10 - 0.075 / 0.003 = 8.3 \text{ second}$

**19. In a reaction  $2\text{A} \rightarrow \text{Products}$ , the concentration of A decreases from  $0.5 \text{ mol L}^{-1}$  in 10 minutes. Calculate the rate during this interval.**


**Ans.** Average rate:  $-\Delta[\text{A}] / 2\Delta t = -1/2(0.4 - 0.5/10) = 5 \times 10^{-3} \text{ M min}^{-1}$

**20. In some cases large number of colliding reactant molecules have energy more than threshold energy even then the reaction is slow . Why**

**Ans.** Because resultant molecules do not collide in proper orientation

**21. Give an example of a reaction having fractional order.**

**Ans.** Decomposition of acetaldehyde (order = 1.5)



**22. Decomposition reaction of ammonia on Pt surface has rate constant =  $2.5 \times 10^{-1} \text{ mol L}^{-1} \text{ sec}^{-1}$ . What is order of reaction ?**

**Ans.** Unit of k explain that it is zero order reaction.

**23. What is order of radioactive decay?**

**Ans.** First order

**24. For a reaction  $A+B \rightarrow \text{product}$ , the rate law is given by  $r=k[A]^{1/2}[B]^2$ . What is the order of the reaction ?**

**Ans.** Order of reaction =  $1/2 + 2 = 2.5$

**25. For a chemical reaction half life period cannot depend on concentration of solution. What is order of reaction?**

**Ans.** First order reaction

**SHORT ANSWER TYPE QUESTIONS (2 or 3 Marks)**

**Q. 1.** The rate of a particular reaction quadruples when the temperature changes from 293 K to 313 K. Calculate activation energy.

**Ans.**  $k_2/k_1 = 4$


$$T_1 = 293 \text{ K}, T_2 = 313 \text{ K}$$

$$\log \frac{k_2}{k_1} = -\frac{E_a}{2.303R} \left[ \frac{1}{T_1} - \frac{1}{T_2} \right]$$

Thus, on calculating and substituting values, we get :

$$E_a = 52.86 \text{ KJ mol}^{-1}$$

**Q. 2.** If the decomposition of nitrogen oxide as



follows a first order kinetics.

(a) Calculate the rate constant for a 0.05M solution if the instantaneous rate is  $1.5 \times 10^{-6} \text{ mol/l/s}$  ?

(b) What concentration of  $\text{N}_2\text{O}_5$  would give a rate of  $2.45 \times 10^{-5} \text{ mol L}^{-1} \text{ s}^{-1}$  ?

**Ans.** (a) Rate =  $k [\text{N}_2\text{O}_5]$

$$k = \frac{\text{Rate}}{[\text{N}_2\text{O}_5]} = \frac{1.5 \times 10^{-6}}{0.05}$$

$$k = 3.0 \times 10^{-5}$$

(b) Rate =  $2.45 \times 10^{-5} \text{ mol L}^{-1} \text{ s}^{-1}$


$$[\text{N}_2\text{O}_5] = \frac{\text{Rate}}{k} = \frac{2.45 \times 10^{-5}}{3.0 \times 10^{-5}} = 0.82 \text{ M}$$

**Q. 3. Write the difference between order and molecularity of reaction.**



**Ans.**

Order	Molecularity
1. It is the sum of the powers of concentration terms in the rate law expression.	It is the number of reacting species undergoing simultaneously collision in a reaction.
2. It is determined experimentally.	2. It is a theoretical concept.
3. Order of reaction need not to be a whole number.	3. It is whole number only.
4. Order of reaction can be zero.	4. It can't be zero or fractional.

**Q. 4. Consider the decomposition reaction :**



This reaction takes place in two steps as given below :



**(a) Determine rate law expression.**

**(b) Determine the order of reaction.**

**Ans.** (a) Rate =  $K[\text{H}_2\text{O}_2][\text{I}^-]$  because second step is rate determining step.

(b) Order = 1 + 1 = 2

**Q. 5. The decomposition of hydrocarbon follows the equation  $K = (4.5 \times 10^{11} \text{ s}^{-1}) e^{-28000\text{K}/T}$ . Calculate  $E_a$ .**

**Ans.**  $K = (4.5 \times 10^{11} \text{ s}^{-1}) e^{-28000\text{K}/T}$

Comparing the equation with Arrhenius equation,

$$K = Ae^{-E_a/RT}$$

$$-\frac{E_a}{R} = -28000 \text{ K}$$

$$E_a = 28000 \times 8.314$$

$$= 232192 \text{ J mol}^{-1}$$


**Q.6. A reaction is of second order with respect to a reactant. How is the rate of reaction affected if the conc. of the reactant is reduced to half. What is the unit or rate constant for such a reaction?**

**Ans.** Rate =  $k[\text{A}]^2$   
 Unit of  $k = \frac{\text{mol L}^{-1}}{\text{S}} = \text{k}(\text{mol}^{-1})^2$   
 $k = \text{mol}^{-1} \text{LS}^{-1}$

**Q.7.** For a first order reaction time taken for half of the reaction to complete is  $t_1$  and  $\frac{3}{4}$  of the reaction to complete is  $t_2$ . How are  $t_1$  and  $t_2$  related ?

**Ans.**  $t_2 = 2t_1$  because for  $\frac{3}{4}$ th of the reaction to complete time required is equal to two half lives.

**Q.8.**



(A) Why is the order of the reaction ?

(b) What is the slope of the curve ?

**Ans.** (a) Zero order reaction.

(b)  $[R] = [R_0] - kt$   $\therefore$  Slope =  $-k$

**Q.9.** Derive an expression to calculate time required for completion of zero order reaction.

**Ans.** For a zero order reaction,

$$R = [R]_0 - kt$$

For completion of the reaction  $[R] = 0$

$$\therefore kt = [R]_0 \quad t = \frac{[R]_0}{k}$$

$$\text{Or} \quad t = \frac{[R]_0}{k}$$

**Q.10.** The rate of a gaseous reaction becomes half when volume of the vessel is doubled. What is the order of reaction ?

**Ans.** Suppose, order of reaction is  $n$  and the reaction is  $A(g) \rightarrow \text{Products}$

$$\text{Rate} = k[A]^n \quad \dots(i)$$

When volume is doubled, molar conc. becomes half and rate of reaction gets halved.

$$\frac{\text{Rate}}{2} = k \left( \frac{A}{2} \right)^n \quad \dots(ii)$$

Dividing equation (i) by equation (ii),

$$(2)^1 = (2)^n \quad n = 1$$

**Q.11.** A reaction which is first order with respect to A has rate constant  $6 \text{ min}^{-1}$ . If we start with  $[A] = 0.5 \text{ mol L}^{-1}$ , when would  $[A]$  reach the value of  $0.05 \text{ ML}^{-1}$  ?

**Ans.** 
$$k = \frac{2.303}{t} \log \frac{[A]_0}{[A]}$$

$$k = 6 \text{ min}^{-1}, [A]_0 = 0.5, [A] = 0.05, t = ?$$

$$t = \frac{2.303}{6} \log \frac{0.5}{0.05} = \frac{2.303}{6} \log 10 = 0.3838 \text{ min}$$

**Q.12.** The conversion of the molecules X to Y follows second order kinetics. If the concentration of X is increased to three times, how will it affect the rate of formation of Y ?

**Ans.** 9 times

**Q.13.** A first order reaction has a rate constant  $1.15 \times 10^{-3} \text{ s}^{-1}$ . How long will 5 gram of this reactant take to reduce to 3 grams ?

**Ans.** 
$$t = \frac{2.303}{k} \log \frac{[R]_0}{[R]}$$

$$t = \frac{2.303}{(1.15 \times 10^{-3}) \text{ s}^{-1}} \log \left( \frac{5}{3} \right)$$

$$= \frac{2.303}{(1.15 \times 10^{-3}) \text{ s}^{-1}} [\log 5 - \log 3]$$

$$= 444 \text{ s}$$

**Q.14.** Distinguish between rate expression and rate constant of a reaction.

**Ans.** Rate expression is the way to express rate of reaction in terms of concentration of reactants. for a chemical reaction  $aA + bB \rightarrow cC + dD$  Rate  $= k[A]^x[B]^y$   
Rate constant ( $k$ ) is defined as the rate of reaction when the concentration of reactants are taken as unity.

**Q.15.** Consider a certain reaction  $A \rightarrow \text{Product}$  with  $K = 2.0 \times 10^{-2} \text{ s}^{-1}$ . Calculate the concentration of A remaining after 100 s, if the initial concentration of A is  $1.0 \text{ mol L}^{-1}$ .

**Ans.** 
$$k = \frac{2.303}{t} \log \frac{[R]_0}{[R]}$$

$$2 \times 10^{-2} = \frac{2.303}{100} \log \frac{1}{[R]}$$

$$\log \frac{1}{[R]} = \frac{2}{2.303} = 0.8684$$

$$\frac{1}{[R]} = \frac{2}{2.303} = \text{Antilog}(0.8684)$$

$$= 7.3853$$

$$[R] = \frac{1}{7.3853} = 0.135 \text{ M}$$

**Q.16.** For the reaction  $\text{NO}_2 + \text{CO} \rightarrow \text{CO}_2 + \text{NO}$ , the experimentally determined rate expression below 400K is rate:  $k[\text{NO}_2]^2$ . What mechanism can be proposed for this reaction?

**Ans.**  $\text{NO}_2 + \text{NO}_2 \xrightarrow{\text{Slow}} \text{NO} + \text{NO}_3$   
 $\text{NO}_3 + \text{CO} \xrightarrow{\text{Fast}} \text{NO}_2 + \text{CO}_2$

**Q.17.** The half life period of a first order reaction is 60 min. What % will be after 240 mins?

Hint: No. of half lives (n) =  $240/60 = 4$

$$\% \text{ of A left} = \frac{[\text{A}]_t}{2^n} = \frac{[\text{A}]_0}{2^n} = 60.25\%$$

**Q.18.** Time for half change for a first order reaction is 25 min. What % will be left after 240 mins. ?


**Ans.**

$$K = \frac{0.693}{t_{1/2}} = \frac{0.693}{25}$$

$$t_{99\%} = \frac{(2.303)}{k} \log \frac{[\text{A}]_0}{1.01[\text{A}]_t}$$

$$t_{99\%} = \frac{(2.303) \times 25 \log 100}{0.693} = 166.16 \text{ min}$$

**Q.19.** The following data were obtained during the first order thermal decomposition of  $\text{N}_2\text{O}_5$  at constant volume :



S. No.	Time per second	Total pressure (atm)
1	0	0.5
2	100	0.512

**Calculate rate constant.**

**Ans.**

	$\text{N}_2\text{O}_5$	$\text{N}_2\text{O}_4$	$+ 1/2\text{O}_2$
Initial	0.5	0	0
Final	$(0.5-x)$	x	$x/2$

$p_t = 0.5 - x + x + x/2 = 0.5 + x/2$   
 $0.5 + x/2 = 0.512$   
 $x = 0.024$

$$k = \frac{2.303}{t} \log \frac{P_i}{P_t}$$

log  $k = \frac{2.303}{100} \log \frac{0.5}{0.5 - 0.024}$

$$k = \frac{2.303 \times 0.0214}{100} = 4.92 \times 10^{-4} \text{ s}^{-1}$$

**Q.20.** The decomposition of A into product has value of  $k$  as  $4.5 \times 10^3 \text{ s}^{-1}$  at  $10^\circ\text{C}$  and energy of activation  $60 \text{ KJ}$ ?


**Ans.**

$$\log \frac{k_2}{k_1} = \frac{E_a}{2.303R} \left( \frac{T_2 - T_1}{T_1 T_2} \right)$$

$$\log \left( \frac{1.5 \cdot 10^4}{4.5 \times 10^3} \right) = \frac{60000}{2.303 \times 8.314} \left[ \frac{T_2 - 283}{283 T_2} \right]$$

$$T_2 = 297 \text{ K}$$

**Q.21.** For a chemical reaction  $R \rightarrow P$ , the variation in the concentration (R) vs time ( $t$ ) plot is given :




- Predict the order of reaction.
- Write down its rate law.
- What is the slope of the curve?

**Ans.** (a) Zero

(b)  $\text{Rate} = k[R]^0$

(c)  $\text{Slope} = -k$

**Q.22.** (a) Write rate law and order of the following reaction:



(b) define energy of activation of a reaction

(c) What is the relationship between rate constant and activation energy of a reaction?

**Ans.** (a)  $\text{Rate} = k [\text{AB}][\text{C}_2]$ , Order =  $1+1=2$

(b) Refer 'points to remember'

(c)  $k = A e^{-E_a/RT}$



**Q.23.** The rate constant for first order reaction is 60/s , How much time will it take to reduce the concentration of the reaction to 1/10 of its initial value ?

**Ans.**

$$t = \frac{2.303}{K} \log \frac{[R]_0}{[R]}$$

$$t = \frac{2.303}{\frac{1}{10}} \log \frac{[R]_0}{[R]}$$

$$t = \frac{2.303}{60} \log 10$$

$$t = \frac{2.303}{60} \\ = 3.38 \times 10^{-2} \text{ s}^{-1}$$

**Q.24.** The rate of most of reaction double when their temperature is raised from 298K to 308K. Calculate the activation energy of such a reaction .

$$\log \frac{K_2}{K_1} = \frac{E_a}{2.303R} \left[ \frac{1}{T_1} - \frac{1}{T_2} \right]$$

$$E_a = \frac{2.303 \times 8.314 \times 298 \times 308 \times 0.3010}{1000}$$

$$E_a = 52.89 \text{ KJ/mol}$$

**Q.25.** A first order reaction takes 69.3 min for 50% completion. Set up an equation for determining the time needed for 80% completion

$$K = \frac{0.693}{t_{1/2}} = \frac{0.693}{69.3} \text{ min} \\ = 10^{-2} \text{ min}^{-1}$$

$$T = \frac{2.303}{K} \log \frac{[R]_0}{[R]}$$

$$= \frac{2.303}{10^{-2}} \log 5$$

$$= 160.9 \text{ min}$$

**Q.26.** The activation energy of a reaction is 94.14 KJ/mol and the value of rate constant at 40° C is  $1.8 \times 10^{-1} \text{ sec}^{-1}$ . Calculate the frequency factor A.

**Ans.** Given,  $E_a = 94.14 \times 10^{-3} \text{ Jmol}^{-1}$ ,  $T = 40 + 273 = 313 \text{ K}$ ,  $K = 1.8 \times 10^{-1} \text{ Sec}^{-1}$

By using,  $K = Ae^{-E_a/RT}$        $\ln K = \ln A - \frac{E_a}{RT}$

Or  $\log K = \log A - \frac{E_a}{2.303RT}$

$$\text{Or } \log(1.8 \times 10^{-1}) + \frac{94.19 \times 10^3}{2.303 \times 8.314 \times 313} = \log A$$

$$\text{Or } A = \text{antilog}(10.9635) = 9.194 \times 10^{10} \text{ Sec}^{-1}$$

**Q.27.** The rate constant of a reaction at 700 K and 760 K are  $0.011 \text{ M}^{-1} \text{ S}^{-1}$  and  $0.105 \text{ M}^{-1} \text{ S}^{-1}$  respectively. Calculate the value of arrhenius parameters

Ans.  $2.824 \times 10^{10}$

**Q.28.** The initial concentration of  $\text{N}_2\text{O}_5$  in the first order reaction  $\text{N}_2\text{O}_5 \rightarrow 2\text{NO}_2 + \frac{1}{2}\text{O}_2$  was  $1.24 \times 10^{-2} \text{ mol L}^{-1}$  at 318 K. The concentration of  $\text{N}_2\text{O}_5$  after 60 minutes was  $0.20 \times 10^{-2} \text{ mol L}^{-1}$ . Calculate the rate constant of the reaction at 318 K.

$$\begin{aligned} \text{Ans. } K &= \frac{2.303}{t} \log \frac{[A]_0}{[A]} = \frac{2.303}{t} \log \frac{[\text{N}_2\text{O}_5]_0}{[\text{N}_2\text{O}_5]_t} = \frac{2.303}{60} \log \frac{1.24 \times 10^{-2}}{0.2 \times 10^{-2}} \\ &= \frac{2.303}{60} \log 6.2 = \frac{2.303}{60} \times 0.7924 \text{ min}^{-1} \\ &= 0.0304 \text{ min}^{-1} \end{aligned}$$

### LONG ANSWER TYPE QUESTIONS (5 Marks)

**Q.1.** (a) Define order of reaction  
(b) Rates of reaction double with every 100 rise in temperature. If this generalization holds for a reaction in the temperature ranges 298 K to 308 K, What would be the value of activation energy for their reaction?  
**R =  $8.314 \text{ J K}^{-1} \text{ mol}^{-1}$**

Ans. (a) Order of Reaction: It is the sum of powers to which to conc. terms are raised in rate law expression.

$$\log \frac{K_2}{K_1} = \frac{E_a}{2.303R} \left[ \frac{1}{T_1} - \frac{1}{T_2} \right]$$

Here,  $T_1 = 298 \text{ K}$ ,  $T_2 = 308 \text{ K}$ ,  $R = 8.314 \text{ J K}^{-1} \text{ Mol}^{-1}$

$$\frac{K_2}{K_1} = 2$$

$$\log 2 = \frac{E_a}{2.303 \times 8.314} \left[ \frac{1}{298} - \frac{1}{308} \right]$$

$$0.3010 = \frac{E_a}{2.303 \times 8.314} \left[ \frac{10}{298 \times 308} \right]$$

$$\begin{aligned}
 E_a &= \frac{0.3010 \times 2.303 \times 8.314 \times 298 \times 308}{10} \\
 &= 52898 \text{ J mol}^{-1} \\
 &= 52.898 \text{ KJ mol}^{-1} \text{ ----}
 \end{aligned}$$

**Q.2. (a) What are pseudo order reaction ? Give example.**


**(b) Rate constant K of a reaction varies with temperature 'T' according to the equation :**

$$\log k = \log A - \frac{E_a}{2.303R} \left( \frac{1}{T} \right)$$

where  $E_a$  is the activation energy. When a graph is plotted for  $\log k$  vs  $1/T$ , a straight line with a slope of  $-4250 \text{ K}$  is obtained. Calculate  $E_a$  for the reaction.

**Ans. (a)** The chemical reaction which look like higher order reaction but in real they follow lower order kinetics.

For example,



$$\text{Rate} = k [\text{CH}_3\text{COOC}_2\text{H}_5]$$

$$\text{Order} = 1$$

$$(b) \quad \text{Slope} = \frac{E_a}{2.303R} = -4250 \text{ K}$$

$$\text{So,} \quad E_a = -2.303 \times R \times \text{Slope} = -2.303 \times 8.314 \text{ J K}^{-1} \text{ mol}^{-1} \times 4250$$

$$= 81375.3 \text{ J mol}^{-1} = 81.375 \text{ kJ mol}^{-1}$$

- Q.3. (a) Determine the units of rate constant for first and zero order reaction.**  
**(b) Show that time required for the completion of 99% of the first order reaction is twice the 90% of completion of the reaction.**

**Ans. (a)**  $k = (\text{mol})^{1-n} \text{L}^{n-1} \text{s}^{-1}$

For zero order,  $n = 0$

So,  $k = (\text{mol})^{1-0} \text{L}^{0-1} \text{s}^{-1} = \text{s}^{-1} \text{mol L}^{-1}$

For first order,  $n = 1$

$$k = (\text{mol})^{1-n} \text{L}^{n-1} \text{s}^{-1}$$

So,  $k = (\text{mol})^{1-1} \text{L}^{1-1} \text{s}^{-1}$   
 $= \text{s}^{-1}$

- (b)** For a first order reaction,

$$t = \frac{2.303}{k} \log \frac{[A]_0}{[A]}$$

$$[A]_0 = a, [A] = a - \frac{a \times 99}{100} = 0.01 a$$

$$t(99\%) = \frac{2.303}{k} \log \frac{a}{0.01a} = \frac{2.303}{k} \log 100 = \frac{2.303}{k} \times 2$$

...(i)

For 90% completion of reaction,

$$[A] = a - \frac{a \times 99}{100} = 0.1a$$

$$t(90\%) = \frac{2.303}{k} \log \frac{a}{0.1a} = \frac{2.303}{k} \times 1$$

...(ii)

Dividing equation (i) by equation (ii), we get

$$t(99\%) = 2 \times t(90\%)$$

Q.4. (a) Define rate constant of reaction.

(b) A first order reaction takes 40 mins for 30% decomposition. Calculate  $t_{1/2}$ .

Ans. (a) **Rate constant** : It is the rate of chemical reaction when the concentration of reactant taken as unity at a given temperature.

(b) Let initial conc. =  $a$

$$\begin{aligned}\text{Conc. after 40 mins.} &= a - \frac{a \times 30}{100} \\ &= 0.70 a\end{aligned}$$


$$k = \frac{2.303}{t} \log \frac{[A]_0}{[A]}$$

$$= \frac{2.303}{40} \log \frac{a}{0.70a} = \frac{2.303}{40} \log \frac{1}{0.70}$$

$$= \frac{2.303}{40} \times 0.1549 = 8.92 \times 10^{-3} \text{ min}^{-1}$$

$$t_{1/2} = \frac{0.693}{k} = \frac{0.693}{8.92 \times 10^{-3}} = 77.7 \text{ min}$$

Q. 5. (a) Determine the order of reaction and also determine the units of rate constant.



(b) The following data were given for thermal decomposition of  $\text{SO}_2\text{Cl}_2$  at a constant volume :



Exp.	Time/s	Total p/atm
1	0	0.5
2	100	0.6

Calculate the rate of the reaction when total pressure is 0.65 atm.

Ans. (a) First order reaction

$$k = \text{mol}^{1-n} \text{L}^{n-1} \text{s}^{-1}$$

$$n = 1$$

$$k = (\text{mol})^{1-1} \text{L}^{1-1} \text{s}^{-1}$$

$$= \text{s}^{-1}$$

$$(b) \quad k = \frac{2.303}{t} \log \frac{P_i}{(2P_i - P_t)}$$

$$= \frac{2.303}{100} \log \frac{0.5}{(2 \times 0.5 - 0.6)}$$

$$= \frac{2.303}{100} \log \frac{0.5}{0.4}$$

$$= \frac{2.303}{100} \times 0.969 = 2.23 \times 10^{-3} \text{s}^{-1}$$

Now, Rate =  $P_{\text{SO}_2\text{Cl}_2}$

Pressure of  $\text{SO}_2\text{Cl}_2$  when total pressure = 0.65 atm


$$P_{\text{SO}_2\text{Cl}_2} = 2P_i - P_t$$

$$= 2 \times 0.5 - 0.65 = 0.35 \text{ atm}$$

$$\text{Rate} = 2.23 \times 10^{-3} \times 0.35 = 7.8 \times 10^{-4} \text{ atm s}^{-1}$$



**Q.6. (a) The activation energy of a reaction is 100 kJ/mol. In the presence of catalyst the activation energy is decreased by 75%. What is the effect on rate constant of the reaction at 20°C ?**



**The rate of disappearance of B is  $1 \times 10^{-2} \text{ mol L}^{-1} \text{ sec}^{-1}$ . What will be**

**(i) rate of reaction (ii) rate of change in the concentration of A and C ?**

Ans. (a)  $2.35 \times 10^{10}$  times

$$[\text{Hint: } \log \frac{K_2}{K_1} = \frac{E_a - E_p}{2.303 RT} \Rightarrow \log \frac{K_2}{K_1} = \frac{75 \times 10^3}{2.303 \times 8.314 \times 293}]$$

(b)  $(5 \times 10^{-3}, 5 \times 10^{-3}, 15 \times 10^{-3}) \text{ ML}^{-1} \text{ sec}^{-1}$

**Q.7. (a) A reaction is of first order in A and of second order in B. Write the differential rate equation for this reaction. How will its initial rate be affected if the concentration of both A and B are together doubled ?**

**(b) The rate constant k of a reaction increases four fold when the temperature changes from 300 K to 320 K. Calculate the activation energy for the reaction. ( $R = 8.314 \text{ J K}^{-1} \text{ mol}^{-1}$ )**

Ans. (a)  $r = k[A]^1 \dots\dots (1)$

$$r = k[B]^2 \dots\dots (2)$$

Differential rate equation is

$$r = k[A]^1[B]^2 \dots\dots (3)$$

When conc. of both A and B is doubled

$$r' = k[2A]^1[2B]^2 \dots\dots (4)$$

Divide (4) by (3),

$$r'/r = 8$$

Thus, rate becomes 8 times.

$$(b) \frac{K_2}{K_1} = 4, T_1 = 300\text{K}, T_2 = 320\text{ K}$$

$$R = 8.314 \text{ J K}^{-1} \text{ mol}^{-1}$$

$$\log \frac{K_2}{K_1} = \frac{E_a}{2.303 R} \left[ \frac{T_2 - T_1}{T_1 T_2} \right]$$

$$\Rightarrow \log 4 = \frac{E_a}{2.303 \times 8.314 \text{ JK}^{-1} \text{ mol}^{-1}} \times \left[ \frac{320 - 300}{300 \times 320} \right] \text{ K}^{-1}$$

$$\Rightarrow 0.6020 = \frac{E_a}{2.303 \times 8.314 \text{ JK}^{-1} \text{ mol}^{-1}} \left[ \frac{20 \text{ K}^{-1}}{96 \times 10^3} \right]$$

$$E_a = 55327.6 \text{ J mol}^{-1} = 55.33 \text{ kJ mol}^{-1}$$

- Q.8.** (a) What are the factors on which rate of the reaction depends ?  
Discuss each factor in brief.  
(b) The following results have been obtained during the kinetics studies of the reaction:  
 $2A + B \rightarrow C + D$

Experiment	[A] mol L <sup>-1</sup>	[B] mol L <sup>-1</sup>	Initial rate of formation of D mol L <sup>-1</sup> min <sup>-1</sup>
I	0.1	0.1	$6.0 \times 10^{-3}$
II	0.3	0.2	$7.2 \times 10^{-2}$
III	0.3	0.4	$2.88 \times 10^{-1}$
IV	0.4	0.1	$2.40 \times 10^{-1}$

**Determine the rate law and the rate constant for the reaction.**

- Ans.** (a) Refer "Points to Remember"  
(b) For the reaction  $2A + B \rightarrow C + D$   
Assume rate law expression as  $\text{Rate} = k[A]^a[B]^b$   
According to question,

$$6.0 \times 10^{-3} = k(0.1)^a(0.1)^b \quad \dots(i)$$

$$7.2 \times 10^{-2} = k(0.3)^a(0.2)^b \quad \dots(ii)$$

$$2.88 \times 10^{-1} = k(0.3)^a(0.4)^b \quad \dots(iii)$$

$$2.40 \times 10^{-2} = k(0.4)^a(0.1)^b \quad \dots(iv)$$

Divide eqn. (iv) by (i), we get

$$4 = (4)^a \therefore a = 1$$

Divide eqn. (iii) by (ii), we get

$$4 = (2)^b$$

$$22 = (2)^b \therefore b = 2$$

Order with respect to A = 1

Order with respect to B = 2

$$\therefore \text{Rate Law} = \frac{d[D]}{dt} = k[A][B]^2$$

On putting the value of 'a' and 'b' into any equation say (i)

$$6.0 \times 10^{-3} \text{ M min}^{-1} = k(0.1 \text{ M})(0.1 \text{ M})^2$$

$$\therefore k = 6 \text{ M}^{-2} \text{ min}^{-1}$$



- Q.9.** (a) Derive the general form of the expression for the half life of a first order reaction.  
 (b) The decomposition of  $\text{NH}_3$  on platinum surface is a zero order reaction. What are the rates of production of  $\text{N}_2$  and  $\text{H}_2$  if  $k = 2.5 \times 10^{-4} \text{ mol L}^{-1} \text{ s}^{-1}$ ?

**Ans.** (a)  $t = \frac{2.303}{k} \log \frac{[\text{R}]_0}{[\text{R}]}$

When  $[\text{R}] = \frac{[\text{R}]_0}{2}$ , then  $t = t_{1/2}$

$\therefore t_{1/2} = \frac{2.303}{k} \log \frac{[\text{R}]_0}{[\text{R}]_{0/2}} = \frac{2.303}{k} \log 2$

or  $t_{1/2} = \frac{2.303}{k} \times 0.3010$

or  $t_{1/2} = \frac{0.693}{k}$

**(b)**



$$-\frac{1}{2} \frac{d[\text{NH}_3]}{dt} = \frac{d[\text{N}_2]}{dt} + \frac{1}{3} \frac{d[\text{H}_2]}{dt}$$

$$\frac{d[\text{NH}_3]}{dt} = \text{rate} = k [\text{NH}_3]^0 = 2.5 \times 10^{-4} \text{ mol L}^{-1} \text{ sec}^{-1}$$

$$\frac{d[\text{N}_2]}{dt} = -\frac{1}{2} \frac{d[\text{NH}_3]}{dt} = \frac{1}{2} \times 2.5 \times 10^{-4} \text{ mol L}^{-1} \text{ sec}^{-1}$$

$$d[\text{H}_2] = -\frac{3}{2} \frac{d[\text{NH}_3]}{dt} = \frac{3}{2} \times 2.5 \times 10^{-4} = 3.75 \times 10^{-4} \text{ mol L}^{-1} \text{ sec}^{-1}$$

$$= -\frac{d[\text{NH}_3]}{dt} = k [\text{NH}_3]^0 = 2.5 \times 10^{-4} \text{ mol L}^{-1} \text{ sec}^{-1}$$

Rate of production of  $\text{N}_2 = 2.5 \times 10^{-4} \text{ mol L}^{-1} \text{ sec}^{-1}$

- Q.10.** (i) What is rate constant ?  
 (ii) On what factors it depends ?  
 (iii) In a pseudo first order reaction of hydrolysis of an ester in  $H_2O$ , the following results were obtained:

t/s	0	30	60	90
Ester (M/L)	0.55	0.31	0.17	0.085

- (a) Calculate the average rate of reaction between the time interval 30 to 60 sec.  
 (b) Calculate the pseudo first order rate constant for the hydrolysis of ester.

**Ans.** (i) rate constant(k) of a chemical reaction is rate of reaction when the concentration of the reactants is unity .  
 (ii) Rate constant (k) depends upon (i) temperature (ii) order of reaction  
 (iii)

(a) Average rate during 30-60 sec.  $= \frac{0.17 - 0.31}{60 - 30} = 4.67 \times 10^{-3} \text{ mol L}^{-1} \text{ sec}^{-1}$

(b)  $k_{30} = \frac{2.303}{t} \log \frac{[A]_0}{[A]} = \frac{2.303}{30} \log \frac{0.55}{0.31}$

$$k_{60} = \frac{2.303}{60} \log \frac{0.55}{0.17}$$

$$k_{90} = \frac{2.303}{90} \log \frac{0.55}{0.085}$$

$$\text{Average } K = 1.98 \times 10^{-2} \text{ sec}^{-1}$$

### CASE STUDY BASED QUESTIONS

**1. Read the passage given below and answer the questions that follow:**

The rate of reaction is the change of concentration of reactant or product with time. The rate law for the reaction  $aA + bB \rightarrow cC + dD$  the rate law is  $\text{rate} = k[A]^a[B]^b$ . The rate of reaction is calculated by knowing  $k$ ,  $a$  and  $b$ . The rate laws are determined experimentally. During the collisions among two A and two B molecules, doubling the number of either type of molecule increases the number of collisions to eight. The species temporarily formed by the reactant molecules as a result of the collision before they form the product is called the *activated complex*. The temperature-dependent rate constant is given by the Arrhenius equation. In many cases, the sum of a series of simple reactions are called elementary steps or elementary reactions because they represent the progress of the overall reaction at the molecular level. The sequence of elementary steps that leads to product formation is called the reaction mechanism. The number of molecules reacting in an elementary step determines the molecularity of a reaction.

- (A) **On which of the following quantities does the rate constant of a reaction depend?**
- |                                 |                        |
|---------------------------------|------------------------|
| (a) Concentrations of reactants | (b) Nature of reactant |
| (c) Temperature                 | (d) All the above      |
- (B) **Instantaneous rate of reaction is measured from the:**
- |  |                                      |
|--|--------------------------------------|
| (a) graph of time versus concentration | (b) molecularity of reaction         |
| (c) integration method                 | (d) reaction mechanism of a reaction |
- (C) **The order of reaction is always defined in terms of:**
- |                            |  |
|----------------------------|--|
| (a) product concentration  | (b) rate constant of a reaction                                      |
| (c) reactant concentration | (d) ratio of the product concentration to the reactant concentration |
- (D) **The number of molecules reacting in an elementary step of a reaction may be:**
- |       |         |        |       |
|-------|---------|--------|-------|
| (a) 1 | (b) 1/2 | (c) 12 | (d) 5 |
|-------|---------|--------|-------|

**2. Read the passage given below and answer the questions that follow:**

The integrated rate equations can be fitted with kinetic data to determine the order of a reaction. The integrated rate equations for zero and first order reaction are:

Zero order :  $[A] = -kt + [A]_0$

First order :  $\log [A] = \frac{-kt}{2.303} + \log [A]_0$

These equations can also be used to calculate the half life periods of different reactions, which give the time during which the concentration of a reactant is reduced to half of its initial concentration i.e. at time  $t_{1/2}$ ;  $[A] = [A]_0/2$

**In the following questions a statement of assertion followed by a statement of reason is given. Choose the correct answer out of the following choices.**

- a) Assertion and reason both are correct statements and reason is correct explanation for assertion.
  - b) Assertion and reason both are correct statements but reason is not correct explanation for assertion.
  - c) Assertion is correct statement but reason is wrong statement.
  - d) Assertion is wrong statement but reason is correct statement.
- (A) **ASSERTION:** For the first order reaction the units of rate constant are  $\text{time}^{-1}$ .  
**REASON:** The rate of first order reaction remains constant throughout.
- (B) **ASSERTION:** In zero order reaction, the conc. versus time graph is a straight line.  
**REASON:** The rate of change of concentration per unit time in zero order reaction remains constant.
- (C) **ASSERTION:** Half-life period is always independent of initial concentration.  
**REASON:** Half-life period is inversely proportional to rate constant.
- (D) **ASSERTION:** The slowest step in the reaction is rate determining step.  
**REASON:** Order of a reaction is given by rate law expression.

## ANSWERS

### I MULTIPLE CHOICE QUESTION:

1. (a) 2. (a) 3. (a) 4. (a) 5. (d) 6. (b) 7. (a) 8. (c) 9. (d) 10. (c) 11. (c)  
12. (a,d) 13. (b) 14. (c) 15. (a) 16. (b) 17. (b) 18. (d) 19. (c) 20. (d)

### II FILL IN THE BLANKS:

1. Pseudo first                      2. Fast                      3. Slowest  
4. 0.693/k                      5. Cannot                      6. First, two  
7. Activation energy                      8.  $\text{min}^{-1}$                       9.  $0.1 \text{ min}^{-1}$   
10. Effective collisions

### III ASSERTION REASON TYPE QUESTIONS:

1. (a) 2. (c) 3. (c) 4. (d) 5. (a) 6. (d) 7. (a) 8. (d) 9. (c) 10. (c)

### IV ONE WORD ANSWER TYPE QUESTIONS:

1. Zero order                      2. Activation energy                      3. First order reaction  
4. Decrease                      5. First order                      6. No effect  
7. Elementary reaction                      8. Second order                      9. Increase  
10. Endothermic                      11. Two  
12. Rate of Disappearance of  $\text{H}_2 = - \frac{d[\text{H}_2]}{dt}$                       13. 3  
14. Zero                      15. 10 Kcal/mol

### CASE STUDY BASED QUESTIONS:

PASSAGE: 1: (A) c (B) a (C) c (D) a

PASSAGE: 2: (A) c (B) a (C) d (D) b

**UNIT TEST-1**  
**CHAPTER - 4**  
**CHEMICAL KINETICS**

**TIME ALLOWED : 1 HR****M.M. : 20**

1. Which of the following statements is not correct for the catalyst? 1
  - (a) It catalyses the forward and backward reaction to the same extent.
  - (b) It alters  $\Delta G$  of the reaction.
  - (c) It is a substance that does not change the equilibrium constant of a reaction.
  - (d) It provides an alternate mechanism by reducing activation energy between reactants and products.
2. For the reaction:  $5\text{Br}^-(\text{aq}) + \text{BrO}_3^-(\text{aq}) + 6\text{H}^+(\text{aq}) \rightarrow 3\text{Br}_2(\text{aq}) + 3\text{H}_2\text{O}(\text{l})$  1  
 Which of the following expression is correct for the rate of the reaction?
  - (a)  $\Delta[\text{Br}^-]/\Delta t = 5\Delta[\text{H}^+]/\Delta t$
  - (b)  $\Delta[\text{Br}^-]/\Delta t = 6\Delta[\text{H}^+]/5\Delta t$
  - (c)  $\Delta[\text{Br}^-]/\Delta t = 5\Delta[\text{H}^+]/6\Delta t$
  - (d)  $\Delta[\text{Br}^-]/\Delta t = 6\Delta[\text{H}^+]/\Delta t$
3. For a zero order reaction will the molecularity be equal to zero? Explain. 1

**ASSERTION REASON TYPE QUESTIONS**

- (a) Both assertion and reason are correct statements, and reason is the correct explanation of the assertion.
  - (b) Both assertion and reason are correct statements, but reason is not the correct explanation of the assertion.
  - (c) Assertion is correct, but reason is wrong statement.
  - (d) Assertion is wrong, but reason is correct statement.
4. **ASSERTION:** The rate of reaction is the rate of change of concentration of a reaction or a product. 1  
**REASON:** Rate of reaction remains constant during the course of reaction.
5. **ASSERTION:** Rate constants determined from Arrhenius equation are fairly accurate for simple as well as complex reactions. 1  
**REASON:** Reactant molecules undergo chemical change irrespective of their orientation during collision.
6.
  - (i) Why does the rate of a reaction increase with increase in temperature? 2
  - (ii) Why is the probability of reaction with molecularity higher than three very rare?


8. After 24 hours, only 0.125 g out of the initial quantity of 1g of a radioactive isotope remains behind. What is its half-life period? 3
9. A first order reaction is 50% completed in 40 minutes at 300K and in 20 minutes at 320K. Calculate the activation energy of the reaction. 3  
(Given:  $\log 2 = 0.3010$ ,  $\log 4 = 0.6021$ ,  $R = 8.314 \text{ JK}^{-1} \text{ mol}^{-1}$ )
10. (i) Define order of reaction. How does order of a reaction differ from molecularity for a complex reaction? 5
- (ii) Define instantaneous rate of reaction
- (iii) Why  $\text{H}_2$  and  $\text{O}_2$  do not react at room temperature?
- (iv) For which type of reactions, order and molecularity have the same value?

**UNIT TEST-2**  
**CHAPTER - 4**  
**CHEMICAL KINETICS**

**TIME ALLOWED : 1 HR****M.M. : 20**

1. Identify the order of reaction from the following unit for its rate constant:  $\text{Lmol}^{-1}\text{s}^{-1}$  (1)
2. for the reaction,  $4\text{NH}_3 + 5\text{O}_2 \rightarrow 4\text{NO} + 6\text{H}_2\text{O}$  If rate of formation of NO is  $6 \times 10^{-4} \text{ atm min}^{-1}$ , calculate the rate of formation of  $\text{H}_2\text{O}$  (1)
3. Write the relationship between  $k$  and  $t_{1/2}$  (half life period ) for first order reaction. (1)
4. Explain with an example, what is a pseudo first order reaction? (1)
5. For the chemical reaction, what is the effect of a catalyst on the rate constant of the reaction? (1)
6. Differentiate between: Average rate and instantaneous rate of a chemical reaction. (2)
7. Observe the graph shown in figure and answer the following question. (2)

- (a) What is the order of the reaction ?
- (b) What is the slope of the curve ?



8. Show that in case of first order reaction, the time required for 99.9% of the reaction to take place is about ten times than that required for half the reaction. (3)
9. The rate constant of a reaction at 500K and 700 K are  $0.02 \text{ s}^{-1}$  and  $0.07 \text{ s}^{-1}$  respectively. Calculate the value of  $E_a$  and  $A$ . (3)
10. (i) Answer the following questions (3)
  - (a) The conversion of molecules A to B follow second order kinetics. If concentration of A is increased to three times, how will it affect the rate of formation of B ?
  - (b) Write the expression of integrated rate equation for zero order reaction.
  - (ii) A first order reaction is 20% complete in 20 minutes. Calculate the time taken for the reaction to go to 80% completion (5)



## UNIT 8

### d- and f- Block Elements

#### Points to Remember

Element having partially filled d-subshell in their elemental or common oxidation state.

Zn, Cd and Hg not considered as transition elements as not having partially filled d - subshell in their elemental or common oxidation state.

General Electronic configuration of the d-block elements is  $(n-1)d^{1-10}ns^{1-2}$ .

**Outer Electronic Configurations of four d-block series elements.**

1st Series										
	Sc	Ti	V	Cr	Mn	Fe	Co	Ni	Cu	Zn
Z	21	22	23	24	25	26	27	28	29	30
4s	2	2	2	1	2	2	2	2	1	2
3d	1	2	3	5	5	6	7	8	10	10

2nd Series										
	Y	Zr	Nb	Mo	Te	Ru	Rh	Pd	Ag	Cd
Z	39	40	41	42	43	44	45	46	47	48
5s	2	2	1	1	1	1	1	0	1	2
4d	1	2	4	5	6	7	8	10	10	10

3rd Series										
	La	Hf	Ta	W	Re	Os	Ir	Pt	Au	Hg
Z	57	72	73	74	75	76	77	78	79	80
6s	2	2	2	2	2	2	2	1	1	2
5d	1	2	3	4	5	6	7	9	10	10

4th Series										
	Ac	Rf	Db	Sg	Bh	Hs	Mt	Ds	Rg	Uub
Z	89	104	105	106	107	108	109	110	111	112
7s	2	2	2	2	2	2	2	2	1	2
6d	1	2	3	4	5	6	7	8	10	10

**Atomic and Ionic Radii :** In a given transition series, from left to right net nuclear charge as well as screening effect increase. Due to this, the atomic and ionic radii for transition elements for a given series show a decreasing trend for first five elements and then screening effect counter balance nuclear charge due to which atomic size becomes almost constant for next five elements of the series. Size of 4d and 5d series elements is almost same due to lanthanoid contraction.


### General Properties of the Transition Elements

**Enthalpies of Atomisation:** Transition elements exhibit higher enthalpies of atomization because of large number of unpaired electrons in their atoms they have stronger interatomic interaction and hence stronger bonding between atoms.

**Ionisation Enthalpies :** • In a series from left to right, ionization enthalpy increases due to increase in nuclear charge.

- The irregular trend in the first ionization enthalpy of the 3d metals, though of little chemical significance, considering that the removal of one electron alters the relative energies of 4s and 3d orbitals.

#### Standard Electrode Potentials:



**Oxidation States :** Transition metals shows variable oxidation state due to two incomplete outermost shells. Only stable oxidation states of the first row transition metals are Sc(+3), Ti(+4), V(+5), Cr(+3,+6), Mn(+2,+7), Fe(+2,+3), Co(+2,+3), Ni(+2), Cu(+2), Zn(+2)

- The transition elements in their lower oxidation states (+2 and +3) usually forms ionic compounds. In higher oxidation state compounds are normally covalent.
- Only Os and Ru show +8 oxidation states in their compounds.
- Ni and Fe in  $\text{Ni(CO)}_4$  and  $\text{Fe(CO)}_5$  show zero oxidation state.

### General Properties of the Transition Elements

#### Magnetic Properties :

- When a magnetic field is applied to substances, mainly two types of magnetic behavior are observed : diamagnetism and paramagnetism. Paramagnetism due to presence of unpaired electrons, each such electron having a magnetic moments associated with its spin angular momentum.
- The magnetic moment is determined by the number of unpaired electrons.

$$\text{Magnetic moment} = \sqrt{n(n+2)}$$

BM


where, n=number of unpaired electron. If all electrons are paired, substance will be diamagnetic and magnetic moment will be zero.

#### Formation of Complex Compounds


- Transition metals have small size high nuclear charge which facilitates the acceptance of lone pair of elements from ligands.
- They have vacant d-orbitals of appropriate energy in order to accommodate the lone pair of electrons.


#### Formation of Coloured

- The d-orbitals are non-degenerated in presence of ligands. When an electron from a lower energy d-orbitals is excited to a higher energy d-orbital, the energy of required wavelength is absorbed and rest light is transmitted out. Therefore, the colour observed corresponds to the complementary colour of the light absorbed.
- In  $\text{V}_2\text{O}_5$ , V is in +5 oxidation state. It is coloured due to defects in crystal lattice.



### Some important Compounds of Transition Elements





### The Inner Transition Elements (f-Block)

The f-block consists of the two series, lanthanoids and actinoids. Lanthanoids are known as rare earth metals and actinoids are known as radioactive elements (Th to Lr).

#### LANTHANOIDS:

General electronic configuration  $[\text{Xe}] 4f^{1-14}, 5d^{0-2}, 6s^2$ .

- Atomic and ionic size from left to right, decreases due to increase in nuclear charge. This is known as lanthanoid contraction.
- All the lanthanoids are silvery white soft metals and tarnish rapidly in air.
  - Many trivalent lanthanoid ions are coloured both in the solid state and in aqueous solutions. Neither  $\text{La}^{3+}$  nor  $\text{Lu}^{3+}$  ion shows any colour but the rest do so.
  - The lanthanoid ions other than the  $f^0$  type ( $\text{La}^{3+}$  and  $\text{Ce}^{4+}$ ) and the  $f^{14}$  type ( $\text{Yb}^{2+}$  and  $\text{Lu}^{3+}$ ) are all paramagnetic. The paramagnetism arises to maximum in neodymium.
  - Oxidation states  $\rightarrow \text{Ce}^{4+}$ ; (Some elements) is favoured by its noble gas configuration, but it is a strong oxidant reverting to the common +3 state. The Eo value for  $\text{Ce}^{4+}/\text{Ce}^{3+}$  is +1.47 V, the reaction rate is very slow and hence,  $\text{Ce(IV)}$  is a good analytical reagent. Pr, Nd, Tb and Dy also exhibit +4 state but only in oxides.  $\text{Eu}^{2+}$  is formed by losing the two s-electrons and its  $f^7$  configuration accounts for the formation of this ion. However,  $\text{Eu}^{2+}$  is a strong reducing agent changing to the common +3 state. Similarly,  $\text{Yb}^{2+}$  which has  $f^{14}$  configuration is a reductant,  $\text{Tb}^{4+}$  has half-filled f-orbitals and is an oxidant.
- Misch metals, contain lanthanoids about 90-95% (Ce 40-5%, Lanthanum and neodymium 44%) iron 4.5%, calcium, carbon and silicon, used in cigarette and gas lighters, toys, tank and tracer bullets.

#### ACTINOIDS:

- General electronic configuration  $[\text{Rn}] 5f^{1-14}, 6d^{0-2}, 7s^2$ .
  - Actinoids exhibit a range of oxidation states due to comparable energies of 5f, 6d and 7s orbitals. The general oxidation state of actinoids is +3.
  - All the actinoids are strong reducing agents and very reactive.
  - Actinoids also react with oxygen, halogen, hydrogen and sulphur, etc. like lanthanoids.
  - Actinoids are radioactive in nature and therefore, it is difficult to study their chemical nature.

## OBJECTIVE TYPE QUESTIONS

### I. MULTIPLE CHOICE QUESTIONS

1. One of the characteristics of transition metals to form the complex ion is:
  - (a) Having unpaired electron in d-subshell
  - (b) Having paired electrons in d-subshells
  - (c) Providing empty d-orbitals
  - (d) Having small charge/size ratio
2. The correct electronic configuration of copper atom is:
  - (a)  $3d^{10}4s^1$
  - (b)  $3d^{10}4s^2$
  - (c)  $3d^94s^2$
  - (d)  $3d^54s^24p^4$
3. Maximum number of unpaired electrons are in :
  - (a) Cr
  - (b) Mn
  - (c)  $Fe^{2+}$
  - (d) Co
4. Manganese exhibits maximum oxidation state in:
  - (a)  $K_2MnO_4$
  - (b)  $KMnO_4$
  - (c)  $MnO_2$
  - (d)  $Mn_3O_4$
5. General electronic configuration of d-block elements is :
  - (a)  $(n-1)d^{1-10}ns^{0-2}$
  - (b)  $(n-1)d^{1-10}ns^{1-2}$
  - (c)  $(n-1)d^{1-10}ns^0$
  - (d)  $(n-1)d^{1-10}ns^1$
6. Electronic configuration of d-block element 'X' in +3 oxidation state in  $[Ar] 3d^5$ , Atomic number of 'X' is :
  - (a) 25
  - (b) 26
  - (c) 27
  - (d) 24
7.  $CrO_3$  dissolves in aqueous NaOH to give:
  - (a)  $CrO_4^{2-}$
  - (b)  $Cr(OH)_3$
  - (c)  $Cr_2O_7^{2-}$
  - (d)  $Cr(OH)_2$
8. The electronic configuration of gadolinium (At. No. 64) is:
  - (a)  $[Xe] 4f^8 5d^0 6s^2$
  - (b)  $[Xe] 4f^7 5d^2 6s^2$
  - (c)  $[Xe] 4f^3 5d^5 6s^2$
  - (d)  $[Xe] 4f^6 5d^2 6s^2$

9. Electronic configuration of a transition element in + 3 oxidation state is  $[\text{Ar}] 3d^5$ . What is its atomic number?
- (a) 25 (b) 26  
(c) 27 (d) 24
10. On addition of small amount of  $\text{KMnO}_4$  to concentrated  $\text{H}_2\text{SO}_4$ , a green oily compound is obtained which is highly explosive in nature. Identify the compound from the following.
- (a)  $\text{Mn}_2\text{O}_7$  (b)  $\text{MnO}_2$   
(c)  $\text{Mn}_5\text{O}_4$  (d)  $\text{Mn}_2\text{O}_3$
11. Which of the following oxidation state is common for all lanthanoids?
- (a) +2 (b) +3  
(c) +4 (d) +5
12. When  $\text{KMnO}_4$  solution is added to oxalic acid solution, the decolourisation is slow in the beginning but becomes instantaneous after some time because:
- (a)  $\text{CO}_2$  is formed as the products  
(b) Reaction is exothermic  
(c)  $\text{MnO}_4^-$  catalysis the reaction  
(d)  $\text{Mn}^{2+}$  acts as autocatalyst
13.  $\text{KMnO}_4$  acts as an oxidising agent in acidic medium. The number of moles of  $\text{KMnO}_4$  that will be needed to react with one mole of sulphide ions in acidic solution is:
- (a) 2/5 (b) 3/5  
(c) 4/5 (d) 1/5
14. Which of the following is amphoteric oxide?  $\text{Mn}_2\text{O}_7$ ,  $\text{CrO}_3$ ,  $\text{Cr}_2\text{O}_3$ ,  $\text{CrO}_3$ ,  $\text{V}_2\text{O}_5$ ,  $\text{V}_2\text{O}_4$ .
- (a)  $\text{V}_2\text{O}_5$ ,  $\text{Cr}_2\text{O}_3$  (b)  $\text{Mn}_2\text{O}_7$ ,  $\text{CrO}_3$   
(c)  $\text{CrO}_3$ ,  $\text{V}_2\text{O}_5$  (d)  $\text{V}_2\text{O}_5$ ,  $\text{V}_2\text{O}_4$
15. The magnetic moment is associated with its spin angular momentum. Spin only magnetic moment value of  $\text{Cr}^{3+}$  ion is.....
- (a) 2.87 B.M. (b) 3.87 B.M.  
(c) 3.47 B.M. (d) 3.57 B.M.



16. Generally transition elements and their salts are coloured due to the product of unpaired electrons in metal ions. Which of the following compounds are coloured?
- (a)  $\text{KMnO}_4$  (b)  $\text{ZnSO}_4$   
 (c)  $\text{TiCl}_4$  (d)  $\text{Cu}_2\text{Cl}_2$
17. Transition elements show magnetic moment due to spin and orbital motion of electrons. Which of the following metallic ions have almost same spin only magnetic moment?
- (a)  $\text{Co}^{2+}, \text{Cr}^{2+}$  (b)  $\text{Cr}^{2+}, \text{Mn}^{2+}$   
 (c)  $\text{Mn}^{2+}, \text{Co}^{2+}$  (d)  $\text{Co}^{2+}, \text{Cr}^{3+}$
18. Which of the following actinoids show oxidation states upto + 7?
- (a) Am (b) Pu  
 (c) U (d) Th
19. Which of the following ions show highest spin only magnetic moment value?
- (a)  $\text{Ti}^{3+}$  (b)  $\text{Mn}^{2+}$   
 (c)  $\text{Fe}^{2+}$  (d)  $\text{Co}^{3+}$

## II FILL IN THE BLANKS

- Hybridisation of Cr in Chromate ion is.....
- Maximum oxidation state exhibited by Mn is.....
- Electronic configuration of  $\text{Cr}^+$  is.....
- Theoretical magnetic moment (spin - only) of  $\text{Ti}^{3+}$  ion is.....
- The equivalent weight of  $\text{K}_2\text{Cr}_2\text{O}_7$  in acidic medium is.....
- Last element in the actinide series is.....
- The general electronic configuration of d—block elements is.....
- The colour of  $\text{KMnO}_4$  is due to.....
- Out of  $\text{Sc}^{2+}$  and  $\text{Sc}^{3+}$  ion, ..... is paramagnetic in nature.
- The oxidation state of Mn in  $\text{K}_2\text{MnO}_4$  is .....

### III ASSERTION REASON TYPE QUESTIONS

In the following questions a statement of assertion followed by a statement of reason is given. Choose the correct answer out of the following choice.

- (a) Both assertion and reason are True, and reason is the correct explanation of the assertion.
  - (b) Both assertion and reason are True, but reason is not the correct explanation of the assertion.
  - (c) Assertion is true, but reason is false.
  - (d) Assertion is false, reason is true.
1. **ASSERTION :** Sc does not show variable oxidation states.  
**REASON :** Sc has only one electron in the 3d subshell.
  2. **ASSERTION :** Separation of Zr and Hf is difficult.  
**REASON :** Because Zr and Hf lie in the same period of the periodic table.
  3. **ASSERTION :** Actinoids form relatively less stable complexes as compared to lanthanoids.  
**REASON :** Actinoids can utilise their 5f orbitals along with 6d orbitals in bonding but lanthanoids do not use their 4f orbital for bonding.
  4. **ASSERTION :** Cu cannot liberate hydrogen from acids.  
**REASON :** Because it has positive electrode potential.
  5. **ASSERTION :** The highest oxidation state of osmium is + 8.  
**REASON :** Osmium is a 5d element.
  6. **ASSERTION :** Highest oxidation state is exhibited by transition metal lying in the middle of the series.  
**REASON :** The highest oxidation state exhibited corresponds to number of (n-1)d electrons.
  7. **ASSERTION :**  $\text{Fe}^{3+}$  is more stable than  $\text{Fe}^{2+}$   
**REASON :**  $\text{Fe}^{3+}$  has  $3d^5$  configuration while  $\text{Fe}^{2+}$  has  $3d^6$  configuration.
  8. **ASSERTION :** Vanadium has the ability to exhibit a wide range of oxidation states.  
**REASON :** The standard potentials Vanadium are rather small, making a switch between oxidation states relatively easy.
  9. **ASSERTION :** Transition metals like Fe, Cr and Mn form oxyions.  
**REASON :** Oxygen is highly electronegative and has a tendency to form multiple bonds.
  10. **ASSERTION :** The highest oxidation states of the 3d metals depends only on electronic configuration of the metal.  
**REASON :** The number of electrons in the (n-1)d and ns subshells determine the oxidation states exhibited by the metal.

### IV ONE WORD ANSWER TYPE QUESTIONS

1. Which element in 3d series shows highest number of oxidation states?
2. Out of cuprous and cupric ions which is coloured?
3. Out of Zn and Cr which is having higher first ionisation enthalpy?
4. Give general Electronic configuration of actinoids.
5. Name the element of 3d series which does not exhibit variable oxidation state.

6. What is the equivalent weight of  $\text{KMnO}_4$  in alkaline medium.
7. What is the maximum oxidation state shown by actinoids.
8. Out of  $\text{CuCl}_2$  and  $\text{Cu}_2\text{Cl}_2$  which is more stable?
9. What is the cause of similar size of Zr and Hf?

**VERY SHORT ANSWER TYPE QUESTIONS (1 Mark)**

Q.1. Explain  $\text{CuSO}_4 \cdot 5\text{H}_2\text{O}$  is blue while  $\text{CuSO}_4$  is colourless ?

Ans. Because water molecules act as ligands and results in crystal field splitting of d-orbitals of  $\text{Cu}^{2+}$  ion.

Q.2. Which element among 3d series exhibit highest oxidation state ?

Ans. Mn

Q.3. In 3d series (Sc to Zn), the enthalpy of atomization of Zn is low. Why ?

Ans. Due to absence of unpaired electrons.

Q.4. Which element among 3d series exhibit only one oxidation state ?

Ans. Sc

Q.5. Why is the 3rd ionization energy of Mn ( $Z = 25$ ) is unexpectedly high ?

Ans. Due to half-filled electronic configuration.

Q.6. Define alloy.

Ans. Alloys are homogeneous solid solutions of two or more metals.

Q.7. Transition metals show zero oxidation state with ligands like CO. Explain.

Ans. CO form synergic bonding with metal ion.

Q.8. Why HCl can not be used to acidify  $\text{KMnO}_4$  solution ?

Ans. Because  $\text{KMnO}_4$  oxidize HCl into  $\text{Cl}_2$ .

Q.9. Name one ore of Mn and Cr.

Ans. Mn :  $\text{MnO}_2$  Cr :  $\text{FeCr}_2\text{O}_4$

Q.10. Why  $\text{Mn}^{2+}$  compounds are more stable than  $\text{Fe}^{2+}$  compounds towards oxidation to their +3 state ?

Ans.  $\text{Mn}^{2+}$  has half-filled electronic configuration ( $d^5$ ).

Q.11. Why do transition metal (elements) show variable oxidation states ?

Ans. Due to presence of partially filled (n-1)d subshell in addition of ns subshell.

Q.12. Write any uses of pyrophoric alloy.

Ans. Making bullets, shells and lighter flints.

Q.13. Which is more basic -  $\text{La}(\text{OH})_3$  or  $\text{Lu}(\text{OH})_3$  ? Why ?

Ans.  $\text{La}(\text{OH})_3$ , due to lanthanoid contraction, lower size, more covalent character, least basic.

Q.14. Find out number of Cr - O - Cr bond/bonds in  $\text{Cr}_2\text{O}_7^{2-}$  ion.

Ans. 1

Q.15. Why is  $\text{Ce}^{3+}$  in aqueous solution a good oxidizing agent ?

Ans. Because Ce is most stable in  $\text{Ce}^{3+}$  state in aqueous solution.

Q.16. Why do Zr and Hf exhibit similar properties ?

Ans. Due to lanthanoid contraction.

Q.17. What is lanthanoid contraction ?

Ans. The regular steady decrease in the atomic or ionic radii of lanthanoids with increasing atomic number.

Q.18. Why is Cu ( $Z = 29$ ) considered a transition metal ?

Ans. Due to its partially filled d-orbital in  $\text{Cu}^{2+}$  state.

Q.19. Arrange the given in increasing order of acidic character :  $\text{CrO}_3$ ,  $\text{CrO}$ ,  $\text{Cr}_2\text{O}_3$ .

Ans.  $\text{CrO}_3 < \text{CrO} < \text{Cr}_2\text{O}_3$

#### SHORT ANSWER TYPE QUESTIONS (2 or 3 Marks)

Q. 1. Chromium is typical hard metal while mercury is a liquid. Explain why ?

Ans. Cr has five unpaired d-electrons. Hence metallic bonds are strong. In Hg, there is absence of unpaired electrons and size is larger.

Q.2. Why  $\text{KMnO}_4$  is used in cleaning surgical instruments in hospitals ?

Ans. This is because  $\text{KMnO}_4$  has a germicidal action.

Q.3. Most of the transition metals do not displace hydrogen from dilute acids, why ?

Ans. Due to their negative reduction potential.

Q.4. Explain why  $\text{Cu}^+$  is not stable in aqueous solution ?

Ans. Due to less negative  $\Delta_{\text{hyd}}H$  of  $\text{Cu}^+$  it cannot compensate 2nd ionization potential of Cu.

Q.5. Why is the highest oxidation state of a metal exhibited in its oxide or fluoride only ?

Ans. Oxygen and fluoride have small size and high electronegativity. They can oxidise the metal.

Q.6. Write electronic configuration of  $\text{Cu}^{2+}$  and  $\text{Co}^{2+}$ .

Ans.  $\text{Cu}^{2+} = [\text{Ar}] 3d^9 4s^0$   $\text{Co}^{2+} = [\text{Ar}] 3d^7$

Q.7. Briefly explain why electronic configuration of lanthanoids are not known with certainty ?

Ans. 4f/5d subshells are very close in energy. So electrons can jump from 4f to 5d or vice-versa.

Q.8. Why Zn, Cd, Hg are soft and have low melting point ?

Ans. Due to weak interatomic attraction/absence of unpaired electrons.

Q.9. Which of the following is/are transition element and why ? Zn, Cd, Ag, Fe, Ni

Ans. Fe, Ni, Ag

Q.10. What are interstitial compounds ? Give example.

Ans. When small atoms like C, H, B and N occupy interstitial site in their lattice.  
Example : TiC.



Q.11. Why is first ionization energy of 5d elements higher than those of 3d and 4d elements ?

Ans. Due to greater effective nuclear charge acting on outer most electrons because of poor shielding of 4f electrons.

Q.12. Explain 'Misch metal' and write its use.

Ans. It is an alloy of 95% lanthanoid and 5% iron and traces of S, C, Ca and Al. Used in lighter flint, bullet tips etc.

Q.13. Write balanced chemical equations :



Q.14. Out of Fe and Cu, which one would exhibit higher melting point ?

Ans. Fe, due to large number of unpaired d-electrons/more interatomic attraction.

Q.15. Sc, the first member of first transition series does not exhibit variable oxidation state. Why?

Ans. Due to noble gas electronic configuration in + 3 oxidation state no other oxidation state is stable.

Q.16. (a) Deduce the number of 3d electrons in the following ions :  $\text{Cu}^{2+}$ ,  $\text{Sc}^{3+}$

(b) Why do transition metals form alloy?

(c) Why  $\text{Zn}^{2+}$  salts are white?

Ans. (a)  $\text{Cu}^{2+}$  : 9 electrons :  $\text{Sc}^{3+}$  : 0 electron

(b) Transition metals have similar atomic radii.

(c) Absence of unpaired electron.

Q.17. (a) Why is separation of lanthanoid elements difficult?

(b) Transition metal exhibit higher enthalpies of atomization. Explain why?

(c) Why the transition metals have ability to form complexes?

Ans. (a) Due to lanthanide contraction, the size of these elements is nearly same.

(b) Transition metal contain large number of unpaired electrons, and they have strong interatomic attractions.

(c) Due to their small size and large nuclear charge.

Q.18. (a) Use Hund's rule to derive the electronic configuration of  $\text{Ce}^{3+}$  ions and calculate its magnetic moment.

(b) Is lanthanum a f-block element?

Ans. (a)  $\text{Ce}(58) = [\text{Xe}] 4f^1 5d^1 6s^2$ ;  $\text{Ce}^{3+} = 4f^1$ ,  $\mu = 1.73 \text{ BM}$   $\mu = \sqrt{n(n+2)}$

(b) No, it is a d-block element.

Q.19. Account for the following :

(a) Silver chloride dissolves in excess of  $\text{NH}_3$ .

(b) Cuprous chloride is diamagnetic while cupric chloride is paramagnetic.

(c) in  $\text{CrO}_4^{2-}$  ion, all the Cr-O bond length are equal.

Ans. (a)  $\text{AgCl}$  forms a soluble complex with  $\text{NH}_3$ ,  $(\text{AgCl} + 2\text{NH}_3 \rightarrow [\text{Ag}(\text{NH}_3)_2]\text{Cl})$

(b)  $\text{Cu}^+$  :  $3d^{10} 4s^0$  - All electrons are paired.  $\text{Cu}^{2+}$  :  $3d^9$  - Here, one unpaired electron is present.

(c) Due to resonance.

Q.20. The  $E^\circ$  values in respect of electrodes of Cr, Mn and Fe are :  $\text{Cr}^{3+}/\text{Cr}^{2+} = -0.4 \text{ V}$   
 $\text{Mn}^{3+}/\text{Mn}^{2+} = +1.5 \text{ V}$   $\text{Fe}^{3+}/\text{Fe}^{2+} = +0.8 \text{ V}$  Compare the feasibilities of further oxidation of these ions.

Ans.  $\text{Cr}^{3+}$  is more stable than  $\text{Cr}^{2+}$ .  $\text{Mn}^{2+}$  is more stable than  $\text{Mn}^{3+}$ .  $\text{Fe}^{3+}$  is more stable than  $\text{Fe}^{2+}$ . Order of feasibility of + 2 oxidation state is :  $\text{Mn}^{2+} > \text{Fe}^{2+} > \text{Cr}^{2+}$

Q.21. Write any three properties of interstitial compounds.

- Ans. (a) They are chemically inert.  
 (b) They retain metallic conductivity.  
 (c) They have high melting point than their pure metals.  
 (d) These are harder and more corrosion resistant.

Q.22. Account for the following :



- (a) All Scandium salts are white.  
 (b) The first ionization energy of the 5d series are higher than 3d and 4d transition elements in respective groups.  
 (c)  $\text{Ce}^{3+}$  can be easily oxidized to  $\text{Ce}^{4+}$ .

- Ans. (a) Sc has only +3 oxidation state, there is no unpaired electron.  
 (b) Due to lanthanoid contraction, effective nuclear charge increase.  
 (c) Due to attainment of noble gas electronic configuration.

### LONG ANSWER TYPE QUESTIONS (5 Marks)

- Q. 1. (a) What is meant by disproportionation of an oxidation state ? Give one example.  
 (b) Draw the structures of  $\text{Cr}_2\text{O}_7^{2-}$ ,  $\text{CrO}_4^{2-}$ ,  $\text{MnO}_4^-$ .  
 (c) What is the effect of lanthanoid contraction beyond lanthanoid ?

- Ans. (a) When any atom or ion undergo oxidation and reduction simultaneously it is called disproportionation.



- (c) Size of respective 4d and 5d series elements becomes comparable from fourth group onwards (e.g., Zr and Hf).

2. On the basis of lanthanoid contraction, explain the following:

- (i) Nature of bonding in  $\text{La}_2\text{O}_3$  and  $\text{Lu}_2\text{O}_3$ .
- (ii) Trends in the stability of oxo salts of lanthanoids from La to Lu.
- (iii) Stability of the complexes of lanthanoids.
- (iv) Radii of 4d and 5d block elements.
- (v) Trends in acidic character of lanthanoid oxides.

Ans. (i) As the size decreases covalent character increases. Therefore,  $\text{La}_2\text{O}_3$  is more ionic and  $\text{Lu}_2\text{O}_3$  is more covalent.

(ii) As the size decreases from La to Lu, stability of oxosalts also decreases.

(iii) Stability of complexes increases as the size of lanthanoids decreases.

(iv) Radii of 4d and 5d block elements will be almost same.

(v) Acidic character of Oxides increases from La to Lu.

3. (a) Answer the following questions:

(i) Which element of the first transition series has highest second ionisation enthalpy?

(ii) Which element of the first transition series has highest third ionisation enthalpy?

(iii) Which element of the first transition series has lowest enthalpy of atomisation?

(b) Identify the metal and justify your answer.

(i) Carbonyl  $\text{M}(\text{CO})_5$

(ii)  $\text{MO}_3\text{F}$

Ans. (a)

(i) Cu, because the electronic configuration of Cu is  $3d^{10}4s^1$ . So second electron needs to be removed from completely filled d-orbital.

(ii) Zn [Hint : As above].

(iii) Zn [Hint : No unpaired electron for metallic bonding]

(b) (i)  $\text{Fe}(\text{CO})_5$

(ii)  $\text{MnO}_3\text{F}$  [Hint : Mn shows + 7 oxidation state; d-electrons are not involved in bonding.]



4. (i) How would you account for the following?
- The oxidising power of oxo-anions are in the order  $\text{VO}^{2+} < \text{Cr}_2\text{O}_7^{2-} < \text{MnO}_4^-$
  - The third ionisation enthalpy of manganese ( $Z = 25$ ) is exceptionally high.
  - $\text{Cr}^{3+}$  is a stronger reducing agent than  $\text{Fe}^{2+}$ .
- (ii) Give reasons for the following:
- $\text{Mn}^{3+}$  is a good oxidising agent.
  - $E^\circ(\text{M}^{2+}/\text{M})$  values are not regular for first row transition metals (3d-series).
  - Although 'F' is more electronegative than 'O' the highest Mn fluoride is  $\text{MnF}_4$ , whereas the highest oxide is  $\text{Mn}_2\text{O}_7$ , reducing agent than  $\text{Fe}^{2+}$ .

- Ans (i) (a) It is due to the fact that V in its lower oxidation state is less stable than Cr which in turn is less stable than Mn. Thus,  $\text{MnO}_4^-$  has a great tendency to get reduced and hence, behave as a good oxidising agent. Similarly  $\text{VO}^{2+}$  has the least oxidising power.
- (b) The third ionisation enthalpy of Mn is very high due to the fact that the third electron has to be removed from stable half-filled configuration, i.e.,  $3d^5$ .
- (c)  $\text{Cr}^{2+}$  is stronger reducing agent than  $\text{Fe}^{2+}$  because  $d^4 \rightarrow d^3$  transition occurs in case of  $\text{Cr}^{2+}$  to  $\text{Cr}^{3+}$  while  $d^6 \rightarrow d^5$  transition occurs in case of  $\text{Fe}^{2+}$  to  $\text{Fe}^{3+}$ . In a medium like water  $d^3$  is more stable as compared to  $d^5$ .
- (ii) (a)  $\text{Mn}^{3+}$  ( $3d^4$ ) is a good electron acceptor as the resulting species is more stable ( $3d^5$ ).
- (b) The  $E^\circ(\text{M}^{2+}/\text{M})$  values are not regular which can be explained from the irregular variation of ionisation enthalpies ( $\Delta_i H_1 + \Delta_i H_2$ ) and also the sublimation enthalpies which are relatively much less for Mn and V.
- (c) Due to multiple bond formation ability of oxygen, Mn can form  $\text{Mn}_2\text{O}_7$ .


5. The elements of 3d-transition series are given as:

**Sc, Ti, V, Cr, Mn, Fe, Co, Ni, Cu, Zn**

**Answer the following :**

- Name the element which shows maximum number of oxidation states. Give reason.
- Which element has the highest melting point?
- Name the element which shows only +3 oxidation state?
- Which element is a strong oxidizing agent in +3 oxidation state and why?

- Ans (I) Mn. It has maximum unpaired electrons.  
 (ii) Cr  
 (iii) Sc  
 (iv) Manganese.  $\text{Mn}^{3+}$  to  $\text{Mn}^{2+}$  results in the stable half filled ( $d^5$ ) configuration.
6. On the basis of the figure given below, answer the following questions:



- Why Manganese has lower melting point than Chromium.
  - Why do transition metals of 3d series have lower melting points as compared to 4d series?
  - In the third transition series, identify and name the metal with the highest melting point.
  - Which element is a strong oxidizing agent in +3 oxidation state and why?
- Ans. (i) Manganese is having lower m.p. as compared to Chromium, as it has highest number of unpaired electrons, strong interatomic metallic bonding, hence no delocalization of electrons.  
 (ii) Due to more frequent metal-metal bonding in compounds of heavy transition metals i.e. 4d and 5d series.  
 (iii) tungsten

### CASE STUDY BASED QUESTIONS

#### 1. Read the passage and answer the following questions.

Potassium dichromate is one of the crystalline inorganic chemical reagents. Hexavalent chromium compounds are harmful to health.  $\text{K}_2\text{Cr}_2\text{O}_7$  is widely used in laboratories and industry as an oxidizing agent because it is not deliquescent. Potassium dichromate looks very bright and red-orange color. In this work different amount of acidic (HCl) and alkaline (NaOH) solutions were added to stock solutions of  $\text{K}_2\text{CrO}_4$  and  $\text{K}_2\text{Cr}_2\text{O}_7$  to show the effect of pH values on their spectra. The results of UV-Visible spectroscopy shows that, the changing of solution pH value when drops of HCl were added led to shift wavelength of  $\text{K}_2\text{CrO}_4$  spectrum while no change has been occurred in  $\text{K}_2\text{Cr}_2\text{O}_7$  spectrum. However, Changing pH values solution by adding drops of NaOH led to change in wavelength red shift for  $\text{K}_2\text{Cr}_2\text{O}_7$  while no changes has been occurred in spectrum of  $\text{K}_2\text{CrO}_4$ .

**Reference:** Effect of the Acidic and Alkaline Solutions on  $\text{K}_2\text{CrO}_4$  and  $\text{K}_2\text{Cr}_2\text{O}_7$  by Ultraviolet and Visible Measurement Mohammad Radi Mohammad, Hasanain Saad Azeez\* Al-Mustansiriyah Journal of Science ISSN: 1814-635X (print), ISSN:2521-3520 (online) Volume 30, Issue 1, 2019, 221-224

- (A) The hybridization of Cr in dichromate ion is :  
 (a)  $d^2sp^3$  (b)  $sp^3$   
 (c)  $dsp^2$  (d)  $sp^3d$
- (B) Colour of potassium dichromate is :  
 (a) purple (b) green  
 (c) yellow (d) orange
- (C) Chemical formula of ferrochrome is :  
 (a)  $FeCrO_2$  (b)  $FeO.Cr_2O_3$   
 (c)  $Fe_2CrO_4$  (d) None of these
- (D) On increasing pH of dichromate, it converts in :  
 (a)  $CrO_4^{2-}$  (b)  $Cr_2O_4^{2-}$   
 (c)  $CrO_4^{-1}$  (d)  $CrO_3$

**2. Read the passage and answer the following questions.**

Heavy rare earth elements crystallize into hexagonally close packed (h.c.p.) structures and share a common outer electronic configuration, differing only in the number of  $4f$  electrons they have. These chemically inert  $4f$  electrons set up localized magnetic moments, which are coupled via an indirect exchange interaction involving the conduction electrons. This leads to the formation of a wide variety of magnetic structures, the periodicities of which are often incommensurate with the underlying crystal lattice. Such incommensurate ordering is associated with a 'webbed' topology of the momentum space surface separating the occupied and unoccupied electron states (the Fermi surface). The shape of this surface—and hence the magnetic structure—for the heavy rare earth elements is known to depend on the ratio of the interplanar spacing  $c$  and the interatomic, intraplanar spacing  $a$  of the h.c.p. lattice. A theoretical understanding of this problem is, however, far from complete. Here, using gadolinium as a prototype for all the heavy rare earth elements, we generate a unified magnetic phase diagram, which unequivocally links the magnetic structures of the heavy rare earths to their lattice parameters. In addition to verifying the importance of the  $c/a$  ratio, we find that the atomic unit cell volume plays a separate, distinct role in determining the magnetic properties: we show that the trend from ferromagnetism to incommensurate ordering as atomic number increases is connected to the concomitant decrease in unit cell volume. This

volume decrease occurs because of the so-called lanthanide contraction, where the addition of electrons to the poorly shielding 4f orbitals leads to an increase in effective nuclear charge and, correspondingly, a decrease in ionic radii.

**Reference:**

**Lanthanide contraction and magnetism in the heavy rare earth elements.**

Hughes, I., Dane, M., Ernst, A. et al.

Nature 446,650-653 (2007). <https://doi.org/10.1038/nature05668>

- (A) Rare earth elements are also called :
- |                   |                   |
|-------------------|-------------------|
| (a) Actinoids     | (d) Lanthanoids   |
| (c) Alkali metals | (d) None of these |
- (B) Radioactive lanthanoids is :
- |        |        |
|--------|--------|
| (a) Pm | (b) Ce |
| (c) Nd | (d) Er |
- (C) Cause of lanthanoid contraction is :
- |                                   |                                   |
|-----------------------------------|-----------------------------------|
| (a) poor shielding of 5f orbitals | (b) poor shielding of 4f orbitals |
| (c) poor shielding of 6f orbitals | (d) All of these                  |
- (D) The common oxidation state of lanthanoids is :
- |        |        |
|--------|--------|
| (a) +4 | (b) +3 |
| (c) +2 | (d) +1 |

3. **Read the passage given below and answer the following questions:**

The d block elements are the 40 elements contained in the four rows of ten columns (3-12) in the periodic table. As all the d block elements are metallic, the term d-block metals is synonymous. This set of d-block elements is also often identified as the transition metals, but sometimes the group 12 elements (zinc, cadmium, mercury) are excluded from the transition metals as the transition elements are defined as those with partly filled d or f shells in their compounds. Inclusion of the elements zinc, cadmium and mercury is necessary as some properties of the group 12 elements are appropriate logically to include with a discussion of transition metal chemistry.

The term transition element or transition metal appeared to derive from early studies of periodicity such as the Mendeleev periodic table of the elements. His horizontal table of the elements was an attempt to group the elements together so that the chemistry of elements might be explained and predicted. In this table there are eight groups labeled 1-VIII with each subdivided into A and B subgroups. Mendeleev recognized that certain properties of elements in Group VIII are related to those of some of the elements in Group VII and those at the start of the next row Group I. In that sense, these elements might be described as possessing properties transitional from one row of the table to the next.

**Reference:** Winter, M. J. **d-Block Chemistry**

(Vol. 27). Oxford University Press, USA.)

**In the following questions, a statement of assertion followed by a statement of reason is given. Choose the correct answer out of the following choices on the basis of the above passage.**

- (a) Assertion and reason both are correct statements and reason is correct explanation for assertion.
- (b) Assertion and reason both are correct statements but reason is not correct explanation for assertion.
- (c) Assertion is correct statement but reason is wrong statement,
- (d) Assertion is wrong statement but reason is correct statement.
- (A) Assertion: Group 12 elements are not considered as transition metals.  
Reason: Transition metals are those which have incompletely filled d shell in their compounds.
- (B) Assertion: All d block elements are metallic in nature.  
Reason: The d —block elements belong to Group 3 -12 of the periodic table.
- (C) Assertion: Nickel is a transition element that belongs to group 10 and period 4 of the modern periodic table.

Reason: Electronic configuration of Nickel is  $[\text{Ar}] 3d^8 4s^2$

4. **Read the passage given below and answer the following questions:**

The transition metals when exposed to oxygen at low and intermediate temperatures form thin, protective oxide films of up to some thousands of Angstroms in thickness. Transition metal oxides lie between the extremes of ionic and covalent binary compounds formed by elements from the left or right side of the periodic table. They range from metallic to semiconducting and deviate by both large and small degrees from stoichiometry. Since d-electron bonding levels are involved, the cations exist in various valence states and hence give rise to a large number of oxides. The crystal structures are often classified by considering a cubic or hexagonal close-packed lattice of one set of ions with the other set of ions filling the octahedral or tetrahedral interstices. The actual oxide structures, however, generally show departures from such regular arrays due in part to distortions caused by packing of ions of different size and to ligand field effects. These distortions depend not only on the number of d-electrons but also on the valence and the position of the transition metal in a period or group.

**Reference:** Smeltzer, W.W., & Young, D. J. (1975). Oxidation properties of transition metals. Progress in Solid State Chemistry, 10,17-54.)

**In the following questions, a statement of assertion followed by a statement of reason is given. Choose the correct answer out of the following choices on the basis of the above passage.**

- (a) Assertion and reason both are correct statements and reason is correct explanation for assertion.
- (b) Assertion and reason both are correct statements but reason is not correct explanation for assertion.
- (c) Assertion is correct statement but reason is wrong statement,
- (d) Assertion is wrong statement but reason is correct statement.
- (A) Assertion: Cations of transition elements occur in various oxidation states  
Reason: Transition metals are those which have incompletely filled d subshell in their compounds.

- (B) Assertion: Crystal structure of oxides of transition metals often show defects.  
Reason: Ligand field effect cause distortions in crystal **structures**.
- (C) Assertion : Transition metals form protective oxide films.  
Reason: Oxides of transition metals are always stoichiometric.
- (D) Assertion: CrO crystallises in a hexagonal close-packed array of oxide ions with two out of every three octahedral holes occupied by chromium ions  
Reason: Transition metal oxide may be hexagonal close-packed lattice of oxide ions with metal ions filling the octahedral voids.

### ANSWERS

#### I. MULTIPLE CHOICE TYPE QUESTIONS

1. d 2. b 3. a 4. a 5. a 6. b 7. a 8. b 9. b 10. a  
11. b 12. d 13. a 14. a 15. b 16. a 17. d 18. b 19. b

#### II. FILL IN THE BLANKS

- |                             |                            |
|-----------------------------|----------------------------|
| 1. $sp^3$                   | 2. +7                      |
| 3. $[Ar]3d^5 4s^1$          | 4. 1.732                   |
| 5. 49g                      | 6. lawrencium              |
| 7. $(n-1)d^{1-10} ns^{0-2}$ | 8. charge transfer complex |
| 9. $Sc^{2+}$                | 10. 6                      |

#### III. ASSERTION REASON TYPE QUESTIONS

- 1.(b) 2.(b) 3.(c) 4.(a) 5.(b) 6.(c) 7.(a) 8.(a) 9.(b) 10.(d)

#### CASE STUDY BASED QUESTIONS

##### I.

- |       |       |
|-------|-------|
| (A) b | (B) d |
| (C) b | (D) a |

##### II.

- |       |       |
|-------|-------|
| (A) b | (B) a |
| (C) b | (D) b |

##### III.

- |       |       |
|-------|-------|
| (A) a | (B) b |
| (C) a |       |

##### IV.

- |       |       |
|-------|-------|
| (A) b | (B) a |
| (C) c | (D) d |

#### IV. ONE WORD TYPE QUESTIONS

- |        |           |             |                              |
|--------|-----------|-------------|------------------------------|
| 1. Mn  | 2. Cupric | 3. Zn       | 4. $5f^{1-14} 6d^{0-1} 7s^2$ |
| 5. Sc  |           |             |                              |
| 6. 58g | 7. 7      | 8. $CuCl_2$ | 9. Lanthanoid Contraction    |

## UNIT TEST-1

## d- and f- block Elements

Maximum marks: 20

Time Allowed: 1 Hour

1. Which element among 3d-transition elements, exhibit the highest oxidation state? (1)
2. Name the transition element which has highest  $E^\circ(M^{2+}/M)$  value (1)
3. Calculate the magnetic moment of  $Cu^{2+}$  ( $Z = 29$ ) on the basis of "spin-only" formula. (1)
4. Name a transition element which does not exhibit variable oxidation state in 3d series. (1)
5. Write the general electronic configuration of d-block elements. (1)
6. Write balanced chemical equations for:
  - (a) Oxidation of  $Fe^{2+}$  by  $Cr_2O_7^{2-}$  in acidic medium
  - (b) Oxidation of  $Mn^{2+}$  by  $MnO_4^-$  in neutral or faintly alkaline medium. (2)
7. Account for the following:
  - (a) Copper shows its inability to liberate hydrogen gas from the dilute acids.
  - (b) Scandium ( $Z = 21$ ) does not exhibit variable oxidation states. (2)
8. Explain lanthanoid contraction with its consequences. (2)
9. Assign reasons for the following: (3)
  - (a) Majority of transition metals form complexes.
  - (b)  $Ce^{3+}$  can be easily oxidised to  $Ce^{4+}$ .
  - (c) Actinoids display a variety of oxidation states.
10. Describe the preparation of potassium permanganate ( $KMnO_4$ ). Write the chemical equations involved in the synthesis. (3)
11. Explain giving reasons: (3)
  - (a) Zn, Cd and Hg are not considered as transition metals.
  - (b) Elements in the middle of transition series have higher melting points.
  - (c) The decrease in atomic size of transition elements in a series is very small.



## UNIT TEST-2

## d- and f- block Elements

Maximum marks: 20

Time Allowed: 1 Hour

1. Explain- zinc is not regarded is transition element. 1
2. Name a lanthanoid well known to exhibit +4 oxidation state. 1
3. Out of  $\text{Sc}^{3+}$ ,  $\text{Co}^{2+}$  and  $\text{Cr}^{3+}$  ions, which ion will be colourless in aqueous solutions?  
(Atomic no. : Co=27, Sc=21 and Cr=24) 1
4. Write general electronic configuration of lanthanoid series. 1
5. Why Zr(Z=40) and Hf(Z=72) shows similar properties? 1
6. Name the 3d element which: 2
  - a) Does not exhibit variable oxidation state
  - b) Exhibits highest oxidation state
  - c) Has highest spin-only magnetic moment in +2 oxidation state
  - d) Has highest  $E^\circ (\text{M}^{2+} | \text{M})$  value.
7. Write any two differences between lanthanoids and actinoids.
8. Explain disproportionation reaction giving one example of a compound/ion containing transition element. 2
9. Explain following in context of transition elements:
  - a) High enthalpy of atomisation
  - b) Catalytic activity
  - c) Coloured complexes
10. Write balanced chemical equations for the following reactions: 3
  - a)  $\text{MnO}_4^- + \text{S}_2\text{O}_3^{2-} + \text{H}_2\text{O} \rightarrow$
  - b)  $\text{Cr}_2\text{O}_7^{2-} + \text{Sn}^{2+} + 4\text{H}^+ \rightarrow$
  - c)  $\text{Fe}^{2+} + \text{MnO}_4^- + 8\text{H}^+ \rightarrow$
11. When  $\text{MnO}_2$  is fused with  $\text{KOH}$  and  $\text{KNO}_3$  (oxidising agent) it gives a dark compound (A). Compound (A) disproportionates in acidic solution to give purple coloured compound (B). An alkaline solution of compound (B) oxidises  $\text{KI}$  to compound (C), whereas an acidified solution of compound (B) oxidises  $\text{KI}$  to (D). Identify A,B,C and D and write reactions involved. 3




## UNIT 9

# Coordination Compounds

### Points to Remember


**Introduction :** Complex compounds or coordination compounds are those molecular compounds which retain their identity in solid as well as in solution are known as complex compounds.

Example,




### Ligands

The ions or molecules bound to the central atom/ion in the coordination entity are called ligands.



Homoleptic Complexes	Heteroleptic Complexes
Complexes in which a metal is bound to only one kind of donor groups are known as homoleptic.	Complexes in which a metal is bound to more than one kind of donor groups are known as heteroleptic.
e.g., $[\text{Co}(\text{NH}_3)_6]^{3+}$	e.g., $[\text{Co}(\text{NH}_3)_4\text{Cl}_2]^+$

## Nomenclature of Coordination Compounds



## Isomerism in Coordination Compounds

**Note:** Stereo isomerism and structural isomerism are the two principal types of isomerisms which are known among coordination compounds.

### Stereo Isomerism

It occurs due to different arrangements of ligands around central metal atom. It is of two types :

**Geometrical  
Isomerism**

**Optical Isomerism.**

### Optical Isomerism

- It arises when mirror images cannot be superimposed on one another. These mirror images are called as enantiomers. The two forms are called dextro (d) and laevo (l).
- Optical isomerism is common in octahedral complexes but at least one didentate ligand must be present.  
e.g.,  $[\text{Co}(\text{en})_3]^{3+}$ ,  $[\text{PtCl}_2(\text{en})_2]^{2+}$  etc.

## Geometrical Isomerism

It arises in heteroleptic complexes due to different possible geometric arrangements of the ligands. Important examples of this behavior are found in square planar and octahedral complexes.


Square planar complex of formula  $[MX_2L_2]$  (X and L are unidentate), The two ligands X may be arranged adjacent to each other in a cis isomer, or opposite to each other in a trans isomer.  
e.g.,  $[Pt(NH_3)_2Cl_2]$

Square planar complex of the type  $[MABXL]$  (where A, B, X, L are unidentates) shows three isomers - two cis and one trans. Such isomerism is not possible for tetrahedral geometry.  
e.g.,  $[Pt(NH_3)(Br)(Cl)(Py)]$

Octahedral complexes of formula  $[MX_2A_4]$  where X are unidentates and A are didentate and form cis and trans isomers.  
e.g.,  $[CoCl_2(en)_2]$

Octahedral complexes of formula  $[MX_2L_4]$  in which the two ligands X may be oriented cis or trans to each other.  
e.g.,  $[Co(NH_3)_4Cl_2]^+$

Octahedral coordination entities of the type  $[Ma_3b_3]$  like  $[Co(NH_3)_3(NO_2)_3]$ . If three donor atoms of the same ligands occupy adjacent positions at the corners of an octahedral face, we have the facial (fac) isomer. When the positions are around the meridian of the octahedron, we get the meridional (mer) isomer.



## Bonding in Coordination Compounds

### Werner's Theory

- (i) In complex compounds, metal atom exhibit two types of valencies - primary valency and secondary valency.
- (ii) Primary valencies are satisfied by anions only while secondary valencies are satisfied by ligands. Primary valency depends upon oxidation number of central metal atom while secondary valency represents coordination number of central metal atom.
- (iii) Primary valencies are ionisable and are non-directional while secondary valencies are non- ionisable and directional. Therefore, geometry of complex is decided by secondary valencies.

### Valence Bond Theory

According to this theory, the metal atom or ion under the influence of ligands form inner orbital and outer orbital complex. These hybridized orbitals are allowed to overlap with ligand orbitals that can donate electron pairs for bonding.

- (i) **Six ligands (unidentate) (octahedral entity)** Generally central atom belongs 3d series and ligands can be monodentate or didentate but coordination number should be six and shape of complexes will be octahedral and form two types of complexes.

#### Inner orbital complexes,


Which are formed due to participation of (n-1)d orbitals in hybridisation is ( $d^2sp^3$ ) and shape of complex will be octahedral.

#### Outer orbital complexes,

Which are formed due to participation of nd orbitals in hybridisation is ( $sp^3d^2$ ).


Generally halides ( $F^-$ ,  $Cl^-$ ,  $Br^-$ ,  $I^-$ ),  $SCN^-$ ,  $S^{2-}$  form outer orbital complexes and other ligands form inner orbital complexes.

e.g., Inner orbital complex,  $[Co(NH_3)_6]^{3+}$



All electrons are paired therefore, complex will be diamagnetic in nature.


e.g., Outer orbital complex,  $[CoF_6]^{3-}$




Complex has unpaired electrons, therefore, complex will be paramagnetic in nature.



Complexes with coordination number: 4



All electrons are paired. Complex will be diamagnetic in nature.




Complex has unpaired electrons. Complex will be paramagnetic in nature.

### Crystal Field Theory

The five d-orbitals are split into lower and higher energy level due to approach of ligands is known as crystal field theory. The five d-orbitals in a gaseous metal atom/ion have same energy.

(i) Crystal field splitting in octahedral coordination entities




- Energy separation is denoted by  $\Delta_o$  (the subscript o is for octahedral).
- The energy of the two  $e_g$  orbitals (higher energy orbitals) will increase by  $(3/5)\Delta_o$ , and that of the three  $t_{2g}$  (lower energy orbitals) will decrease by  $(2/5)\Delta_o$ .
- If  $\Delta_o < P$ , the fourth electron enters one of the  $e_g$  orbitals giving the configuration  $t_{2g}^3 e_g^1$ . Ligands for which  $\Delta_o < P$  are known as weak field ligands and form high spin complexes.

- If  $\Delta_o < P$ , it becomes more energetically favourable for the fourth electron to occupy  $t_{2g}$  orbital with configuration  $t_{2g}^4 e_g^0$ .

Ligands which produce this effect are known as strong field ligands which produce this effect are known as strong field ligands which produce this effect are known as strong field ligands and form low spin complexes, where  $P$  represents the energy required for electron pairing in a single orbital.

(ii) Crystal field splitting in tetrahedral coordination entities.



- In tetrahedral coordination entities,  $\Delta_t = (4/9)\Delta_o$ .

Consequently the orbital splitting energies are not sufficiently large for forcing pairing, therefore, low spin configurations are rarely observed.

- Due to less crystal field stabilization energy, it is not possible to pair electrons and so all the tetrahedral complexes are high spin.

## 8. Colour in Coordination Compounds

- In complex compounds d-orbitals split in two sets  $t_{2g}$  and  $e_g$ . These have different energies. The difference in energies lies in visible region and electron jump from ground state  $t_{2g}$  level to higher state  $e_g$  level. This is known as d-d transition and it is responsible for colour of coordination compounds.
- d-d transition takes place in  $d^1$  to  $d^9$  ions, so the ions having  $d^1$  to  $d^9$  configuration are coloured. On the other hand, the ions  $d^0$  and  $d^{10}$  configuration do not show d-d transition.

### Importance and Applications of Coordination Compounds.


- Hardness of water is estimated by simple titration with  $\text{Na}_2\text{EDTA}$ . The  $\text{Ca}^{2+}$  and  $\text{Mg}^{2+}$  ions form stable complexes with EDTA.
- Some important extraction processes of metals, like those of silver and gold make use of complex formation.
- Similarly, purification of metals can be achieved through formation and subsequent decomposition of their coordination compounds. For example, impure nickel is converted to  $[\text{Ni}(\text{CO})_4]$ , which is decomposed to yield pure nickel.
- Coordination compounds are used as catalysts for many industrial processes. Examples include rhodium.

### Supplementary List of Ligands

Ligand	$\text{F}^-$	$\text{Cl}^-$	$\text{Br}^-$	$\text{I}^-$	$\text{OH}^-$	$\text{CN}^-$	$\text{O}^{2-}$	$\text{O}_2^{2-}$	$\text{O}_2^-$	$\text{CO}_3^{2-}$	$\text{PH}_3$
Name	fluoro	chloro	bromo	iodo	hydroxo	cyano	oxo	peroxo	super oxo	carbonato	phosphine
Ligand	$\text{SO}_4^{2-}$	$\text{NO}_2^-$	$\text{ONO}^-$	$\text{SCN}^-$	$\text{NCS}$	$\text{CH}_3\text{COO}^-$	$\text{C}_5\text{H}_5\text{N}$	$\text{S}^{2-}$	$\text{S}_2\text{O}_3^{2-}$	$\text{NO}_3^-$	$\text{SO}_3^{2-}$
Name	sulphato	nitro	nitrito	thiocyanato	isothiocyanato	acetate	pyridine (Py)	sulphido	thiosulphato	nitrate	sulphite
Ligand	$\text{NC}^-$	$(\text{C}_6\text{H}_5)_3\text{P}$	$\text{CS}$	$\text{NH}_2^-$	$\text{NH}^{2-}$	$\text{H}_2\text{NCSNH}_2$	$\text{C}_2\text{O}_4^{2-}$	$\text{H}_2\text{O}$	$\text{NH}_3$	$\text{CO}$	$\text{NO}$
Name	isocyanato	triphenyl phosphine	thiocarbonyl	amido	imido	thiourea (tu)	oxalate (ox)	aqua	ammine	carbonyl	nitrosyl




ethane-1, 2- diamine (en)



Ethylenediaminetriacetate ion  $\text{EDTA}^{3-}$



Diethylenetriamine(diene)



Ethylenediaminetetracetate ion  $\text{EDTA}^{4-}$

## OBJECTIVE TYPE QUESTIONS

### I. MULTIPLE CHOICE QUESTIONS

1. Which of the following compounds formed by  $\text{Cu}^{2+}$  ions is most stable?
  - (a)  $\text{Cu}^{2+} + 4\text{NH}_3 \rightarrow [\text{Cu}(\text{NH}_3)_4]^{2+}$ ;  $\log K = 11.6$
  - (b)  $\text{Cu}^{2+} + 4\text{CN}^- \rightarrow [\text{Cu}(\text{CN})_4]^{2-}$ ;  $\log K = 27.3$
  - (c)  $\text{Cu}^{2+} + 2\text{en} \rightarrow [\text{Cu}(\text{en})_2]^{2+}$ ;  $\log K = 15.4$
  - (d)  $\text{Cu}^{2+} + 4\text{H}_2\text{O} \rightarrow [\text{Cu}(\text{H}_2\text{O})_4]^{2+}$ ;  $\log K = 8.9$
2. The colour of the coordination compounds depends on the crystals field splitting. What will be the correct order of absorption of wavelength of light in the visible region for the complexes  $[\text{Co}(\text{NH}_3)_6]^{3+}$ ;  $[\text{Co}(\text{CN})_6]^{3-}$ ;  $[\text{Co}(\text{H}_2\text{O})_6]^{3+}$ .
  - (a)  $[\text{Co}(\text{CN})_6]^{3-} > [\text{Co}(\text{NH}_3)_6]^{3+} > [\text{Co}(\text{H}_2\text{O})_6]^{3+}$
  - (b)  $[\text{Co}(\text{NH}_3)_6]^{3+} > [\text{Co}(\text{H}_2\text{O})_6]^{3+} > [\text{Co}(\text{CN})_6]^{3-}$
  - (c)  $[\text{Co}(\text{H}_2\text{O})_6]^{3+} > [\text{Co}(\text{NH}_3)_6]^{3+} > [\text{Co}(\text{CN})_6]^{3-}$
  - (d)  $[\text{Co}(\text{CN})_6]^{3-} > [\text{Co}(\text{NH}_3)_6]^{3+} > [\text{Co}(\text{H}_2\text{O})_6]^{3+}$
3. When 0.1 mol  $\text{CoCl}_3 \cdot 5\text{NH}_3$  is treated with excess of  $\text{AgNO}_3$ , 0.2 mol of  $\text{AgCl}$  are obtained. The conductivity of solution will correspond to-
  - (a) 1 : 3 electrolyte                      (b) 1 : 2 electrolyte
  - (c) 1 : 1 electrolyte                      (d) 3 : 1 electrolyte
4. The correct IUPAC name of  $[\text{Pt}(\text{NH}_3)_2\text{Cl}_2]$  is-
  - (a) Diamminedichloridoplatinum (II)
  - (b) Diamminedichloridoplatinum (IV)
  - (c) Diamminedichloridoplatinum (0)
  - (d) Diamminedichloridoplatinum (IV)
5. Out of the following is the most stable complex species is
  - (a)  $[\text{Fe}(\text{CO})_5]$                       (b)  $[\text{Fe}(\text{CN})_6]^{3-}$
  - (c)  $[\text{Fe}(\text{C}_2\text{O}_4)_3]^{3-}$                       (d)  $[\text{Fe}(\text{H}_2\text{O})_6]^{3+}$
6. The CFSE for octahedral  $[\text{CaCl}_6]^{4-}$  is  $18,000 \text{ cm}^{-1}$ . The CFSE for tetrahedral  $[\text{CoCl}_4]^{2-}$  will be:
  - (a)  $18,000 \text{ cm}^{-1}$                       (b)  $16,000 \text{ cm}^{-1}$
  - (c)  $8,000 \text{ cm}^{-1}$                       (d)  $20,000 \text{ cm}^{-1}$

7. Which of the following species is not likely to be a ligand?
  - (a) NO
  - (b)  $\text{NH}_4^+$
  - (c)  $\text{NH}_2\text{CH}_2\text{NH}_2$
  - (d) CO
8. IUPAC name of  $[\text{Pt}(\text{NH}_3)_2\text{Cl}(\text{NO}_2)]$  is:
  - (a) Platinum diaminechloritrite
  - (b) Chloronitrito-N-ammine platinum(II)
  - (c) Diamminechloridonitrito-N-platinum (II)
  - (d) Diamminechlornitrite-N-platinate(II)
9. Atomic number of Mn, Fe and Co are 25, 26 and 27 respectively. Which of the following inner orbital octahedral complex ions are diamagnetic?
  - (a)  $[\text{Co}(\text{NH}_3)_6]^{3+}$
  - (b)  $[\text{Mn}(\text{CN})_6]^{3-}$
  - (c)  $[\text{Fe}(\text{CN})_6]^{4-}$
  - (d)  $[\text{Fe}(\text{CN})_6]^{3-}$
10. Which of the following options are correct for  $[\text{Fe}(\text{CN})_6]^{3-}$  complex?
  - (a)  $\text{sp}^3$  hybridisation
  - (b)  $\text{sp}^3 \text{d}^2$  hybridisation
  - (c) paramagnetic
  - (d) diamagnetic
11. The denticity of 'PPh<sub>3</sub>' and 'en' are respectively-
  - (a) 1,1
  - (b) 2,1
  - (c) 3,2
  - (d) 1,2
12. The type of isomerism exhibited by complex  $[\text{Co}(\text{NH}_3)_4\text{Br}_2]\text{Cl}$ ?
  - (a) Geometrical and Ionization
  - (b) Geometrical and Optical
  - (c) Optical and Ionization
  - (d) Geometrical only
13. Complex which will be colourless is -
  - (a)  $[\text{Ti}(\text{H}_2\text{O})_6]^{3+}$
  - (b)  $[\text{Ti}(\text{NO}_3)_4]$
  - (c)  $[\text{Cr}(\text{NH}_3)_6]^{3+}$
  - (d)  $[\text{Fe}(\text{CN})_6]^{4-}$
14.  $\text{K}_3[\text{CoF}_6]$  is a high spin complex. The hybridisation of Co in complex will be-
  - (a)  $\text{sp}^3 \text{d}$
  - (b)  $\text{sp}^3 \text{d}^2$
  - (c)  $\text{d}^2 \text{sp}^3$
  - (d)  $\text{dsp}^2$
15. Complex which is paramagnetic as well as tetrahedral is-
  - (a)  $[\text{Ni}(\text{CO})_4]$
  - (b)  $[\text{NiCl}_4]^{2-}$
  - (c)  $[\text{Ni}(\text{CN})_4]^{2-}$
  - (d) None of these
16. On dissolving  $\text{K}_4[\text{Fe}(\text{CN})_6]$  in water, presence of ..... ions are detected.
  - (a)  $\text{K}^+$
  - (b)  $\text{Fe}^{2+}$
  - (c)  $\text{CN}^-$
  - (d) Both  $\text{K}^+$  and  $\text{CN}^-$

17. The complex ion, which does not have any 'd' electron in central metal atom is-
- (a)  $[\text{MnO}_4]^-$  (b)  $[\text{Co}(\text{NH}_3)_6]^{3+}$   
 (c)  $[\text{Fe}(\text{CN})_6]^{3-}$  (d)  $[\text{Cr}(\text{H}_2\text{O})_6]^{3+}$
18. Two isomers of a complex are **A** and **B**. if **A** gives white precipitate on reacting with  $\text{BaCl}_2$ . While **B** gives precipitate of  $\text{AgCl}$  on reaction with  $\text{AgNO}_3$ . The complex **B** and type of isomerism are.
- (a)  $[\text{Co}(\text{NH}_3)_5\text{Cl}]\text{SO}_4$ , Ionization (b)  $[\text{Co}(\text{NH}_3)_5\text{Cl}]\text{SO}_4$ , coordination  
 (c)  $[\text{Co}(\text{NH}_3)_5\text{SO}_4]\text{Cl}$ , Ionization (d)  $[\text{Co}(\text{NH}_3)_5\text{SO}_4]\text{Cl}$ , Coordination
19. The complex which shows optical isomerism is-
- (a)  $\text{trans-}[\text{Co}(\text{en})_2\text{Cl}_2]\text{Cl}$  (b)  $[\text{Co}(\text{NH}_3)_4\text{Cl}_2]\text{Cl}$   
 (c)  $[\text{Co}(\text{NH}_3)_3\text{Cl}_3]$  (d)  $\text{cis-}[\text{Co}(\text{en})_2\text{Cl}_2]\text{Cl}$

## II FILL IN THE BLANKS

- The IUPAC name of linkage isomer of  $[\text{Co}(\text{NH}_3)_5\text{NO}_2]\text{Cl}_2$  is.....
- The oxidation state of Ni in  $\text{K}_2[\text{Ni}(\text{CN})_4]$  is.....
- The metal present in vitamin  $\text{B}_{12}$  is.....
- $\text{NO}$  is named as.....
- Octahedral crystal field splitting the three d orbitals having lower energy are collectively called.....
- The number of isomers of  $[\text{Pt}(\text{NH}_3)_2\text{Cl}_2]$  are.....
- In complex  $[\text{Fe}(\text{H}_2\text{O})_6]^{3+}$  number of unpaired electrons are.....
- Number of ions produced per mole of the complex  $[\text{Co}(\text{NH}_3)_5\text{Cl}]\text{Cl}_2$ ..... are
- Out of cis and trans isomer of  $[\text{RhCl}(\text{en})_2]^+$ , .....isomer shows optical activity.
- Triethylenetetraammine is.....ligand.

## III ASSERTION REASON TYPE QUESTIONS

Note: In the following questions a statement of assertion followed by a statement of reason is given. Choose the correct answer out of the following choice.

- (a) Both assertion and reason are True, and reason is the correct explanation of the assertion.
- (b) Both assertion and reason are True, but reason is not the correct explanation of the assertion.

(c) Assertion is True, but reason is False.

(d) Both assertion and reason are False.

1. **ASSERTION:** Toxic metal ions are removed by the chelating ligands.  
**REASON:** Chelate complex tend to be more stable.
2. **ASSERTION:**  $[(\text{Fe}(\text{CN})_6)]^{3-}$  ion shows magnetic moment corresponding to two unpaired electrons.  
**REASON:** Because it has  $d^2sp^3$  type hybridisation.
3. **ASSERTION:** Carbon monoxide forms low spin complexes with metals.  
**REASON:** Carbon monoxide is neutral oxide.
4. **ASSERTION:** Carbon monoxide is a deadly poison.  
**REASON:** CO can form strong complexes with haemoglobin.
5. **ASSERTION :** Tetrahedral complexes cannot exhibit geometrical isomerism.  
**REASON:** Tetrahedral complexes are chiral in nature.
6. **ASSERTION:** An octahedral complex with two bidentate ligands is always optically active.  
**REASON:** The cis - isomer of the octahedral complex is chiral.
7. **ASSERTION:** Ionisation isomerism is shown by complexes having ambidentate ligands.  
**REASON:** Ambidentate ligands have two possible ligating centres.
8. **ASSERTION:**  $[\text{Fe}(\text{CN})_6]^{3-}$  is more paramagnetic than  $[\text{FeF}_6]^{3-}$ .  
**REASON:**  $[\text{FeF}_6]^{3-}$  has more number of unpaired electrons.
9. **ASSERTION:**  $[\text{Co}(\text{NH}_3)_6]^{3+}$  is an inner orbital complex.  
**REASON:**  $[\text{Co}(\text{NH}_3)_6]^{3+}$  forms an octahedral complex.
10. **ASSERTION:**  $[\text{Fe}(\text{ox})_3]^{3-}$  is more stable than  $[\text{Fe}(\text{H}_2\text{O})_6]^{3+}$   
**REASON:**  $[\text{Fe}(\text{ox})_3]^{3-}$  is a chelate complex.

#### IV ONE WORD TYPE QUESTIONS

1. What is the oxidation state of Ni in  $[\text{Ni}(\text{CO})_4]$ ?
2. Write IUPAC name of  $[\text{Ni}(\text{CN})_4]^{2-}$ .
3. What is the hybridisation of Co in the complex  $[\text{CoF}_6]^{3-}$ ?
4. Write the chemical formula of potassiumtrioxalatochromate(III).
5. Name the transition element present in haemoglobin.



6. Give an example of hexadentate ligand.
7. Out of  $\text{NH}_3$  and  $\text{H}_2\text{O}$  which is strong field ligand?
8. Write electronic configuration of complex  $[\text{Fe}(\text{CN})_6]^{4-}$ .
9. What is the coordination number of Cr in  $[\text{Cr}(\text{en})_2\text{Cl}_2]\text{Cl}$ ?
10. How many geometrical isomers are possible for  $[\text{Cr}(\text{ox})_3]^{3-}$ ?
11. Give the linkage isomer of  $[\text{Pd}(\text{PPh}_3)_2(\text{NCS})_2]$ ?
12. How many unpaired electrons are present in  $[\text{Fe}(\text{NH}_3)_6]\text{Cl}_3$ ?
13. Name the imperfect complex compound which ionizes completely in solutions.
14. Give an example of hexadentate ligand.
15. Write the coordination number of 'Fe' in  $\text{K}[\text{Fe}(\text{CN})_2(\text{en})_2]$

**VERY SHORT ANSWER TYPE QUESTIONS (1 Mark)**

- Q.1. What is ambidentate ligand? Give one example.  
 Ans. Monodentate ligands contain more than one coordinating atoms. Example,  $\text{CN}^-$ .
- Q.2. Write the IUPAC name of  $[\text{PtCl}_2(\text{en})_2(\text{NO}_3)_2]$ .  
 Ans. Dichloridobis(ethane-1, 2-diamine) nitratoplatinum (IV)
- Q.3. What is a chelate ligand? Give one example.  
 Ans. The complex in which ligand binds through two or more donor sites simultaneously to form cyclic complexes. Example: ethane-1-2 diamine.
- Q.4. How many geometrical isomers are possible for the  $[\text{Ni}(\text{NH}_3)_4]^{2+}$ ?  
 Ans. Not possible, because all 4 ligands are same.
- Q.5. Define coordination polyhedron.  
 Ans. The spatial arrangement of the ligand with the central metal ion.
- Q.6. Give the chemical formula of potassium hexacyano ferrate (II).  
 Ans.  $\text{K}_4[\text{Fe}(\text{CN})_6]$ .
- Q.7. Name the metal present in : (i) Chlorophyll (ii) cis platin  
 Ans. (i) Mg (ii) Pt
- Q.8. Which of the two is more stable —  $\text{K}_4[\text{Fe}(\text{CN})_6]$  Or  $\text{K}_3[\text{Fe}(\text{CN})_6]$ ?  
 Ans.  $\text{K}_4[\text{Fe}(\text{CN})_6]$  because Fe has  $d^6$  configuration in this case.
- Q.9. Arrange the following complexes in order of increasing electrical conductivity :  
 $[\text{Co}(\text{NH}_3)_3\text{Cl}_3]$ ,  $[\text{Co}(\text{NH}_3)_5\text{Cl}]\text{Cl}_2$ ,  $[\text{Co}(\text{NH}_3)_4]\text{Cl}_3$   
 Ans.  $[\text{Co}(\text{NH}_3)_3\text{Cl}_3] < [\text{Co}(\text{NH}_3)_5\text{Cl}]\text{Cl}_2 < [\text{Co}(\text{NH}_3)_4]\text{Cl}_3$   
 More number of ions, more electrical conductivity

Q.10. Give an example of the role of coordination compounds in biological system.

Ans. Haemoglobin, the red pigment of blood acts as oxygen carrier. It is a coordination compound of iron.

Q.11. Why is CO a stronger ligand than  $\text{Cl}^-$ ?

Ans. CO is a stronger ligand than  $\text{Cl}^-$  due to back bonding between its empty p-orbitals and filled d-orbitals of central metal ion which in turn increases  $\Delta_o$  value.

Q.12. What do you understand by denticity of a ligand?

Ans. Denticity of a ligand is the number of coordinating or ligating groups sites present in a ligand.

Q.13. What is the coordination number of central metal ion in  $[\text{Fe}(\text{C}_2\text{O}_4)_3]^{3-}$ ?

Ans. Coordination number = Number of ligands  $\times$  Denticity =  $3 \times 2 = 6$

Q.14 Name two properties of the central metal atom/ion which enable it to form stable complex entities.

Ans. (a) Small size of metal ion  
(b) High charge on the metal ion.

Q.15 Write the IUPAC name of the complex  $[\text{Cr}(\text{en})_3][\text{Co}(\text{CN})_6]$ .

Ans. Tris (ethane- 1, 2-diamine) chromium(III) hexacyanidocobaltate(III)

Q.16. A blue coloured solution of  $[\text{CoCl}_4]^{2-}$  ion changes to pink on reaction with  $\text{HgCl}_2$ . Why?

Ans.  $\text{CoCl}_4^{2-} + \text{HgCl}_2 \rightarrow \text{Co}[\text{HgCl}_4] + 2\text{Cl}^-$

[Blue]

[Pink]

**SHORT ANSWER TYPE QUESTIONS (2 or 3 Marks)**

Q.1. Explain the following :

(i)  $\text{NH}_3$  act as a ligand but  $\text{NH}_4^+$  does not.

(ii)  $\text{CN}^-$  is a ambidentate ligand.

Ans. (i)  $\text{NH}_3$  has one lone pair while  $\text{NH}_4^+$  does not.


(ii) Because it has two donor atoms in a monodentate ligand.

Q. 2. Mention the main postulates of Werner theory.

Ans. (i) Metal ion has two types of valency. (ii) Primary valency and secondary valency.  
(iii) Secondary valency is equal to coordination number.

Q. 3. Draw the structure of : (i)  $[\text{Ni}(\text{CO})_4]$  (ii)  $[\text{Fe}(\text{H}_2\text{O})_6]^{3+}$

Ans.



Q.4. How does EDTA help as a cure for lead poisoning?

Ans. Calcium in Ca-EDTA complex is replaced by lead in the body. The more soluble complex Pb-EDTA is eliminated in urine.

Q.5. Define homoleptic and heteroleptic complexes.

Ans. Homoleptic : When metal atom/ion is linked with one type of ligands. Example,  $[\text{Ni}(\text{CO})_4]$ .

Heteroleptic : With the more than one kind of ligands. Example,  $[\text{Co}(\text{NH}_3)_4\text{Cl}_2]^+$

Q.6.  $[\text{NiCl}_4]^{2-}$  is paramagnetic while  $[\text{Ni}(\text{CO})_4]$  is diamagnetic though both are tetrahedral. Why?

Ans. In  $[\text{NiCl}_4]^{2-}$ , Ni has  $3d^8 4s^0$  configuration,  $\text{Cl}^-$  can not pair up while in  $[\text{Ni}(\text{CO})_4]$ , Ni has  $3d^8 4s^2$  configuration, CO pair up electrons

Q.7. The oxidation number of cobalt in the complex : (i)  $\text{K}[\text{Co}(\text{CO})_4]$  (ii)  $[\text{Co}(\text{C}_2\text{O}_4)_3]^{3-}$

Ans. (i) - 1 (ii) + 3

Q.8. What are  $t_{2g}$  and  $e_g$  orbitals?

Ans. In a free transition metal ion, the d-orbitals are degenerate. When it form complex, the degeneracy is split and d-orbitals split into  $t_{2g}$  and  $e_g$  orbitals.

Q.9. What is the solution in which photographic film is washed? What reaction takes place?

Ans. Hypo solution.  $\text{AgBr} + 2\text{Na}_2\text{S}_2\text{O}_3 \rightarrow \text{Na}_3[\text{Ag}(\text{S}_2\text{O}_3)_2] + \text{NaBr}$

Q.10. What is spectrochemical series?

Ans. The arrangement of ligands in order of their increasing crystal field splitting field strength.

Q.12. What are the assumptions of Crystal Field theory.

Ans. (i) Ligand act as a point charge. (ii) Metal ion has electrostatic attraction force with the ligand. (Or any other)

Q.11.  $\text{CuSO}_4$  is colourless while  $\text{CuSO}_4 \cdot 5\text{H}_2\text{O}$  is coloured. Why ?

Ans.  $\text{CuSO}_4$  does not has any ligand, so splitting of d-orbital take place while  $\text{CuSO}_4 \cdot 5\text{H}_2\text{O}$  has water ligand.

Q.12. Differentiate between inner and outer orbital complexes.

Ans. Inner sphere complex : When d-orbital of inner shell take part in hybridisation.

Outer sphere complex : When d-orbital of outermost shell take part in hybridisation.

Q.13. How is stability of coordination compound determined in aqueous solution ?

Ans. By using stability constant. More stability constant, more stability.

Q.14. In a complex ion  $[\text{Co}(\text{NH}_3)_5\text{NO}_2]\text{Cl}_2$ ,

(i) Identify the ligand. (ii) Oxidation number of metal ion.

Ans. (i)  $\text{NH}_3$ ,  $\text{NO}_2^-$ . (ii) +3


Q.15. Explain how the nature of ligand affects the stability of complex ion.

Ans. Strong ligand : More stability Weak ligand : Less stability

### LONG ANSWER TYPE QUESTIONS (5 Marks)

Q.1. A metal ion  $\text{M}^{n+}$  having  $d^4$  valence electronic configuration combines with three didentate ligands to form a complex compound. Assuming  $\Delta_o > P$  : (i) Draw the diagram showing d-orbital splitting during this complex formation. (ii) What type of hybridisation will  $\text{M}^{n+}$  have ? (iii) Name the type of isomerism exhibited by this complex. (iv) Write the electronic configuration of metal  $\text{M}^{n+}$ .

Ans. (i) If  $\Delta_o > P$  then



(ii)  $d^2 sp^3$


(iii)  $[\text{M}(\text{AA})_3]$  type complex show optical isomerism.

(iv)  $t_{2g}^4 e_g^0$

- Q.2. (i) Discuss the nature of bonding in metal carbonyls.  
 (ii) Draw figure to show the splitting of d-orbitals in an octahedral crystal field and write electronic configuration of  $M^{2+}$  ion when : (a)  $P > \Delta_o$  (b)  $\Delta_o > P$

Ans. (i) The metal carbon bond in metal carbonyls possess both  $\sigma$  and  $\pi$  character. The M-C  $\sigma$  bond is formed by the  $M \leftarrow C \equiv O$  while M-C  $\pi$  bond is formed by the donation of a pair of electron from filled d-orbital of metal to antibonding  $\pi^*$  orbital of CO.

(ii)



(a)  $t_{2g}^3 e_g^2$                       (b)  $t_{2g}^5 e_g^0$

- Q.3. (i)  $[Fe(CN)_6]^{4-}$  and  $[Fe(H_2O)_6]^{2+}$  are of different colours in dilute solution. Why?  
 (ii) A complex is prepared by mixing  $CoCl_3$  and  $NH_3$  in the molar ratio of 1 : 4. A 0.1M solution of this complex was found to freeze at  $-0.372^\circ C$ . What is the formula of the complex? ( $K_f = 1.86^\circ C/M$ )

Ans. (i) In both the cases Fe is in +2 state, it has 4 unpaired electrons but  $CN^-$  and  $H_2O$  has different crystal field splitting energy.

(ii)  $\Delta T_f = i \cdot K_f \cdot m$ ,  $i = 2$  means complex dissociate into two ions. Hence the formula is  $[Co(NH_3)_4 Cl_2]Cl$ .

- Q.4.  $CoSO_4 \cdot 5NH_3$  exists in two isomeric forms 'A' and 'B'. Isomer 'A' reacts with  $AgNO_3$  to give white precipitate, but does not react with  $BaCl_2$ . Isomer 'B' gives white precipitate with  $BaCl_2$  but does not react with  $AgNO_3$ .

Answer the following questions: (i) Identify 'A' and 'B' and write their structural formulae. (ii) Name the type of isomerism involved. (iii) Give the IUPAC name of 'A' and 'B'.

Ans. (i)  $A = [Co(NH_3)_5 SO_4]Cl$ ,  $B = [Co(NH_3)_5 Cl]SO_4$

- (ii) Ionisation isomerism
- (iii) (A) = Pentaamminesulphatocobalt(III) chloride
- (B) = Pentaamminechloridocobalt(III) sulphate

Q.5. For the complexes  $[\text{Fe}(\text{CN})_6]^{4-}$  and  $[\text{FeF}_6]^{3-}$  write -

- (i) Oxidation state of Fe
- (ii) IUPAC names
- (iii) Magnetic behaviour
- (iv) Spin type
- (v) Which complex has higher  $\lambda_{\text{max}}$  of absorption?

Ans.

S.No.	$[\text{Fe}(\text{CN})_6]^{4-}$	$[\text{FeF}_6]^{3-}$
(i)	+2	+3
(ii)	Hexacyanatoferate (II)	Hexafluoridoferate (III)
(iii)	Diamagnetic	Paramagnetic
(iv)	Low spin	High spin
(V)	low $\lambda_{\text{max}}$	Higher $\lambda_{\text{max}}$

## CASE STUDY BASED QUESTIONS

## 1. Read the passage and answer the following question.

According to the VBT, the metal atom or ion under the influence of ligands can use its  $(n-1)d$ ,  $ns$ ,  $np$  or  $ns$ ,  $np$ ,  $nd$  orbitals for hybridisation to yield a set of equivalent orbitals of definite geometry such as octahedral, tetrahedral, square planar and so on. These hybridised orbitals are allowed to overlap with ligand orbitals that can donate electron pairs for bonding. The geometry of a complex from the knowledge of its magnetic behaviour on the basis of the valence bond theory In the diamagnetic octahedral complex,  $[\text{Co}(\text{NH}_3)_6]^{3+}$ , the cobalt ion is in +3 oxidation state and has the electronic configuration  $3d^6$ . Six pairs of electrons, one from each  $\text{NH}_3$  molecule, occupy the six hybrid orbitals. Thus, the complex has octahedral geometry and is diamagnetic because of the absence of unpaired electron. In the formation of this complex, since the inner  $d$  orbital ( $3d$ ) is used in hybridisation, the complex,  $[\text{Co}(\text{NH}_3)_6]^{3+}$  is called an inner orbital or low spin or spin paired complex. The paramagnetic octahedral complex,  $[\text{CoF}_6]^{3-}$  uses outer orbital ( $4d$ ) in hybridisation ( $sp^3d^2$ ). It is thus called outer orbital or high spin or spin free complex.

**Source : NCERT**

(A) Number of unpaired electrons present in  $[\text{CoF}_6]^{3-}$  are :

- |       |       |
|-------|-------|
| (a) 4 | (b) 3 |
| (c) 2 | (d) 1 |

(B) Paramagnetic complex is :

- |                                     |                                     |
|-------------------------------------|-------------------------------------|
| (a) $[\text{Ni}(\text{CN})_4]^{2-}$ | (b) $[\text{Ni}(\text{Cl})_4]^{2-}$ |
| (c) $[\text{Ni}(\text{CO})_4]$      | (d) All of these                    |

(C) Outer orbital complex is :

- |                                     |                                       |
|-------------------------------------|---------------------------------------|
| (a) $[\text{CoF}_6]^{3-}$           | (b) $[\text{Co}(\text{NH}_3)_6]^{3+}$ |
| (c) $[\text{Ni}(\text{CN})_4]^{2-}$ | (d) None of these                     |

(D) Inner orbital complex involves :

- (a) nd orbitals                      (b) (n-1)d orbitals
- (c) ns orbitals                      (d) None of these

**2. Read the passage and answer the following question.**

The degeneracy of the d orbitals has been removed due to ligand electron-metal electron repulsions in the octahedral complex to yield three orbitals of lower energy,  $t_{2g}$  set and two orbitals of higher energy,  $e_g$  set. This splitting of the degenerate levels due to the presence of ligands in a definite geometry is termed as crystal field splitting and the energy separation is denoted by  $\Delta_o$  (the subscript o is for octahedral). Thus, the energy of the two  $e_g$  orbitals will increase by  $(3/5) \Delta_o$  and that of the three  $t_{2g}$  will decrease by  $(2/5) \Delta_o$ . The crystal field splitting,  $\Delta_o$ , depends upon the field produced by the ligand and charge on the metal ion. Some ligands are able to produce strong fields in which case, the splitting will be large whereas others produce weak fields and consequently result in small splitting of d orbitals.

(A) The cause of removal of degeneracy of the d orbitals in coordination compounds is :

- (a) Due to higher nuclear charge
- (b) Due to ligand electron-metal electron repulsions
- (c) Atomic size
- (d) None of these

(B) The crystal field splitting is :

- (a) The splitting of the degenerate levels due to the presence of ligands in a definite geometry.
- (b) splitting of f-orbitals
- (c) splitting of p-orbitals
- (d) all of these



(C) Energy of two  $e_g$  orbitals in octahedral complexes increased by :

- (a)  $3/5 \Delta_o$  (b)  $2/5 \Delta_o$   
(c)  $1/5 \Delta_o$  (d)  $5/3 \Delta_o$

(D) Correct relation is :

- (a)  $\Delta_t = 9/4 \Delta_o$  (a)  $\Delta_t = 4/9 \Delta_o$   
(c)  $\Delta_t = 2 \Delta_o$  (d)  $\Delta_t = 5 \Delta_o$

### 3. Read the passage and answer the following question.

The ions or molecules bound to the central atom/ion in the coordination entity are called ligands. These may be simple ions such as  $\text{Cl}^-$ , small molecules such as  $\text{H}_2\text{O}$  or  $\text{NH}_3$ , larger molecules such as  $\text{H}_2\text{NCH}_2\text{CH}_2\text{NH}_2$  or  $\text{N}(\text{CH}_2\text{CH}_2\text{NH}_2)_3$  or even macromolecules, such as proteins. When a ligand is bound to a metal ion through a single donor atom, as with  $\text{Cl}^-$ ,  $\text{H}_2\text{O}$  or  $\text{NH}_3$ , the ligand is said to be unidentate. When a ligand can bind through two donor atoms as in  $\text{H}_2\text{NCH}_2\text{CH}_2\text{NH}_2$  (ethane-1,2-diamine) or  $\text{C}_2\text{O}_4^{2-}$  (oxalate), the ligand is said to be didentate and when several donor atoms are present in a single ligand as in  $\text{N}(\text{CH}_2\text{CH}_2\text{NH}_2)_3$ , the ligand is said to be polydentate.

Ethylenediaminetetraacetate ion ( $\text{EDTA}^{4-}$ ) is an important hexadentate ligand. It can bind through two nitrogen and four oxygen atoms to a central metal ion. When a di- or polydentate ligand uses its two or more donor atoms to bind a single metal ion, it is said to be a chelate ligand. The number of such ligating groups is called the denticity of the ligand. Such complexes, called chelate complexes tend to be more stable than similar complexes containing unidentate ligands. Ligand which can ligate through two different atoms is called ambidentate ligand. Examples of such ligands are the  $\text{NO}_2^-$  and  $\text{SCN}^-$  ions.  $\text{NO}_2^-$  ion can coordinate either through nitrogen or through oxygen to a central metal atom/ion. Similarly,  $\text{SCN}^-$  ion can coordinate through the sulphur or nitrogen atom.

(A) Ambidentate ligand is :

- (a)  $\text{CN}^-$  (b)  $\text{CO}$   
(c)  $\text{NH}_3$  (d)  $\text{H}_2\text{O}$

(B) Total number of donor sites in  $\text{EDTA}^{4-}$  are :

- (a) 6 (b) 4  
(c) 5 (d) 3

(C) Ligand with maximum denticity is-

- (a)  $\text{NC}^-$  (b)  $\text{C}_2\text{O}_4^{2-}$   
(c)  $\text{NH}_3$  (d)  $\text{H}_2\text{O}$

(D) Chelating ligand is :

- (a) EDTA (b)  $\text{NH}_3$   
(c)  $\text{CN}^-$  (d)  $\text{H}_2\text{O}$

## ANSWERS

### I MULTIPLE CHOICE QUESTIONS

1. (b) 2. (c) 3. (b) 4. (a) 5. (c) 6. (a) 7. (b) 8. (c) 9. (a) 10. (c)

### II FILL IN THE BLANKS

1. pentaamminenitrito-O-cobalt(III) chloride 2. 2 3. Cobalt  
4. Nitrosyl 5.  $t_{2g}$  6. 4 7. 5 8. 3  
9. cis 10. tetradenate 11. (d) 12. (a) 13. (b)  
14. (b) 15. (b) 16. (a) 17. (a) 18. (c)  
19. (d)

### III ASSERTION REASON TYPE QUESTIONS

1. (a) 2. (b) 3. (b) 4. (a) 5. (c) 6. (d) 7. (d) 8. (d) 9. (b) 10. (a)

### IV ONE WORD QUESTION ANSWERS

1. 0 2. tetracyanonickelate(II) ion  
3.  $(\text{CoF}_6)^{3-}$ ,  $(sp^3d^2)$  4.  $(\text{K}_3[\text{Cr}(\text{C}_2\text{O}_4)_3])$   
5. Iron 6. EDTA  
7.  $\text{NH}_3$  8.  $t_{2g}^6, e_g^0$   
9. 6 10. 0  
11.  $[\text{Pd}(\text{PPh}_3)_2(\text{NCS})_2]$  12. 1  
13. Double salt 14.  $[\text{EDTA}]^{4-}$   
15. 6

**CASE STUDY BASED QUESTIONS**

**PASSAGE I**

- |       |       |
|-------|-------|
| (A) a | (B) b |
| (C) a | (D) b |

**PASSAGE II**

- |       |       |
|-------|-------|
| (A) b | (B) a |
| (C) a | (D) b |

**PASSAGE III**

- |       |       |
|-------|-------|
| (A) a | (B) a |
| (C) b | (D) a |

## UNIT TEST-1

## Coordination Compounds

Maximum Marks: 20

Time Allowed : 1 hour

1. Explain-  $[\text{Cr}(\text{en})_3]^{2+}$  is more stable than  $[\text{Cr}(\text{H}_2\text{O})_6]^{2+}$ . 1
2. Give two examples of didentate ligands. 1
3. A metal ion  $\text{M}^{n+}$  having  $d^4$  valence electronic configuration combines with three identate ligands to form a complex compound. Assuming  $\Delta_o > P$ , write the electronic configuration of the valence electrons of the metal  $\text{M}^{n+}$  ion in terms of  $t_{2g}$  and  $e_g$ . 1
4. Predict the number of unpaired electrons in the square planar  $[\text{Pt}(\text{CN})_4]^{2-}$  ion. 1
5. For the complex  $[\text{CoBr}_2(\text{en})_2]^{2+}$ , write coordination number and oxidation state of Co. 1
6. Name the isomerism exhibited by following molecules: 2
  - a)  $[\text{Co}(\text{NH}_3)_5\text{SO}_4]\text{Br}$
  - b)  $[\text{Fe}(\text{SCN})_6]^{3-}$
  - c)  $[\text{Co}(\text{en})_3]^{2+}$
  - d)  $[\text{Cr}(\text{en})_2(\text{H}_2\text{O})_2]^{2+}$
7. Define following terms with suitable example. 2
  - a) Spectrochemical Series
  - b) Synergic Bonding
8. Write IUPAC names of following complexes: 2
  - a)  $[\text{Fe}(\text{NO}_2)_2(\text{C}_2\text{O}_4)_2]^{2-}$
  - b)  $[\text{Cr}(\text{en})_2(\text{H}_2\text{O})_2]^{2+}$
9. Explain hybridization, geometry, and magnetic behaviour of following Complexes (Attempt any 2) 2
  - a)  $[\text{Ni}(\text{CO})_4]$
  - b)  $[\text{Mn}(\text{CN})_6]^{3-}$
  - c)  $[\text{FeCl}_6]^{3-}$
10. Arrange the following in increasing order of given properties: 3
  - a)  $[\text{Co}(\text{NH}_3)_3\text{Cl}_3]$ ,  $[\text{Co}(\text{NH}_3)_5\text{Cl}]\text{Cl}_2$ ,  $[\text{Co}(\text{NH}_3)_6]\text{Cl}_3$  (Electrical Conductivity)
  - b)  $[\text{Co}(\text{NH}_3)_6]^{3+}$ ,  $[\text{Co}(\text{CN})_6]^{3-}$ ,  $[\text{Co}(\text{H}_2\text{O})_6]^{3+}$  (Wavelength of absorption)
  - c)  $[\text{Co}(\text{NH}_3)_6]^{3+}$ ,  $[\text{Mn}(\text{CN})_6]^{3-}$ ,  $[\text{Fe}(\text{CN})_6]^{3-}$  (spin-only magnetic moment)
11. Explain following: 3
  - a)  $[\text{Ni}(\text{H}_2\text{O})_6]^{2+}$  is green in colour while  $[\text{Ni}(\text{CN})_6]^{2-}$  is colourless.
  - b)  $[\text{Fe}(\text{H}_2\text{O})_6]^{3+}$  is highly paramagnetic while  $[\text{Fe}(\text{CN})_6]^{3-}$  is weakly paramagnetic.
  - c)  $[\text{NiCl}_4]^{2-}$  is paramagnetic and  $[\text{Ni}(\text{CO})_4]$  is diamagnetic, even though both are tetrahedral

## UNIT TEST-2

## Coordination Compounds

Maximum Marks: 20

Time Allowed : 1 hour

1. Write coordination number and oxidation number of Cr in  $[\text{Cr}(\text{ox})_2\text{Cl}_2]^{3+}$ . 1
2. Give two examples of ambidentate ligands. 1
3. Tetrahedral complexes does not exhibit geometrical isomerism. Explain.
4. Predict the hybridisation of  $[\text{FeCl}_4]^-$  1
5. Mention the denticity of  $\text{C}_2\text{O}_4^{2-}$  &  $\text{EDTA}^{4-}$  of ligands. 1
6. Name the isomerism exhibited by following molecules: 2
  - a)  $[\text{Fe}(\text{CN})_6]^{3-}$
  - b)  $[\text{Co}(\text{NH}_3)_5\text{Cl}]\text{SO}_4$
  - c)  $[\text{Co}(\text{en})_3]^{3+}$
  - d)  $[\text{Co}(\text{NH}_3)_6][\text{Cr}(\text{CN})_6]$
7. Define following terms giving examples: 2
  - a) Ambidentate ligands
  - b) Chelate effect
8. The molar conductivity of the complex  $\text{CrCl}_3 \cdot 4\text{NH}_3 \cdot 2\text{H}_2\text{O}$  is found to be same as that of 3:1 electrolyte. Write its structural formula, name and geometrical isomers of the complex. 2
9. Explain hybridization, geometry, and magnetic, behaviour of following Complexes (Attempt any 2) 3
  - a)  $[\text{Ni}(\text{Cl})_4]^{2-}$
  - b)  $[\text{Co}(\text{CN})_6]^{3-}$
  - c)  $[\text{Cr}(\text{H}_2\text{O})_6]^{3+}$
10.
  - a) Differentiate low spin and high spin complexes giving examples.
  - b) How homoleptic complexes are different from heteroleptic complexes ?
  - c) Write the chemical formula of potassium hexacyanatoferate (II). 3
11. Explain following: 3
  - a) CO is stronger ligand than  $\text{NH}_3$ .
  - b) Low spin octahedral complexes of nickel are not known.
  - c) Aqueous solution of  $[\text{Ti}(\text{H}_2\text{O})_6]^{3+}$  is coloured.


# UNIT 10

## Haloalkanes and Haloarenes

### Points to Remember

#### Classification

- No. of halogen atoms
- $C_2H_5X$   
Monohaloalkane



Monohaloarene




Dihaloarene



Trihaloarene

- Compounds containing  $sp^3$  C-X bond

(a) Alkyl halides



(b) Allylic halides



(c) Benzylic halides



- Compounds containing  $sp^2$  C-X bond

(a) Vinylic halides



(b) Aryl halides



#### Nomenclature

**Common name :** Alkyl group followed by halides.

Dihalogen derivatives of arenes, prefix *o*-, *p*- are used


**IUPAC name :** Numerals are used for position of halogen.

**Nature of C-X bond :**

Carbon-halogen bond is polarised due to more electronegativity of halogen

#### Preparation

**From alcohol:**




**From hydrocarbons:**


(a) By free radical halogenation




(b) By electrophilic substitution





(c) Sandmeyer's reaction



(d) From alkanes



(e) Halogen exchange




### Properties and Reactions of Haloalkanes and Haloarenes

#### Reactions

(a) Nucleophilic substitution




(i) Resonance effect






(ii) Hybridisation of C in C-X bond :

Haloalkane  $\rightarrow sp^3$  Haloarene  $\rightarrow sp^2$



(iii) Phenyl cation unstabilised by resonance



(b) Electrophilic substitution




Friedel - Crafts reaction







(c) Reaction with metals

Wurtz - Fittig reaction



Fittig reaction



**Mechanisms of  $S_N1$  and  $S_N2$ :** **$S_N1$** **Step - 1****Step 2** **$S_N2$** 

	$S_N1$	$S_N2$
Carbon ( $sp^3$ ) Electrophile	Favored by $3^\circ$ alkyl halides or benzylic, or allylic can also react by an $S_N1$ mechanism ( More stable carbocation)	Favored by $-CH_3$ and $1^\circ$ alkyl halides ( less sterically hindered)
Nucleophile	Nature of the nucleophile has no affect on rate. In general, $S_N1$ use neutral, weak nucleophiles	Favored by more reactive nucleophiles $RS^- > NC^- > RO^- > HO^- > Cl^-$
Solvent Effect	Favored by polar, protic solvents.	Favored by polar, aprotic solvents.

**IMPORTANT POINTS**


- In haloarenes electrophilic substitution reaction occurs at o-& p-position.
- Haloarenes exhibit nucleophilic substitution reaction at extremely slow rate due to :
  - (i) Partial double bond character of C-X bond.
  - (ii) Benzene ring is electron rich.
  - (iii) Phenyl carbocation is not stable.
  - (iv) More electronegativity of  $sp^2$  hybridised carbon in haloarene.
  - (v) Steric hindrance due to benzene ring.






## OBJECTIVE TYPE QUESTIONS

## I. MULTIPLE CHOICE QUESTIONS


1. The Number of chiral carbons in given molecule is/are-




- (a) 1 (b) 2  
(c) 3 (d) 4
2. The correct increasing order for rate of reaction towards nucleophilic substitution for following is:



- (a)  $a < b < c$  (b)  $a < c < b$   
(c)  $b < a < c$  (d)  $c < b < a$
3. Which of the following molecules is chiral?
- (a) 2-Bromobutane (b) 1-Bromobutane  
(c) 2-Bromopropane (d) 2-Bromopropan-2-ol
4. Reaction of  $C_6H_5CH_2Br$  with aqueous sodium hydroxide follows.....
- (a)  $S_N1$  mechanism  
(b)  $S_N2$  mechanism  
(c) Either  $S_N1$  or  $S_N2$  mechanism depending on temperature  
(d)  $E_1$  mechanism
5. The correct IUPAC name for following molecule is-




- (a) 1-Bromo-2-ethylpropane  
 (b) 1-Bromo-2-ethyl-2-methylethane  
 (c) 1-Bromo-2-methylbutane  
 (d) 2-Methyl-1-bromobutane
6. The correct increasing order of boiling points of the following compounds is-
- (a) Bromobenzene < 1-Bromobutane < 1-Bromopropane < 1-Bromoethane  
 (b) Bromobenzene < 1-Bromoethane < 1-Bromopropane < 1-Bromobutane  
 (c) 1-Bromopropane < 1-Bromobutane < 1-Bromoethane < Bromobenzene  
 (d) 1-Bromoethane < 1-Bromopropane < 1-Bromobutane < Bromobenzene
7. Alkyl fluorides are synthesised by heating an alkyl chloride/bromide in presence of \_\_\_\_\_ or \_\_\_\_\_.
- (a)  $\text{CaF}_2, \text{CaF}_2$  (b)  $\text{CoF}_2, \text{Hg}_2\text{F}_2$   
 (c)  $\text{Hg}_2\text{F}_2, \text{CaF}_2$  (d)  $\text{NaF}, \text{CaF}_2$
8. Which of the following haloalkanes reacts with aqueous KOH most easily?
- (a) 1-Bromobutane (b) 2-Bromobutane  
 (c) 2-Bromo-2-methylpropane (d) 2-Chlorobutane
9. Toluene reacts with halogen in the presence of  $\text{FeCl}_3$  giving *ortho* and *para* compounds. The reaction is :
- (a) electrophilic elimination (b) electrophilic substitution  
 (c) free radical addition (d) nucleophilic substitution
10. Arrange the following compounds in the increasing order of their densities.



- (a) (A) < (B) < (C) < (D) (b) (A) < (C) < (D) < (B)  
 (c) (D) < (C) < (B) < (A) (d) (B) < (D) < (C) < (A)

11. Chlorobenzene is formed by reaction of chlorine with benzene in presence of  $\text{AlCl}_3$ . Which of the following species attacks the benzene ring in this reaction ?  
 (a)  $\text{Cl}^-$  (b)  $\text{Cl}^+$   
 (c)  $\text{AlCl}_3$  (d)  $\text{AlCl}_4^-$
12. Which of the following statement is correct ?  
 (a) Benzyl halides are more reactive than vinyl and aryl halides.  
 (b) Vinyl halides are more reactive than alkyl halides.  
 (c) Aryl halides are less reactive than alkyl halides.  
 (d) Aryl halides are more reactive than benzyl halides
13. A new carbon-carbon bond is possible in the following reaction:  
 (a)  $\text{C}_6\text{H}_6 + \text{CH}_3\text{Cl} (\text{anhyd. AlCl}_3) \rightarrow$   
 (b)  $\text{CH}_3\text{CH}_2\text{Br} + \text{CH}_3\text{CH}_3\text{NH}_2 \rightarrow$   
 (c)  $\text{CH}_3\text{-Br} + \text{CH}_3\text{CH}_2\text{-ONa} \rightarrow$   
 (d)  $\text{CH}_3\text{CH}_2\text{-Br} + \text{KOH (alc.)} \rightarrow$
14. Alcoholic  $\text{AgNO}_3$  does not give precipitate with-  
 (a)  $\text{C}_6\text{H}_5\text{CH}_2\text{Cl}$  (b)  $\text{CH}_3\text{CH}_2\text{Cl}$   
 (c)  $\text{C}_6\text{H}_5\text{Cl}$  (d)  $\text{CH}_3\text{-CHCl-CH}_3$
15. Reaction intermediate of  $\text{E}_1$  reaction is-  
 (a) Benzyne (b) Carbocation  
 (c) Carbanion (d) Free radical
16. Best reagent for preparation of a chloroalkane from alcohol is -  
 (a)  $\text{SOCl}_2$  (b)  $\text{HCl/ZnCl}_2$   
 (c)  $\text{PCl}_3$  (d)  $\text{Cl}_2/\text{CCl}_4$
17. Highest nucleophilicity is shown by:-  
 (a)  $\text{F}^-$  (b)  $\text{OH}^-$   
 (c)  $\text{CH}_3^-$  (d)  $\text{NH}_2^-$
18. Chloroform on reaction with oxygen form poisonous gas  
 (a)  $\text{CO}_2$  (b)  $\text{COCl}_2$   
 (c)  $\text{Cl}_2$  (d)  $\text{HCl}$
19. The order of reactivity of alkyl halides for  $\text{S}_\text{N}2$  reactions is-  
 (a)  $\text{RF} > \text{RCl} > \text{RBr} > \text{RI}$  (b)  $\text{RF} > \text{RBr} > \text{RCl} > \text{RI}$   
 (c)  $\text{RCl} > \text{RBr} > \text{RF} > \text{RI}$  (d)  $\text{RI} > \text{RBr} > \text{RCl} > \text{RF}$
20. An  $\text{S}_\text{N}2$  reaction at an asymmetric carbon gives-  
 (a) enantiomer of substrate (b) product with opposite optical rotation  
 (c) mixture of diastereomers (d) single stereoisomer
21. The product in the following reaction is -



**II FILL IN THE BLANKS**

1. Chlorobenzene may be converted into diphenyl by.....reaction.
2. The mixture containing two enantiomers in equal amount, having zero optical rotation is called.....
3. Reaction of bromopropane with magnesium metal in dry ether forms the category of compounds called.....
4. The major product of Friedel-Crafts acetylation of chlorobenzene is.....
5. Polyhalogen compound having antiseptic property is.....
6. Alkyl halides are..... in water but.....in organic solvents.
7. Bimolecular nucleophilic substitution of optically active haloalkanes leads to the .....of the configuration.
8. The molecules which rotates the plane-polarized light in clockwise direction are called.....
9. Halogens.....the aromatic ring towards electrophilic substitution due to -I effect of halogens.

**III ASSERTION REASON TYPE QUESTIONS**

In these questions, a statement of assertion followed by a statement of reason is given. Choose the correct answer out of the following choices:

- (A) Assertion and reason both are correct statements and reason is correct explanation for assertion.
  - (B) Assertion and reason both are correct statements but reason is not correct explanation for assertion.
  - (C) Assertion is correct statement but reason is wrong statement.
  - (D) Assertion is wrong statement but reason is correct statement.
1. **ASSERTION :**  $S_N2$  reaction is accompanied by the inversion of configuration.  
**REASON :**  $S_N2$  reaction occurs in two step.
  2. **ASSERTION :** Treatment of chloroethane with saturated solution of AgCN give ethylisocyanide as major product.  
**REASON :** Cyanide ion ( $CN^-$ ) is an ambidentate nucleophile.
  3. **ASSERTION :** The boiling points of alkyl halides decreases in the order :  $RI > RBr > RCl > RF$   
**REASON :** The boiling points of alkyl chlorides, bromides and iodides are considerably higher than that of the hydrocarbon of comparable molecular mass.
  4. **ASSERTION :** tert-Butyl bromide undergoes Wurtz reaction to give 2, 2, 3, 3-tetramethylbutane.  
**REASON :** In Wurtz reaction, alkyl halides react with sodium in dry ether to give hydrocarbon containing double the number of carbon atoms present in the halide.
  5. **ASSERTION :** Presence of a nitro group at *ortho* or *para* position increases the reactivity of haloarenes towards nucleophilic substitution.  
**REASON :** Nitro group, being an electron withdrawing group decreases the electron density over the benzene ring.
  6. **ASSERTION :** In monohaloarenes, further electrophilic substitution occurs at *ortho* and *para* positions.


7. **ASSERTION :** Aryl iodides can be prepared by reaction of arenes with iodine in the presence of an oxidising agent.  
**REASON :** Oxidising agent oxidises  $I_2$  into HI.
8. **ASSERTION:** The nucleophilic substitution of vinyl chloride is difficult than ethyl chloride.  
**REASON:** Vinyl group is electron donating group.
9. **ASSERTION:** Silver nitrite gives nitro alkane when it reacts with an alkyl halide.  
**REASON:** Silver nitrite is an ionic compound.
10. **ASSERTION:** Chloroform is generally stored in brown bottles which are filled to brims.  
**REASON :**  $CHCl_3$  get oxidised to phosgene in atmosphere.
11. **ASSERTION:** Neopentyl chloride undergoes  $S_N2$  reaction easily.  
**REASON:** Neopentyl chloride is a tertiary halide.
12. **ASSERTION :** It is difficult to replace chlorine by -OH in chlorobenzene in comparison to that in chloroethane.  
**REASON :** Chlorine-carbon (C-Cl) bond in chlorobenzene has a partial double bond character due to resonance.
13. **ASSERTION:** 2-Bromobutane reacts with aqueous KOH forming racemic mixture.  
**REASON:** 2-Bromobutane forms secondary carbocation which leads to bimolecular substitution reaction.

#### IV **ONE WORD ANSWER TYPE QUESTIONS**

1. Name the category of nucleophiles to which  $CN^-$ ,  $NO_2^-$  belongs, which may attack through two different sites.
2. Name the reaction which converts aniline into chlorobenzene.
3. Which isomer of dichlorobenzene has highest boiling point?
4. If elimination takes place according to Zaitsev (Saytzeff) rule then from which carbon removal of -H takes place?
5. Which mechanism  $Ph_3C-Cl$  follows when it reacts with aqueous NaOH?
6. Name the instrument used to measure optical rotation.
7. Name the category of the molecules related by non-superimposable mirror images of each other.
8. Which gas is formed by the reaction of chloroform with oxygen in presence of light?
9. Name the first chlorinated organic insecticide used against mosquito during world war II?
10. Write name of major product formed by reaction of ethyl magnesium bromide with water :
11. What will be major product when 2-Bromopentane reacts with alcoholic KOH ?
12. What will be the sign of optical rotation for a laevorotatory molecule?

**VERY SHORT ANSWER TYPE QUESTIONS (1 Mark)**

1. Give IUPAC name of:



Ans : 1-Chloro-2,3-dibromo-2-methylpentane


2. Identify A and B in each of the following process :



Ans : A:  $\text{CH}_3\text{-CH}_2\text{-CN}$ ; B:  $\text{CH}_3\text{CH}_2\text{CH}_2\text{NH}_2$


3. Draw the structure of 4-bromo-3-methylpent-2-ene.

Ans:



4. Why Grignard reagent should be prepared under anhydrous condition?


Ans: It reacts with water and converts into corresponding hydrocarbon.



5. Chloroform is stored in dark coloured and sealed bottles. Why ?

Ans: It is slowly oxidised by air in presence of light to form  $\text{COCl}_2$  (Phosgene) which is a poisonous gas.  $(\text{CHCl}_3 + \frac{1}{2} \text{O}_2 \rightarrow \text{COCl}_2 + 2\text{HCl})$

6. An alkyl halide having molecular formula  $\text{C}_4\text{H}_9\text{Cl}$  is optically active. Write its structure.



7. An organic compound 'A' on treatment with KCN gave 'B' which on hydrolysis with dil. HCl gave acetic acid. Identify A.

Ans:  $\text{CH}_3\text{Cl}$

8. Arrange the following in order of their increasing reactivity in nucleophilic substitution reaction :  $\text{CH}_3\text{F}$ ,  $\text{CH}_3\text{I}$ ,  $\text{CH}_3\text{Br}$ ,  $\text{CH}_3\text{Cl}$

Ans:  $\text{CH}_3\text{F}$ ,  $\text{CH}_3\text{Cl}$ ,  $\text{CH}_3\text{Br}$ ,  $\text{CH}_3\text{I}$

- Ans: Due to more stability of allyl carbocation than n-propyl carbocation.

- ClCC1=CC=CC=C1.[Na]>>[Na]C1=CC=CC=C1

11. Give one chemical test to distinguish between chlorobenzene and benzyl chloride?

Ans: Nitro group withdraws electrons by -R, -I effect resulting the formation of more stable carbocation.

- Ans: The by-products are gaseous  $\text{SO}_2$  and  $\text{HCl}$  which can be easily removed to give pure haloalkane.

- 

Ans: Due to  $sp^2$  hybridised carbon in chlorobenzene which is more electronegative and reduces polarity of C-Cl bond.

16. Name the compound which will be formed by reaction of (+)-2-chlorobutane with KOH?

Ans: (+)-butan-2-ol


17. What happens when methylchloride is treated with KCN?

Ans:  $\text{CH}_3\text{-CN}$  will be formed

18. Identify X and Y :  $\text{R-Br} + \text{Mg} \xrightarrow{\text{Dry Ether}} \text{X} \xrightarrow{\text{D}_2\text{O}} \text{Y}$

Ans. X:  $\text{R-MgBr}$       Y:  $\text{R-D}$


19. Complete by writing structures of P and Q:



20. Name the product formed when 2,4,6- trinitrochlorobenzene reacts with warm water.

Ans. Picric acid(2,4,6-trinitrophenol)

21. Write structure of product formed when following undergoes  $\text{S}_\text{N}2$  reaction taking  $\text{OH}^-$  as nucleophile:



22. Optical rotation of an enantiomer is  $+12.5^\circ$ . Write optical rotation of :

(i) Its mirror image

(ii) mixture of enantiomer & mirror image (1:1)

Ans. (i)  $-12.5^\circ$       (ii) Zero

23. Out of pentan-2-ol and pentan-3-ol, which is chiral molecule?

Ans. Pentan-2-ol



SHORT ANSWER TYPE QUESTIONS (2 or 3 Marks)

1. Why is sulphuric acid not used during the reaction of alcohols with KI?

Ans:  $\text{H}_2\text{SO}_4$  cannot be used along with KI in the conversion of an alcohol to an alkyl iodide as it converts KI to corresponding acid, HI which is then oxidised by it to  $\text{I}_2$ .

2. Which compound in each of the following pairs will react faster in  $\text{S}_{\text{N}}2$  reaction with  $\text{OH}^-$ ?

(i)  $\text{CH}_3\text{Br}$  or  $\text{CH}_3\text{I}$                       (ii)  $(\text{CH}_3)_3\text{CCl}$  or  $\text{CH}_3\text{Cl}$


Ans: (i) With  $\text{OH}^-$  ion,  $\text{CH}_3\text{I}$  will react faster than  $\text{CH}_3\text{Br}$ , as iodide is a better leaving group.

(ii)  $\text{CH}_3\text{Cl}$  as it is primary haloalkanes with less steric hindrance.


3. Carry out the following conversions in not more than two steps :

(i) Toluene to Benzyl alcohol

(ii) Benzyl alcohol to phenylethanenitrile



4. Which of the following compounds would undergo  $\text{S}_{\text{N}}1$  Reaction faster and why?



Ans: **a** will be more reactive due to higher stability of benzyl carbocation.

6. Complete the following reaction :


- (i)  $\text{C}_6\text{H}_5\text{N}_2^+\text{Cl}^- + \text{KI} \rightarrow$   
 (ii)  $\text{CH}_2=\text{CH}_2 + \text{Br}_2 \rightarrow$   
 (iii)  $\text{CH}_3\text{CH}_2\text{OH} + \text{SOCl}_2 \rightarrow$

Ans: (i)  $\text{C}_6\text{H}_5\text{I}$ , (ii)  $\text{CH}_2\text{Br}-\text{CH}_2\text{Br}$ , (iii)  $\text{CH}_3\text{CH}_2\text{Cl}$

7. Convert :

- (i) Benzene to *p*-nitrochlorobenzene  
 (ii) Benzene to diphenyl

Ans:



8. What happen when :

- (i) Propene is treated with HBr in presence of peroxide.  
 (ii) Benzene is treated with methyl chloride in presence of  $\text{AlCl}_3$ .


Ans: (i) 1-Bromopropane is formed (Anti-Markonikov addition)

- (ii) Toluene is formed (Friedel-Crafts Alkylation)

9. Write short note on :

- (i) Sandmeyer reaction                      (ii) Finkelstein reaction

Ans:



10. Name the reagent used to convert:

- (i) 2-Chloropropane to 2-nitropropane  
 (ii) Chloroethane to butane

Ans: (i)  $\text{AgNO}_2$                       (ii)  $\text{Na}$  / dry ether

11. Give reasons :

- (i) Boiling point of alkyl bromide is higher than alkyl chloride.
- (ii) Alkyl halides are better solvents than aryl halides.


Ans: (i) High magnitude of van der Waals forces in alkyl bromides.

- (ii) C - X is more polar in alkyl halides.

12. Carry out the following conversion :

- (i) But-1-ene to n-Butyl iodide
- (ii) 2-Bromopropane to 1-Bromopropane


Ans:



13. Identify and indicate the presence of center of chirality (if any) in the following molecules. How many stereoisomers are possible for those containing chiral center?



- (i) 1,2-Dichloropropane
- (ii) 3-Bromopent-1-ene

Ans:



[\*marked carbon are chiral centres]

14. Complete the reactions:



Ans :



15. Haloalkanes react with KCN to form alkyl cyanides as main product while AgCN form isocyanide as the chief product. Explain

Ans: KCN is predominantly ionic and provides cyanide ions in solution. Although both carbon and nitrogen atoms are in a position to donate electron pairs, the attack takes place mainly through carbon atom and not through nitrogen atom since C-C bond is more stable than C-N bond. However, AgCN is mainly covalent in nature and nitrogen is free to donate electron pair forming isocyanide as the main product.

16. An organic compound **A** react with  $\text{PCl}_5$  to give compound **B**, compound **B** react with Na/ether to give n-butane. What are compounds **A** and **B**?

Ans: **A** =  $\text{C}_2\text{H}_5\text{OH}$ , **B** =  $\text{C}_2\text{H}_5\text{Cl}$

17. The treatment of alkyl chlorides with aqueous KOH leads to the formation of alcohols but in presence of alcoholic KOH, alkenes are major products. Explain.

Ans: In aqueous medium i.e. water, KOH will produce strong nucleophile  $\text{OH}^-$  which will bring about the substitution of alkyl halides to form alcohols. At the same time, the  $\text{OH}^-$  ions will be highly hydrated also. They will not be able to abstract a proton ( $\text{H}^+$ ) from the  $\beta$ -carbon atom to form alkenes.


In alcoholic KOH, the solution will also contain ethoxide ions ( $\text{C}_2\text{H}_5\text{O}^-$ ) in addition to  $\text{OH}^-$  ions. They being a stronger base than  $\text{OH}^-$  ions, will abstract a  $\text{H}^+$  ion from the  $\beta$ -carbon atom giving alkene as the product as a result of dehydrohalogenation.

### LONG ANSWER TYPE QUESTIONS (5 Marks)

1. Haloarenes are extremely less reactive towards nucleophilic substitution reactions. Explain.


Ans: Aryl halides are extremely less reactive towards nucleophilic substitution reactions due to the following reasons:

- (i) Resonance effect : In haloarenes, the electron pairs on halogen atom are in conjugation with  $\pi$ -electrons of the ring and the following resonating structures are possible.



C-Cl bond acquires a partial double bond character due to resonance. As a result, the bond cleavage in haloarene is difficult than haloalkane and therefore, they are less reactive towards nucleophilic substitution reaction.

- (ii) Difference in hybridisation of carbon atom in C-X bond: In haloalkane, the carbon atom attached to halogen is  $sp^3$  hybridised while in case of haloarene, the carbon atom attached to halogen is  $sp^2$ -hybridised.




The  $sp^2$  hybridised carbon with a greater s-character is more electronegative and can hold the electron pair of C-X bond more tightly than  $sp^3$ -hybridised carbon in haloalkane with less s-character. Since it is difficult to break a shorter bond than a longer bond therefore, haloarenes are less reactive than haloalkanes towards nucleophilic substitution reaction.

- (iii) Instability of phenyl cation: In case of haloarenes, the phenyl cation formed as a result of self-ionisation will not be stabilised by resonance and therefore,  $S_N1$  mechanism is ruled out.

- (iv) Because of the possible repulsion, it is less likely for the electron rich nucleophile to approach electron rich arenes.

2. Although chlorine is an electron withdrawing group, yet it is *ortho*- and *para*-directing in electrophilic aromatic substitution reaction. Explain.

Ans: Chlorine withdraws electrons through inductive effect and releases electrons through resonance. Through inductive effect, chlorine destabilises the intermediate carbocation formed during the electrophilic substitution.



Through resonance, halogen tends to stabilise the carbocation and the effect is more pronounced at *ortho*- and *para*- positions. The inductive effect is stronger than resonance and causes net electron withdrawal and thus causes net deactivation. The resonance effect tends to oppose the inductive effect for the attack at *ortho*- and *para*-positions and hence makes the deactivation less for *ortho*- and *para*-attack. Reactivity is thus controlled by the stronger inductive effect and orientation is controlled by resonance effect.

3. A primary alkyl halide (**A**),  $C_4H_9Br$  reacted with hot alcoholic KOH to give compound (**B**). Compound (**B**) reacted with HBr to give (**C**), which is an isomer of (**A**). When (**A**) was reacted with sodium metal, it gave a compound (**D**),  $C_8H_{18}$  which was different than the compound when n-butyl bromide was reacted with sodium. Give the structural formula of (**A**) and write equations of all the reactions.

Ans: (A) 1-Bromo-2-methylpropane

(B) 2-Methylprop-1-ene

(C) tert-butylbromide

(D) 2,5-Dimethylhexane


3. Write wedge and Dash representation of 2-Bromobutane then write structures of molecules when:

Ans: (i) It undergoes retention on reaction with HCl


(ii) It undergoes inversion on reaction with  $OH^-$

(iii) It undergoes  $S_N1$  reaction with  $H_2O$

(iv) It placed in front of mirror and then rotated  $180^\circ$



5. Predict A to E in following sequence of reactions:



Ans. A:  $\text{CH}_3-\text{CH}=\text{CH}_2$     B:  $\text{CH}_3-\text{CH}=\text{CH}_2$     C:  $\text{CH}_3-\text{CH}_2-\text{CH}_2-\text{Br}$   
 D:  $\text{CH}_3-\text{CN}$     E:  $\text{CH}_3-\text{COOH}$

6. Give reasons:


- (i)  $(\pm)$ -Butan-2-ol is optically inactive
- (ii) Iodoethane is more reactive than chloroethane in  $\text{S}_\text{N}2$  reaction
- (iii) *p*-dichlorobenzene has higher m.p. than *o*-& *m*-isomers.
- (iv) Haloalkanes easily dissolve in organic solvents.
- (v) Grignard's reagent is stored in moisture proof containers.

Ans. (i) Rotation due to one enantiomer cancelled out by another in racemic mixture.  
 (ii) Because I is better leaving group than Cl  
 (iii) Due to symmetry of *p*-isomer, it fits in lattice better  
 (iv) New intermolecular attractions between haloalkanes and solvent have much same strength as the ones being broken.  
 (v) Grignard's reagent reacts with water forming corresponding alkane.

## CASE STUDY BASED QUESTIONS

## 1. Read the passage and answer the following questions:

The bimolecular nucleophilic substitution ( $S_N2$ ) reactions are among the fundamental and most important organic reactions. Traditionally, the mechanism of the  $S_N2$  reactions is studied using qualitative transition state theory. The functionalized  $sp^3$  hybridized carbon in a substrate molecule functions as an electrophilic center. This electrophilicity is considered due to a partial positive charge created on carbon by the electronegative functional group. A nucleophile ( $Nuc^-$ ) attacks the  $sp^3$  hybridized carbon from the opposite side of the leaving group (-LG). This nucleophilic attack results in a transition state in which the carbon atom becomes  $sp^2$  hybridized with the C-LG bond partially broken and the Nuc-C bond partially formed. Finally, the C-LG bond is broken completely coincident with formation of the Nuc-C bond, giving the nucleophilic substitution product.



**Reference :** Xiaoping Sun, **Mechanistic Studies of Nucleophilic Substitution and  $\beta$ -Elimination Reactions**, *Symmetry* **2010**, 2(1), 201-212; <https://doi.org/10.3390/sym2010201>

(A) Which of the following undergoes  $S_N2$  reaction fastest?

- |                   |                   |
|-------------------|-------------------|
| (a) Chloropropane | (b) Bromopropane  |
| (c) Iodopropane   | (d) Chlorobenzene |


(B)  $S_N2$  reaction accompanied by.....of the configuration.

- |                  |                 |
|------------------|-----------------|
| (a) Retention    | (b) Inversion   |
| (c) Racemisation | (d) Tautomerism |

(C) Reaction of  $(CH_3)_3C-Cl$  with aq. NaOH gives:

- |                     |                     |
|---------------------|---------------------|
| (a) $(CH_3)_3CH-OH$ | (b) $(CH_3)_2CH-OH$ |
| (c) (a) and (b)     | (d) But-2-ene       |

(D) The correct order of rate of reaction in  $S_N2$  for following molecules are:





- (I)  $1 > 2 > 3$  (ii)  $2 > 1 > 3$   
 (iii)  $1 > 3 > 2$  (iv)  $3 > 2 > 1$

2. **Read the passage and answer the following questions:**


Halogenation of alcohols is a useful reaction as alcohol-starting materials are cheap and large varieties are commercially available. Many procedures are known that convert alcohols to halides however, they can involve harsh conditions and hazardous chemicals. For example, the chlorination of alcohols is traditionally performed using HCl gas or thionyl chloride, both of which are hazardous to human health and often produce many side reactions. Although improvements have been made upon these halogenation reactions, further green methodology is desired. On a laboratory scale, ammonium halide salts ( $\text{NH}_4\text{X}$ ) are solid reagents, which means they are easy to handle and measure.

Discovering a mild, neutral, highly selective and environmentally friendly system for catalytic dehydroxyhalogenation is desirable but it is not a trivial task. Compromises in the journey towards an ideal sustainable process have to be made. This is exhibited by some of the trends found in the literature regarding chlorination of alcohols. Some reports involve the use of tripbenylphosphine ( $\text{PPh}_3$ ) as a superstoichiometric additive in dehydroxychlorination reactions.

**Reference :** Petten, C.F., Kalviri, H.A. & Kelton, F.M. **Halodehydroxylation of alcohols to yield benzylic and alkyl halides in ionic liquids.** *Sustain Chem Process* 3, 16 (2015). <https://doi.org/10.1186/s40508-015-0043-4>


- (A) Which of the following reaction may be used for the formation of 2-Fluorobutane?
- (a) Finkelstein Reaction (b) Swarts Reaction  
 (c) Sandmeyer Reaction (d)  $\text{S}_{\text{N}}2$  Reaction
- (B) For the reaction,  $\text{R-OH} + \text{HX} \rightarrow \text{R-X} + \text{H}_2\text{O}$  the rate of reaction will be highest with the-
- (a) Ethanol (b) Propan-1-ol  
 (c) Propan-2-ol (d) 2-Methylbutan-2-ol
- (C) When primary, secondary and tertiary alcohols reacts with Lucas' reagent ( $\text{HCl} + \text{ZnCl}_2$ ), the incorrect observation is-

- (a) Primary alcohol gives turbidity immediately.  
 (b) Secondary alcohols gives turbidity in 5-10 minutes.  
 (c) Tertiary alcohols are most reactive.  
 (d) Reaction follows  $S_N1$  pathway.  
 (D) The correct product of following reaction is-




3. **Read the passage and answer the following questions:**

Bimolecular nucleophilic substitution ( $S_N2$ ) reactions constitute one of the most widely-used organic chemistry reactions, both in chemistry and biology. The general reaction scheme is summarized in Scheme I, where a nucleophile  $Nu^q$  attacks the central atom A and simultaneously a leaving group LG is displaced. The reaction can proceed for either anionic species (typically  $q_1=q_4<0$ ), neutral (radical) species (typically  $q_1=q_2=q_3+q_4=0$ ), or cationic species (typically  $q_2=q_3>0$ ), together with a wide range of nucleophiles, leaving groups and central atoms. The number and nature of the substituents around the central atom play a major role in determining reactivity.



$S_N2$  substitution is, in principle, always in competition with base induced elimination (E2), and the two pathways may occur as unwanted side reactions of each other.




**Reference :** T.A. Hamlin, M. Swart, F. M. Bickelhaupt, **Nucleophilic Substitution ( $S_N2$ ): Dependence on Nucleophile, Leaving Group, Central Atom, Substituents, and Solvent** *ChemPhysChem* **2018**, 19, 1315.

(A) Which of the following molecule leads to fastest reaction according to  $S_N2$  mechanism.

- (a) 2-Chloropropane                      (b) 1-Chloropropane  
(c) Chlorophenylmethane              (d) 2-Chlorotoluene

(B) In given reactions, X and Y are respectively-



- (a) Pent-1-ene, Pent-2-ene
- (b) Pent-2-ene, Pentan-2-ol
- (c) Pentan-2-ol, Pent-2-ene
- (d) Pent-1-ene, Pentan-2-ol
- (C) Which of the following haloalkane reacts with nucleophile forming racemic mixture?
  - (a) 1-Chloropropane
  - (b) Chlorobenzene
  - (c) 2-Chlorobutane
  - (d) 3-Chlorobutane
- (D) Which of the following statement is incorrect about  $S_N2$  reaction?
  - (a) It takes place in single step.
  - (b) It is accompanied by the inversion of configuration.
  - (c) Stability of carbocation determines the rate of reaction.
  - (d) Iodopropane undergoes  $S_N2$  faster than Chloropropane.

4. **Read the passage and answers the following questions:**

Nucleophilic substitution reaction of haloalkane can be conducted according to both  $S_N1$  and  $S_N2$  mechanisms. However, which mechanism it is based on is related to such factors as the structure of haloalkane, and properties of leaving group, nucleophilic reagent and solvent.

**Influences of halogen :** No matter which mechanism the nucleophilic substitution reaction is based on, the leaving group always leave the central carbon atom with electron pair. This is just the opposite of the situation that nucleophilic reagent attacks the central carbon atom with electron pair. Therefore, the weaker the alkalinity of leaving group is, the more stable the anion formed is and it will be more easier for the leaving group to leave the central carbon atom; that is to say, the reactant is more easier to be substituted. The alkalinity order of halogen ion is  $F < Br < Cl < I$  and the order of their leaving tendency should be  $I > Br > Cl > F$ . Therefore, in four halides with the same alkyl and different halogens, the order of substitution reaction rate is  $RI > RBr > RCl > RF$ . In addition, if the leaving group is very easy to leave, many carbocation intermediates are generated in the reaction and the reaction is based on  $S_N1$  mechanism. If the leaving group is not easy to leave, the reaction is based on  $S_N2$  mechanism.

**Influences of solvent polarity:** In  $S_N1$  reaction, the polarity of the system increases from the reactant to the transition state, because polar solvent has a greater stabilizing effect on the transition state than the reactant, thereby reduce activation energy and accelerate the reaction. In  $S_N2$  reaction, the polarity, of the system generally does not change from the reactant to the transition state and only charge dispersion occurs. At this time, polar solvent has a great stabilizing effect on Nu than the transition state, thereby increasing activation energy and low down the reaction rate. For example, the decomposition rate ( $S_N1$ ) of tertiary chlorobutane in 25°C water (dielectric constant 79) is 300000 times faster than in ethanol (dielectric constant 24). The reaction rate ( $S_N2$ ) of 2-bromopropane and NaOH in ethanol containing 40% water is twice slower than in absolute ethanol. In a word, the level of solvent polarity has influence on both  $S_N1$  and  $S_N2$  reactions, but with different results. Generally speaking, weak polar solvent is favorable for  $S_N2$  reaction, while strong polar solvent is favorable for  $S_N1$  reaction, because only under the action of polar solvent can halogenated hydrocarbon dissociate into carbocation and halogen ion and solvents with a strong polarity is favorable for solvation of carbocation, increasing its stability. Generally speaking, the substitution reaction of tertiary haloalkane is based on  $S_N1$  mechanism in solvents with a strong polarity (for example, ethanol containing water).

**Reference** A Brief Discussion on Nucleophilic Substitution Reaction on Saturated Carbon Atom. In *Applied Mechanics and Materials* (Vol. 312, pp. 433-437). Trans. Tech Publications Ltd.) (Ding, Y. (2013).

(A)  $S_N1$  mechanism is favoured in which of the following solvents:

- (a) benzene
- (b) carbon tetrachloride
- (c) acetic acid
- (d) carbon disulphide

(B) Nucleophilic substitution will be fastest in case of:

- (a) 1-Chloro-2,2-dimethyl propane
- (b) 1-Iodo-2,2-dimethyl propane
- (c) 1-Bromo-2,2-dimethyl propane
- (d) 1-Fluoro-2,2-dimethyl propane

(C)  $S_N1$  reaction will be fastest in which of the following solvents?

- (a) Acetone (dielectric constant 21)
- (b) Ethanol (dielectric constant 24)
- (c) Methanol (dielectric constant 32)
- (d) Chloroform (dielectric constant 5)

- (D) Polar solvents make the reaction faster as they:  
 (a) destabilize transition state and decrease the activation energy  
 (b) destabilize transition state and increase the activation energy  
 (c) stabilize transition state and increase the activation energy  
 (d) stabilize transition state and decrease the activation energy
- (E)  $S_N1$  reaction will be fastest in case of:  
 (a) 1-Chloro-2-methyl propane  
 (b) 1-Iodo-2-methyl propane  
 (c) 1-Chlorobutane  
 (d) 1-Iodobutane

### ANSWERS

#### I MULTIPLE CHOICE QUESTIONS

1. b 2. a 3. a 4. a 5. c 6. d 7. b 8. a 9. b 10. a 11. b 12. a 13. a 14. c 15. b  
 16. a 17. c 18. b 19. d 20. d 21. a

#### II FILL IN THE BLANKS

- |                      |                                  |
|----------------------|----------------------------------|
| 1. Fittig reaction.  | 2. Racemic mixture.              |
| 3. Grignard reagent. | 4. <i>p</i> -chloroacetophenone. |
| 5. Iodoform.         | 6. Insoluble, soluble.           |
| 7. Inversion.        | 8. Dextrorotatory.               |
| 9. Deactivates       |                                  |

#### III ASSERTION REASON TYPE QUESTIONS

- |       |       |       |
|-------|-------|-------|
| 1. C  | 2. B  | 3. B  |
| 4. D  | 5. A  | 6. B  |
| 7. D  | 8. C  | 9. C  |
| 10. A | 11. D | 12. A |
| 13. C |       |       |

#### IV ONE WORD TYPE QUESTIONS

- |                          |                        |
|--------------------------|------------------------|
| 1. Ambident nucleophile. | 2. Sandmeyer reaction. |
| 3. para-dichlorobenzene. | 4. $\beta$ -carbon.    |
| 5. $S_N1$ .              | 6. Polarimeter.        |
| 7. Enantiomers.          | 8. Phosgene            |
| 9. DDT                   | 10. Ethane             |
| 11. Pent-2-ene           | 12. Negative           |

**CASE STUDY BASED QUESTIONS**

1. (A) - (c)      (B) - (b)      (C) - (a)      (D) - (a)
2. (A) - (b)      (B) - (d)      (C) - (a)      (D) - (b)
3. (A) - (a)      (B) - (b)      (C) - (c)      (D) - (c)
4. (A) - (c)      (B) - (b)      (C) - (c)      (D) - (c)      (E) - (b)

## UNIT TEST-1


## HALOALKANES AND HALOARENES

Maximum Marks : 20

Time Allowed : 1 Hr

1. Write IUPAC name of following:

1



2. The correct ascending order of nucleophilic substitution reactions for following molecules is-

1

(I) Chlorobenzene

(II) 4-Nitrochlorobenzene

(III) 4-Chlorotoluene

(a) I,II,III

(b) I,III,II

(c) III,I,II

(d) III,II,I

3. How will you convert aniline into iodobenzene?

1

4. Chlorobenzene on reaction with chloromethane in presence of zinc chloride forms-

1

(a) *o*-dichlorobenzene(b) *o*-chlorotoluene(c) *p*-chlorotoluene(d) *o*-chlorotoluene and *p*-chlorotoluene

5. A solution of a molecule (X) rotates plane polarised light
- $32^\circ$
- clockwise. Which of the following is incorrect?

1

(a) X is optically active


(b) X is dextrorotatory

(c) X has all achiral carbons


(d) X forms non-superimposable

6. Which of the following undergoes
- $S_N2$
- reaction faster and Why?

2



OR







7. How will you synthesize following from aniline? Write chemical equations. 2  
(i) Chlorobenzene (ii) Iodobenzene
8. Explain following giving examples: 2  
(i) racemic mixture (ii) enantiomers
9. What happens when: 3  
(i) Bromoethane reacts with sodium ethoxide  
(ii) Chloroethane reacts with AgCN.  
(iii) 2-Bromo-2-methylbutane is heated with ethanolic KOH.
10. Explain following: 3  
(i) Aryl chlorides and bromides can be easily prepared by electrophilic substitution of aromatic hydrocarbons with chlorine and bromine respectively in the presence of Lewis acid catalysts but for preparation of aryl iodides presence of an oxidising agent is required.  
(ii) Grignard reagent is kept under anhydrous conditions.  
(iii) Reactions through  $S_N2$  mechanisms are accompanied by inversion of configuration.
11. Convert the following: 3  
(i) Ethanol to iodoethane  
(ii) Toluene to benzyl alcohol  
(iii) Benzene to diphenyl.







## UNIT TEST-2

## HALOALKANES AND HALOARENES

Maximum Marks : 20

Time Allowed : 1 Hr

- Write the IUPAC name of  1
- 

Identify A & B 1
- Define ambident nucleophile with suitable examples. 1
- How will you convert chloroethane into butane? 1
- Out of  $\text{C}_6\text{H}_5\text{CH}_2\text{Cl}$  and  $\text{C}_6\text{H}_5\text{CHCl}$ , which is more easily hydrolysed by aqueous KOH? 1
- (i) Chloroform is kept in completely filled bottles in dark. Explain the reason. 2  
 (ii) Mention one application of Iodoform 2
- How Grignard's reagent is prepared using alkyl halides? Why it is kept under anhydrous conditions? 2
- Explain following terms giving examples: 2
  - Chirality
  - Inversion of configuration
- Discuss mechanisms of  $\text{S}_{\text{N}}1$  and  $\text{S}_{\text{N}}2$  taking appropriate examples of optically active alkyl halides. 3
- Which of the following undergoes  $\text{S}_{\text{N}}2$  faster and why? 3
  - 
 Or 
  - 
 Or 
  - 
 Or 
- Convert following: 3
  - Aniline to phenylisocyanide
  - Benzene to 4-Bromonitrobenzene
  - Propene to Propan-1-ol


## UNIT 7

### Alcohols, Phenols and Ethers


#### Points to Remember

##### Classification of Alcohols:

(a) Containing  $sp^3$  C-OH bond. (i)  $-CH_2-OH$  ( $1^\circ$ ), (ii)  $>CH-OH$  ( $2^\circ$ ), (iii)  $\text{---}C\text{---}OH$  ( $3^\circ$ )  
Allylic:  $CH_2=CH-CH_2OH$


(iv) Benzylic: 

(b) Containing  $sp^2$  C-OH bond - Vinylic (i)  $CH_2=CH-OH$

(ii) Phenols: 

##### Classification of ethers:

Simple/Symmetric: Alkyl/aryl groups attached to oxygen are same e.g.



Mixed/Unsymmetric: Two different alkyl/aryl groups attached to oxygen e.g.



#### Classification and Nomenclature of Alcohols and ethers

##### Nomenclature:

**Alcohols:** Common names (Alkyl group + alcohol)

**IUPAC** (Alkan+ol, substituting -e by -ol e.g. ethanol)

**Phenols:** Common names (as derivatives of phenol with position like ortho-, meta-, para-)


**IUPAC** (Derivatives of phenols with numbers like 1,2-, 1,3-e.g.

2-Nitrophenol or o-nitrophenol)



**Ethers:** Common names (alkyl/aryl groups in alphabetical order followed by ether e.g. ethyl methyl ether) IUPAC (In alkyl/aryl group -e is replaced by oxy followed by parent hydrocarbon e.g. methoxyethane)

**Preparation of Alcohols****1. From alkenes:**



a) By acid catalyzed hydration Markonikov's addition)





b) By hydroboration-Oxidation(Anti-Markonikov)

**2. From carbonyl compounds**

a) By reduction of aldehydes and ketones










b) By reduction of carboxylic acids and esters

**3. From Grignard reagent**

For Primary alcohol  $\rightarrow$  methanal (formaldehyde)

Secondary  $\rightarrow$  aldehydes other than methanal

Tertiary  $\rightarrow$  Ketones are used with appropriate Grignard reagent



**Preparation of Phenol****1. From Haloalkanes****2. From Benzene sulphonic acid****3. From Diazonium salts****4. From Cumene****Preparation of Alcohols, Phenols and Ethers****Preparation of Ethers****1. By dehydration of alcohols****2. Williamson's Synthesis**

### Physical Properties and Reactions of Alcohols and phenols




#### (a) Physical properties:



- Boiling point increases with increase of carbon chain
- Solubility decreases with increase in size of alkyl aryl group

#### (b) Chemical properties:





**Order:** Primary < Secondary < Tertiary




**Mechanisms:****a) Hydration of Alkenes****Mechanism of hydration of ethene**


The mechanism of the reaction involves the following three steps:


**Step 1: Protonation of alkene to form carbocation by electrophilic attack of  $\text{H}_3\text{O}^+$ .****Step 2: Nucleophilic attack of water on carbocation.****Step 3: Deprotonation to form an alcohol.****b) Reaction of Grignard reagent**

The first step of the reaction is the nucleophilic addition of Grignard reagent to the carbonyl group to form an adduct. Hydrolysis of the adduct yields an alcohol.





The overall reactions using different aldehydes and ketones in dry ether medium are as follows:



**c) Dehydration of Alcohols****Step 1: Formation of protonated alcohol.**




**Step 2: Formation of carbocation: It is the slowest step and hence, the rate determining step of the reaction.**

**Step 3: Formation of ethene by elimination of a proton.**

The acid used in step 1 is released in step 3. To drive the equilibrium to the right, ethene is removed as it is formed.

**d) Dehydration of alcohol at 413K to form alkene**

The formation of ether is a nucleophilic bimolecular reaction ( $\text{S}_{\text{N}}2$ ) involving the attack of alcohol molecule on a protonated alcohol, as indicated below:




Acidic dehydration of alcohols, to give an alkene is also associated with substitution reaction to give an ether.

**c) Reaction of ethers with HI**

The reaction of an ether with concentrated HI starts with with protonation of ether molecule.


Step 1:



Step 2:


Iodide is a good nucleophile. It attacks the least substituted carbon of the oxonium ion formed in step 1 and displaces an alcohol molecule by  $\text{S}_{\text{N}}2$  mechanism.

Thus, in the cleavage of mixed ethers with two different alkyl groups. The alcohol and alkyl iodide formed, depend on the nature of alkyl groups. When primary or secondary alkyl groups are present, it is the lower alkyl group that forms alkyl iodide ( $\text{S}_{\text{N}}2$  reaction).



When HI is in excess and the reaction is carried out at high temperature, ethanol reacts with another molecule of HI and is converted to ethyl iodide.

Step 3:





### OBJECTIVE TYPE QUESTIONS

#### I. MULTIPLE CHOICE QUESTIONS

1. Arrange the following compound in decreasing order of boiling point?

(i) Propan-1-ol

(ii) Butan-2-ol

(iii) Butan-1-ol

(iv) Pentan-1-ol


(a)  $i > iii > ii > iv$

(b)  $i > ii > iii > iv$

(c)  $iv > iii > ii > i$

(d)  $iv > ii > iii > i$

2. What is the correct order of reactivity of alcohols in the following reaction?




(a)  $1^\circ > 2^\circ > 3^\circ$

(b)  $1^\circ > 3^\circ > 2^\circ$

(c)  $3^\circ > 1^\circ > 2^\circ$

(d)  $3^\circ > 2^\circ > 1^\circ$

3. IUPAC name of the compound :






(a) 1-Methoxy-1-methyl ethane


(b) 2-Methoxy-2-methyl ethane

(c) 2-Methoxypropane








(d) Isopropylmethyl ether

4. The correct order of decreasing acid strength of the following compound is:




- (a)  $i > ii > iii$  (b)  $iii > ii > i$   
 (c)  $ii > i > iii$  (d)  $ii > iii > i$
5. The major product obtained on reaction of phenol with NaOH followed by  $\text{CO}_2$  and acidification is:  
 (a) Benzoic acid (b) Salicylaldehyde  
 (c) Salicylic acid (d) Phthalic acid
6. Which of the following alcohol on dehydration with conc.  $\text{H}_2\text{SO}_4$  gives but-2-ene?  
 (a) Butan-1-ol (b) Butan-2-ol  
 (c) 2-Methylpropan-1-ol (d) Both (a) and (b)
7. Which of the following alcohol give iodoform test?  
 (a) Butan-1-ol (b) Propan-1-ol  
 (c) Propan-2-ol (d) Pentan-3-one
8. In the following sequence of reaction. The product D is
-   $\xrightarrow{\text{PCl}_3}$  A  $\xrightarrow[\text{ether}]{\text{Mg}}$  B  $\xrightarrow{\text{HCHO}}$  C  $\xrightarrow{\text{H}_3\text{O}^+}$  D
- (a) n-Butyl alcohol (b) n-Propyl alcohol  
 (c) Propanal (d) Butanal
9. Correct statement in case of n-butanol and t-Butanol is:  
 (a) both are having equal solubility in water  
 (b) 1-Butanol is more soluble in water  
 (c) boiling point of t-butanol is lower than n-butanol  
 (d) boiling point of n-butanol is lower than t-butanol
10. The major products of following reaction are:




- (a)  +  $\text{CH}_3\text{-I}$  (b)  + 
- (c)  +  (d)  + 


11. Write the IUPAC name of the compound given below.



- (a) 3-Methylpent-2-ene-1,2-diol  
 (b) 2-Methylpent-2-ene-1,2-diol  
 (c) 3-Methylpent-3-ene-2,3-diol  
 (d) 3-Methylpent-3-ene-4,5-diol
12. Which of the following are used to convert  $\text{RCHO}$  into  $\text{RCH}_2\text{OH}$  ?  
 (a)  $\text{H}_2/\text{Pd}$  (b)  $\text{LiAlH}_4$   
 (c)  $\text{NaBH}_4$  (d) All of the above
13. Monochlorination of toluene in sunlight followed by hydrolysis with aq.  $\text{NaOH}$  yields-  
 • (a) *o*-cresol (b) *m*-cresol  
 (c) 2,4-Dihydroxytoluene (d) Benzyl alcohol
14. The product C in given reaction will be



- (a) (b)   
 (c) (d)

15. Which of the following species can act as the strongest base?
- $^-\text{OH}$
  - $^-\text{OR}$
  - $^-\text{OC}_6\text{H}_5$
  - 
16.  $\text{CH}_3\text{CH}_2\text{OH}$  can be converted into  $\text{CH}_3\text{CHO}$  by.....
- catalytic hydrogenation
  - treatment with  $\text{LiAlH}_4$
  - treatment with pyridinium chlorochromate (PCC)
  - treatment with  $\text{KMnO}_4$
17. Which of the following reactions will yield phenol?
- Fusion of chlorobenzene with  $\text{NaOH}$  at 300 atm.
  - Diazotization of aniline followed by heating with water.
  - Sulphonation of benzene followed by treatment with  $\text{NaOH}$  then acidification.
  - All of the above
18. Williamson's synthesis is used to prepare
- alcohol
  - ethers
  - Aldehydes
  - Amines
19. Which of the following compound will be most readily attacked by electrophile?
- Chlorobenzene
  - Benzene
  - Phenol
  - Toluene
20. Among the following sets of reactants which one produces anisole?
- $\text{CH}_3\text{CHO}$  ;  $\text{RMgX}$
  - $\text{C}_6\text{H}_5\text{OH}$  ;  $\text{NaOH}$  ;  $\text{CH}_3\text{Br}$
  - $\text{C}_6\text{H}_5\text{OH}$  ; Neutral  $\text{FeCl}_3$
  - $\text{C}_6\text{H}_5\text{-CH}_3$  ;  $\text{CHCOCl}$  ;  $\text{AlCl}_3$

## . II FILL IN THE BLANKS

- Phenol on reaction with bromine water gives white precipitate of.....
- Ethanoic acid on reaction with  $\text{LiAlH}_4$  forms.....
- Reaction of.....with Grignard reagent gives primary alcohols.
- Phenols are commercially manufactured by oxidation followed by acidification of.....
- Reaction of alcohols/phenols with carboxylic acids is termed as.....
- .....test is used to distinguish primary, secondary and tertiary alcohols.
- Dehydration of tertiary alcohols is .....than primary alcohols.
- Dehydration of alcohols to form ether is ..... type reaction.
- Benzyl alcohol on reaction with  $\text{KMnO}_4$  followed by acidification forms.....
- $\text{SOCl}_2$  converts Propan-1-ol to .....

### III ASSERTION REASON TYPE QUESTIONS


In the following questions a statement of Assertion followed by a statement of Reason is given. Choose the correct answer out of the following choices.

- (A) Assertion and reason both are correct and reason is correct explanation of assertion
- (B) Both assertion and reason are correct statement but reason is not correct explanation of assertion.
- (C) Assertion is correct statement but reason is wrong statement.
- (D) Assertion is wrong statement but reason is correct statement.

1. **ASSERTION :** *p*-Nitrophenol is more acidic than phenol.

**REASON :** Nitro group helps in the stabilization of the phenoxide ion by dispersal of negative charge due to resonance.

2. **ASSERTION :** IUPAC name of the compound is 2-Ethoxy-2-methylethane.



**REASON :** In IUPAC nomenclature, ether is regarded as hydrocarbon derivative in which a hydrogen atom is replaced by -OR or -OAr group [where, R= alkyl group and Ar = aryl group].

3. **ASSERTION :** Bond angle in ethers is slightly less than the tetrahedral angle.

**REASON :** There is a repulsion between the two bulky (-R) groups.

4. **ASSERTION :** *o*-Nitrophenol is less soluble in water than the *m* and *p*-isomers.

**REASON:** *m* and *p*-Nitrophenols exist as associated molecule.

5. **ASSERTION :** Like bromination of benzene, bromination of phenol is also carried out in the presence of Lewis acid.

**REASON:** Lewis acid polarises the bromine molecule.

6. **ASSERTION :** Ethanol is a weaker acid than phenol.

**REASON :** Sodium ethoxide may be prepared by the reaction of ethanol with aqueous NaOH.

7. **ASSERTION :** Phenols give *o*- and *p*-Nitrophenol on nitration with dil.  $\text{HNO}_3$ .

**REASON :** -OH group in phenol is *o*-, *p*- directing.


8. **ASSERTION :**  $(\text{CH}_3)_3\text{C}-\text{O}-\text{CH}_3$  gives  $(\text{CH}_3)_3\text{C}-\text{I}$  and  $\text{CH}_3\text{OH}$  on treatment with HI.  
**REASON :** The reaction occurs by  $\text{S}_\text{N}1$  mechanism
9. **ASSERTION :** Protonation of phenol is difficult than ethanol.  
**REASON :** Ethyl group in ethanol is electron releasing.
10. **ASSERTION :** Tertiary alcohols give turbidity almost immediately on treatment with  $\text{ZnCl}_2/\text{HCl}$ .  
**REASON :** Tertiary carbocation formed is very stable and undergoes substitution easily.

#### IV ONE WORD TYPE QUESTIONS

- Which reagent is used to convert primary alcohol to carboxylic acid?
- Out of *ortho* and *para*-Nitrophenol which is more steam volatile?
- What is the major product of bromination of anisole?
- Which reaction is used to convert phenol to salicylaldehyde?
- Out of ethanol and phenol, which is more acidic?
- Which reagent could be used to reduce aldehyde selectively in presence of ester group?
- Which reagent(s) are best to convert propene to propan-1-ol?
- Out of primary, secondary and tertiary alcohols which is most acidic?
- Name the product formed, when phenol is treated with cone.  $\text{HNO}_3$ .
- Draw structure of hex-1-en-3-ol compound
- Write the name of chemical test to distinguish between pentan-3-one and pentan-2-one
- \_\_\_\_\_ is added into commercial ethanol to provide a foul smelling.
- \_\_\_\_\_ bonding is responsible for solubility of ethanol in water.
- Name the product formed when phenol is treated with  $\text{Na}_2\text{Cr}_2\text{O}_7/\text{H}_2\text{SO}_4$
- What is the IUPAC name of Aspirin.

**VERY SHORT ANSWER TYPE QUESTIONS (1 Mark)**


1. Write IUPAC name of the following compound:



Ans: 2,5-Dimethylhexan-1,3-diol

2. How is phenol obtained from aniline ?

Ans:



3. Why phenol is acidic in nature ?

Ans: Due to stability of phenoxide ion by resonance

4. Among HI, HBr and HCl which is most reactive towards alcohols. Why?

Ans: Due to lowest bond dissociation energy of HI, it is most reactive.


5. Name a compound which is used as antiseptic as well as disinfectant'.

Ans: Solution of phenol : 0.2% antiseptic, 2% disinfectant

6. What is nitrating mixture for monosubstitution of phenol?

Ans: Dilute  $\text{HNO}_3$

7. Arrange the following in decreasing order of their acidic character:




Ans: (c) > (b) > (a)


8. Why lower alcohols are soluble in water while higher alcohols are not?

Ans: Due to formation of hydrogen bonds, lower alcohols are soluble but increase in hydrophobic chain decreases solubility.

9. Complete the following reaction :



Ans:



10. What happens when  $\text{CH}_3\text{CH}_2\text{OH}$  heated with red P and HI ?


Ans.:



11. Ethanol has higher boiling point than Methoxymethane. Give reason.


Ans : Because of H-bonding in ethanol.

12. Explain Kolbe's reaction with example.




13. How could you convert ethanol to ethene ?

Ans.




14. Explain Reimer-Tiemann reaction.







15. How will you get benzoquinone from phenol?




16. Predict the major product of acid catalysed dehydration of 1-Methylcyclohexanol



17. What is the significance of pyridine in following reaction.




Ans. To remove HCl from product side and make the reaction to shift forward.




Ans:- 1-Ethoxy-2-nitrocyclohexane.

19. How is acetone obtained from 2-bromopropane ?



20. Which alcohol contain the -OH group attach to a  $sp^3$  hybridised carbon atom next to an aromatic ring.

Ans. Benzylic  $\text{CH}_2 - \text{OH}$



21. Which chemical is used as wood spirit ?

Ans. Methanol  $\text{CH}_3\text{OH}$

22. Why boiling point of ethers are much lower than those of alcohols of comparable molecular masses.

Ans. There is no H-bonding in ethers which is present in alcohol.

23. Which test is used to distinguish between phenol and Benzyl Alcohol.


Ans. Neutral  $\text{FeCl}_3$

Phenol gives violet colour with aq.  $\text{FeCl}_3$  while benzyl alcohol does not.

24. Why does phenol not undergo protonation easily.

Ans. In phenol, there is +ve charge, in its resonance structure, therefore does not undergo protonation.



25. Write the steps involved in Conversion of phenol to anisole.



**SHORT ANSWER TYPE QUESTIONS (2 or 3 Marks)**



- Out of t-Butyl alcohol and n-Butanol, which one will undergo acid catalysed dehydration faster and why?  
 Ans: t-Butyl alcohol will undergo dehydration faster due to formation of stable tertiary carbocation intermediate.
- Carry out the following conversions:
  - Phenol to salicylaldehyde
  - t-Butylchloride to t-Butyl ethyl ether
  - Propene to propan-1-ol
- Write the steps involved in the mechanism for the formation of ethanol from ethene.
- Predict the reagent for carrying out the following conversions:
  - Phenol to Benzoquinone
  - Anisole to *p*-Bromoanisole
  - Phenol to 2,4,6-Tribromophenol
- Write one chemical reaction to illustrate the following:
  - Reimer-Tiemann reaction
  - Williamson synthesis

Ans:






- Complete the following the equations and name the products:





Ans: (i)  $[\text{Fe}(\text{C}_6\text{H}_5\text{O})_3]^{3-} + \text{HCl}$




7. Write an example for the following name reactions :

(a) Friedel-Crafts reaction

(b) Coupling reaction

Ans: (a)



(b)



8. Account for the following:

- (a) Phenol has a smaller dipole moment than methanol.  
 (b) Phenol undergoes electrophilic substitution reactions faster than benzene.

Ans: (a) Due to delocalization of electrons of oxygen in phenol.

- (b) Due to +R effect of -OH group in phenol which activates phenyl nucleus by increasing electron density as compared to benzene.

9. Give one reaction of alcohol involving cleavage of:

- (a) C-O bond (b) O-H bond

Ans: (a)





(b)



10. Ethereal solution of an organic compound 'X' when heated with Mg gave 'Y' which on treatment with  $\text{CH}_3\text{CHO}$  followed by acid hydrolysis gave 2-Propanol. Identify the compound 'X'. What is 'Y' known as?

Ans:



11. Phenol is more acidic than alcohol:

Ans: Due to resonance stabilised phenoxide ion.

12. While separating a mixture of *o*- and *p*-Nitrophenols by steam distillation, name the isomer which is steam volatile? Give reasons.



Ans: *o*-Nitrophenol is steam volatile because it has intramolecular H-bonding.

13. Write the reactions and conditions involved in the conversion of:

(a) Propene to propan-2-ol


(b) Phenol to salicylic acid

Ans:



14. Write the chemical reaction of HI with Methoxymethane.

Ans:




15. Ethers are relatively inert. Justify

Ans: Due to absence of any active site in their molecules, divalent oxygen is linked to carbon atoms on both sides (C-O-C).

16. How will you distinguish between  $\text{CH}_3\text{OH}$  and  $\text{C}_2\text{H}_5\text{OH}$ ?


Ans:  $\text{C}_2\text{H}_5\text{OH} + 4\text{I}_2 + 3\text{Na}_2\text{CO}_3 \xrightarrow{\text{heat}} \text{CH}_3\text{I} + \text{HCOONa} + 5\text{NaI} + 2\text{H}_2\text{O} + 3\text{CO}_2$

17. Which of the following is an appropriate set of reactants for the preparation of 1-Methoxy-4-nitrobenzene and why?



Ans:

18. Arrange in order of boiling points :



Ans: (a)  $C_4H_9COOH > C_4H_9OH > C_2H_5-O-C_2H_5$






19. Describe the following reactions with examples :

(a) Reimer-Tiemann reaction

(b) Kolbe's reaction

(c) Friedel Crafts acylation of anisole

Ans:




20. Give equations of the following reactions:

(a) Bromine in  $CS_2$  with phenol

(b) Treating phenol with chloroform in presence of aqueous  $NaOH$


(c) Oxidation of propan-1-ol with alkaline  $KMnO_4$  solution.



21. Write the structure of the major products of the following:

- Mononitration of 3-Methylphenol
- Dinitration of 3-Methylphenol
- Mononitration of phenyl ethanoate

Ans : -OH and -CH<sub>3</sub> are *o*- and *p*-directing groups. The products are:





22. Dehydration of alcohols to form an alkene is always carried out with conc.  $\text{H}_2\text{SO}_4$  and not with conc.  $\text{HCl}$  or  $\text{HNO}_3$ . Explain.

Ans: In acidic medium alcohols protonated then loses  $\text{H}_2\text{O}$  to form a carbocation. If  $\text{HCl}$  is used which is strong nucleophile causes nucleophilic substitution and  $\text{HNO}_3$  causes oxidation.

23. Name the reagents which are used in the following conversions:

- Primary alcohol to an aldehyde
- Butan-2-one to Butan-2-ol
- Phenol to 2,4,6-Trinitrophenol

Ans: (i) PCC, a complex of chromium trioxide with pyridine and  $\text{HCl}$ .



(ii)  $\text{NaBH}_4$ , sodium borohydride

(iii) Conc  $\text{HNO}_3 + \text{H}_2\text{SO}_4$

24. Write major products of following reactions:



Ans: (i)  $\text{CH}_3\text{CH}_2\text{CH}_2\text{CH}_2\text{OH}$





25. How will you convert :

(a) Phenol to Cyclohexanol

(b) Benzyl chloride to Benzyl alcohol

Ans: (b) Anisole to phenol



### LONG ANSWER TYPE QUESTIONS (5 Marks)


1. An alcohol A ( $\text{C}_4\text{H}_{10}\text{O}$ ) on oxidation with acidified  $\text{K}_2\text{Cr}_2\text{O}_7$  gives carboxylic acid 'B' ( $\text{C}_4\text{H}_8\text{O}_2$ ). Compound 'A' when dehydrated with conc.  $\text{H}_2\text{SO}_4$  at 443K gives compound 'C' with aqueous  $\text{H}_2\text{SO}_4$ . 'C' gives compound 'D' ( $\text{C}_4\text{H}_{10}\text{O}$ ) which is an isomer of 'A'. Compound 'D' is resistant to oxidation but compound 'A' can be easily oxidised. Identify A, B, C and D and write their structure.

Ans: A:  $(\text{CH}_3)_2\text{CHCH}_2\text{OH}$  C:  $(\text{CH}_3)_2\text{C}=\text{CH}_2$



B:  $\text{CH}_3\text{CH}(\text{CH}_3)\text{COOH}$  D:  $(\text{CH}_3)_3\text{C-OH}$

2. An ether 'A' ( $\text{C}_5\text{H}_{12}\text{O}$ ) when heated with excess of hot conc. HI produced two alkyl halides which on hydrolysis form compound 'B' and 'C'. Oxidation of B gives an acid 'D' whereas oxidation of 'C' gave a ketone E. Deduce the structure of A, B, C, D and E.

Ans:



3. a) Which of the following compounds gives fastest reaction with HBr and why?



Ans: (i)  $(\text{CH}_3)_3\text{C-OH}$



Due to formation more stable of carbocation

b) How are following obtained

(i) Toluene from Phenol


(ii) Phenol from Aniline.

Ans. (i)




4. Phenol,  $C_6H_5OH$  when reacts with concentrated sulphuric acid, forms 'Y'. The compound, 'Y' is reacted with concentrated nitric acid to form 'Z'. Identify 'Y' and 'Z' Explain why phenol is not converted commercially to Z by reacting it with cone.  $HNO_3$

Ans:








(Y)




(Z)

5. Write the structure of the main product in the following reactions:




Ans. (i)



An organic compound 'A' ( $C_2H_6O$ ) reacts with sodium to form a compound 'B' with the evolution of  $H_2$  and gives a yellow compound 'C' on reacting with Iodine and NaOH. When heated with conc.  $H_2SO_4$  on reaction with conc. HI at 373 K gives compound 'E'. The compound 'D' is also obtained when 'B' is heated with 'E'. Identify A, B, C, D, E. Write the equation for the reaction involved.

Ans. A- $CH_3CH_2OH$




7. Two compounds [A] and [B] have molecular formula  $C_2H_6O$  on reacting with HI, [A] gives alkyl iodide and water while [B] give alkyl iodide and alcohol. What are the compounds [A] and [B] write the reaction involved.

Ans. A- $C_2H_5OH + HI \longrightarrow C_2H_5I + H_2O$   
 B- $CH_3OCH_3 + HI \longrightarrow CH_3I + CH_3OH$


8. A compound [A] $C_4H_{10}O$  is found to be soluble in sulphuric acid. [A] does not react with Na or  $KMnO_4$ . On heating with excess of HI, it is converted into single alkyl halide, what is compound [A]

Ans. [A]- $CH_3CH_2O-CH_2CH_3 + HI \longrightarrow 2 CH_3CH_2I$

9. (a) Write the equation for the following reaction  
 (i) Friedel-craft reaction - alkylation in anisole  
 (ii) Nitration of anisole  
 (iii) Bromination of anisole in ethanolic medium  
 (b) Explain alkyl aryl ether directs the incoming substituents towards ortho and para position in the ring



(b) Due to resonance




10. (i) Write the chemical distinguish test between followings
- Ethanol and Propan-2-ol
  - Butan-1-ol and Phenol
  - Propan-2-ol and pentan-3-ol
  - Phenol and Anisole
- (ii) Name the reagent needed to perform the following reaction;
- Dehydrogenation of ethanol to ethanal
  - Phenol to Benzene

Ans. (i) a) Lucas Test (anhy  $\text{ZnCl}_2 + \text{HCl}$ )  
 (b) aq.  $\text{FeCl}_3$  solution  
 (c) Iodoform test  
 (ii) a)  $\text{Cu}/573\text{K}$   
 (b) zinc dust

### CASE-STUDY BASED QUESTIONS

**1. Read the passage and answer the following questions:**

In the past few decades, phenols have received great attention in modern synthetic chemistry since ever Runge and Laurent made the first discovery in 1834 and 1841, respectively with regard to this motif, which is frequently found in natural products, flavonoids and pharmaceutically important compounds associated with certain bioactivities, such as antibacterial, antifungal, antibiotic, anti-inflammatory, antiviral, anxiolytic and antioxidant activities. Conventional methods for the large-scale synthesis of phenols include the Hock process, diazotization of aromatic amines and nucleophilic substitution reactions. Academicians have focused on the development of alternative approaches, for example, C-H activation of arenes and oxidation of C-Si bonds and C-halo bonds. Recently, the direct hydroxylation of aryl boronic acids to phenols has gained a lot of attention. In this context, a variety of oxidative methods employing metal catalysts,  $\text{Cu}(\text{OAc})_2\text{-H}_2\text{O}_2$ ,  $\text{CuSO}_4\text{-phenanthroline}$ ,  $\text{CuCl}_2\text{-miceller systems}$ ,  $\text{Cu}_2\text{O-NH}_3$ ,  $[\text{Ru}(\text{bpy})_3\text{Cl}_2]\cdot 6\text{H}_2\text{O}$ ,  $\text{Al}_2\text{O}_3\text{-H}_2\text{O}_2$ , and  $\text{H}_3\text{BO}_3\text{-H}_2\text{O}_2$  has been developed. On the other hand, the metal-free oxidative process are also competitive, Oxone,  $n\text{Bu}_4\text{NHSO}_5$ ,  $\text{NH}_2\text{OH}$ ,  $\text{H}_2\text{O}_2\text{-poly(N-vinylpyrrolidone)}$ ,  $\text{I}_2\text{-H}_2\text{O}_2$ , Amberlite IR-120- $\text{H}_2\text{O}_2$ , N-oxides, MCPBA,  $\text{NaClO}_2$ , photoredox catalysis, electrochemical oxidation,  $(\text{NH}_4)_2\text{S}_2\text{O}_8$ , PEG-400- $\text{H}_2\text{O}_2$ , WERSA- $\text{H}_2\text{O}_2$ , WEBPA- $\text{H}_2\text{O}_2$ , nanoparticles of Ag,  $\text{Cu}_2\text{O}$ , and  $\text{Fe}_2\text{O}_3/\text{silica gel}$  and TBHP/ $\text{C}_{13}\text{CCN}$ . Despite these efficient oxidative processes, developing a new methodology free from metal oxidants and organic solvents is highly desirable. As part of our research interest involving metal-free oxidation reactions herein, a new protocol for the direct hydroxylation of aryl boronic acids with TBHP in the aqueous medium is reported (Scheme 1).



Scheme 1: Hydroxylation of aryl boronic acids.


**Reference :** Tanveer MahmadAlli Shaikh, **Synthesis of Phenols via Metal-Free Hydroxylation of Aryl Boronic Acids with Aqueous TBHP**, *Journal of Chemistry*, vol. 2020, Article ID 1543081, 7 pages, 2020. <https://doi.org/10.1155/2020/1543081>

- (A) Which of the following method of preparation of phenols is least likely to take place at 298K?
- Nucleophilic substitution in chlorobenzene
  - Reaction of diazonium salt with water.
  - Oxidation followed by acidification of cumene
  - Reaction of benzene sulphonic acid with NaOH followed by acidification
- (B) In which of the following haloarene, nucleophilic substitution will be fastest to yield corresponding phenol?
- Chlorobenzene
  - p*-Chloronitrobenzene
  - p*-Chlorotoluene
  - p*-Chloroanisole
- (C) Aniline on reaction with  $\text{NaNO}_2 + \text{HCl}$  forms (X). (X) converts to (Y) on reaction with KI. (X) and (Y) are respectively-
- Benzene diazonium chloride, iodobenzene
  - Iodobenzene, Ethoxybenzene
  - Iodobenzene, Benzene
  - Benzene diazonium chloride, Phenol
- (D) Phenols on reaction with bromine water forms-
- Colourless, 2-Bromophenol
  - Dark coloured mixture of 2-Bromophenol and 4-Bromophenol
  - White precipitate of 2,4,6-Tribromophenol
  - Yellow colouration of 2,4-Dibromophenol

2. **Read the passage and answer the following questions:**

On the basis of the investigation of the combinational effect of quaternary ammonium salts and organic bases, an added-metal-free catalytic system for nucleophilic addition reactions of a variety of Grignard reagents to diverse ketones in the solvent has been developed to produce tertiary alcohols in good to excellent yields. By using tetrabutylammonium chloride ( $\text{NBu}_4\text{Cl}$ ) as a catalyst and diglyme (DGDE) as an additive, this system strongly enhances the efficiency of





addition at the expense of enolization and reduction.  $\text{NBu}_4\text{Cl}$  should help to shift the Schlenk equilibrium of Grignard reagents to the side of dimeric Grignard reagents to favor the additions of Grignard reagents to ketones via a favored six-membered transition state to form the desired tertiary alcohols, and DGDE should increase the nucleophilic reactivities of Grignard reagents by coordination. This catalytic system has been applied in the efficient synthesis of Citalopram, an effective U.S. FDA-approved antidepressant, and a recyclable version of this catalytic synthesis has also been devised.

**Reference :** Hua Zong, Huayin Huang, Junfeng Liu, Guangling Bian, and Ling Song **Added-Metal-Free Catalytic Nucleophilic Addition of Grignard Reagents to Ketones** *J. Org. Chem.* **2012**, 77, 10, 4645-4652

- (A) Which ketone and Grignard reagent can be used to form 2-methylbutan-2-ol?
- (B) Write structure and IUPAC name of product formed reaction of allyl magnesium bromide with acetophenone?
- (C) Which reaction will take place at faster rate and why?
  - (i) Benzaldehyde + Propyl magnesium bromide
  - (ii) Propanal + Benzyl magnesium bromide
- (D) Why Grignard reagent is stored under anhydrous conditions?





3. **Read the passage and answer the following questions:**

Phenols are compounds that possess a hydroxyl group directly attached to an aromatic carbocyclic nucleus. Phenol is the trivial name for monohydroxybenzene. The *o*-, *m*-, and *p*-cresols are monohydroxytoluenes ( $\text{CH}_3\text{-C}_6\text{H}_4\text{OH}$ ) and are distinct in their properties and reactions from the isomeric side-chain hydroxy compound, benzyl alcohol ( $\text{C}_6\text{H}_5\text{-CH}_2\text{OH}$ ), which is a typical aromatic alcohol. Simple monohydric phenols are either corrosive liquids or low melting solids. The dihydric and trihydric phenols are solids. The mono-hydroxy compounds are only slightly soluble in water but are miscible with organic solvents. Water solubility increases and solubility in organic solvents

decreases with the introduction of additional hydroxyl groups. They are all characterized by, and distinguished from, the aliphatic or aromatic alcohols by their ready solubility in aqueous alkali. Phenols and the cresols are widely used as antiseptics and disinfectants; the cresols are contained in the wood preserving fluid, creosote. Many phenols have wide application in the industrial production of plastics, dyestuffs, insecticides, selective weedkillers, and germicides.

**Reference :** P.W.G. Smith, A.R. Tatchell, **Phenols**, *Aromatic Chemistry*, 1969

(A) Which of the following is not a phenol?



(B) Which of the following phenol has highest  $pK_a$  value?

(a) Phenol (b) p-Nitrophenol


(c) o-Nitrophenol (d) o-cresol

(C) Phenols may be characterized by the reaction with-

(a)  $FeCl_3$  (b)  $Br_2$  water

(c)  $NaHCO_3$  (d) Both  $FeCl_3$  and  $NaHCO_3$

(D) Write IUPAC name of following phenol.



## ANSWERS

### I MULTIPLE CHOICE QUESTIONS

1. c 2. d 3. c 4. c 5. c 6. d 7. d 8. b 9. c 10. c

11. a 12. d 13. d 14. a 15. b 16. c 17. d 18. b 19. c 20. b

**II FILL IN THE BLANKS**

- |                         |                     |
|-------------------------|---------------------|
| 1. 2,4,6-Tribromophenol | 2. Ethanol          |
| 3. Methanal             | 4. Cumene           |
| 5. Esterification       | 6. Lucas            |
| 7. Easier               | 8. $S_N2$           |
| 9. Benzoic acid         | 10. 1-Chloropropane |

**III ASSERTION REASON TYPE QUESTIONS**

1. (A) 2. (D) 3. (D) 4. (C) 5. (D) 6. (C) 7. (A) 8. (A) 9. (B) 10. (A)

**IV ONE WORD ANSWER**

- PCC
- ortho*-Nitrophenol
- para*-bromoanisole
- Reimer-Tiemann reaction
- Phenol
- $\text{NaBH}_4$
- $\text{B}_2\text{H}_6/\text{H}_2\text{O}_2, \text{OH}^-$
- Primary
- 2,4,6-Trinitrophenol (Picric Acid)
- $\text{H}_3\text{C}=\overset{1}{\text{C}}=\overset{2}{\text{CH}}-\overset{3}{\underset{\text{OH}}{\text{C}}}-\overset{4}{\text{CH}_2}-\overset{5}{\text{CH}_2}-\overset{6}{\text{CH}_3}$
- Iodoform Test
- Pyridine
- Intermolecular H-bond
- Benzoquinone
- 2-Acetoxybenzoic acid

**CASE STUDY QUESTIONS**

PASSAGE: 1 : (A) b (B) b (C) a (D) c

PASSAGE: 2:

(A)  $\text{CH}_3\text{COCH}_3, \text{C}_2\text{H}_5\text{MgCl}$ 

(B)  $\text{C}_6\text{H}_5-\overset{\text{OH}}{\underset{\text{CH}_3}{\text{C}}}-\text{CH}_2-\text{CH}=\text{CH}_2$  IUPAC (i) 1-Methyl-1-phenyl but -3-ene-1-ol

(C) i

(D) It react with water to form alkane  $\text{R-MgX} + \text{H-OH} \rightarrow \text{R-H} + \text{Mg-X-OH}$ 


PASSAGE: 3: (A) d (B) d (C) a (D) 3-Ethyl-5-Chloro Phenol

**UNIT TEST**  
**CHAPTER-10**

Maximum Marks : 20

Time : 1 Hour

1. Write the IUPAC name of:




2. A and B in the following reaction are :




- |     |    |  |    |  |
|-----|----|--|----|--|
| (a) | A. |  | B. |  |
| (b) | A. |  | B. |  |
| (c) | A. |  | B. |  |
| (d) | A. |  | B. |  |

3. The correct order of acidic strength of following phenols:




- (a)  $A < B < C < D$
- (b)  $B < A < C < D$
- (c)  $D < A < B < C$
- (d)  $D < C < B < A$

4. Identify the product: a



5.  $(\text{CH}_3)_3\text{C}-\text{OCH}_3 + \text{HI} \rightarrow ?$
6. Give reactions for following name reactions
- (I) Kolbe's reaction
  - (ii) Williamsons synthesis
7. Write the products formed by nitration of phenol. Which of the product has higher boiling point and why?
8. Identify the product formed when ethanol is heated at 413 K. Write the mechanism of the reaction.
9. Complete the reaction sequence, A to E. Also name the reaction involved in conversion of B to D.



10. How will you convert :
- (i) Ethanol to propan-2-ol
  - (ii) Aniline to phenol
  - (iii) Methanol to methoxyethane
11. Explain the following:
- (i) Alcohols are more soluble in water than ethers of comparable molar mass.
  - (ii) t-Butylchloride on heating with sodium methoxide gives 2-Methylpropene instead of t-Butylmethly ether.
  - (iii) Reaction of phenol with bromine leads to formation of 2,4,6- Tribromophenol


## UNIT TEST-2

## CHAPTER-10



Maximum Marks : 20

Time : 1 Hour

- Write the IUPAC name of  $C_6H_5-CH_2-CH_2-OH$
- Out of t-Butyl alcohol and n-Butanol, which one will undergoes acid catalysed dehydration faster.
- Write 'A' and 'B' in following reaction.



- Name the reagent used for conversion of primary alcohol to an aldehyde.
- Out of phenol and methanol which one has smaller dipole moment
- Write the mechanism of Conversion of propan-1-ol into propene.
- Write short note on williamson synthesis
- Complete the reaction.



- Convert the following
  - Phenol to picric acid
  - Propan-2-ol to 2-methyl propan-2-ol
  - Phenol to aspirin
- Write the reason of following observation
  - alcohols are generally soluble in water but alkyl halides are not.
  - Phenol exhibit an acidic character
  - o-nitrophenol is more volatile than p-nitrophenol
- An ether 'A' ( $C_5H_{12}O$ ) when heated with excess of hot conc H<sub>1</sub> produced two alkyl halides which on hydrolysis form compound 'B' and 'C' Oxidation of 'B' gives acid 'D' whereas oxidation of 'C' give ketone 'E'. Write A, B, C, D, E.

## UNIT 8

# Aldehydes, Ketones and Carboxylic

### Points to Remember

#### Nomenclature

##### Aldehydes and Ketones

**Common names:** replace -e from alkyl group by aldehyde or ketone e.g.  $\text{CH}_3\text{-CHO}$  is acetaldehyde


**IUPAC names:** replace -e by -al for aldehydes and -one for ketones e.g.  $\text{CH}_3\text{-CHO}$  is ethanal and  $\text{CH}_3\text{-COCH}_3$  is propanone.

##### Carboxylic Acids

**Common names:** end with -ic acid e.g.  $\text{CH}_3\text{CH}_2\text{COOH}$  is propionic acid

**IUPAC names:** replace -e in corresponding alkanes by -oic acid e.g.  $\text{CH}_3\text{CH}_2\text{COOH}$  is propanoic acid.



#### Structure of Carbonyl group


















Carbonyl carbon is  $\text{sp}^2$  Hybridised and planar in shape, with one sigma and one  $\pi$  bond between  $\text{C}=\text{O}$

**General Methods:****(a) Controlled oxidation/dehydrogenation of primary and secondary alcohols**


Primary alcohol gives aldehyde, secondary gives ketones, tertiary undergoes dehydration

**(b) Hydration of Alkynes**

## Preparation of Aldehydes, Ketones and Carboxylic Acids


**Aldehydes:****(a) From acyl chloride**  
(Rosenmund reduction)**(b) From nitriles and esters**  
(Stephen reaction)**(c) From hydrocarbons (Etard Reaction)****(d) (Side chain chlorination)****(e) (Gatterman-Koch reaction)****Ketones:****(a) From acyl chlorides****(b) from nitriles**  $\text{CH}_3\text{CH}_2\text{CN} + \text{C}_6\text{H}_5\text{MgBr}$ **(c) From benzene****Carboxylic Acids:****(a) From Primary alcohols/aldehydes****(b) From alkylbenzene****(c) From nitriles and amides****(d) From Grignard reagent****(e) From acyl halides and anhydrides****(f) From esters**




**Aldehyde and Ketones****Nucleophilic Addition reaction**

Reduction: Aldehydes and ketones convert to primary and secondary alcohols respectively in presence of  $\text{NaBH}_4$  /  $\text{LiAlH}_4$ .  $\text{NaBH}_4$  selectively reduces aldehydes and ketones only in presence of other carbonyl containing functional groups including carboxylic acids, esters etc.

Clemmensen reduction:





Wolf - Kishner Reduction:




Oxidation:

Tollens' Test.




Silver mirror

Fehling's Test



Red Ppt

Haloform reaction: given by methyl ketones





Reactions of Aldehydes,  
Ketones and Carboxylic Acids


**Carboxylic Acids**

**Acidic nature:**  $\text{R-COOH} + \text{NaHCO}_3 \rightarrow \text{RCOONa} + \text{H}_2\text{O} + \text{CO}_2$  Brisk effervescence

Formation of carboxylic acid derivatives:





This reaction is useful in reducing the size of carbon chain




## OBJECTIVE TYPE QUESTIONS

## I. MULTIPLE CHOICE QUESTIONS

1. Which product is formed when benzaldehyde is treated with concentrated KOH solution?



2. Structure of 'A' and type of isomerism in the above reaction are respectively-




- (a) Prop-1-en-2-ol, metamerism  
 (b) Prop-1-en-1-ol, tautomerism  
 (c) Prop-2-en-2-ol, geometrical  
 (d) Prop-1-en-2-ol, tautomerism
3. Compound A and C in the following reaction are:-







- (a) Identical                                      (b) Position isomer  
 (c) Functional group isomer              (d) Optical isomer
4. Toluene  $\xrightarrow{\text{KMnO}_4}$  A  $\xrightarrow{\text{SOCl}_2}$  B  $\xrightarrow[\text{BaSO}_4]{\text{H}_2/\text{Pd}}$  C  
 the product 'C' is:-





5. Among the following which has the lowest  $\text{pK}_a$  value:-




B The structure of 'B' is:-





7. The IUPAC name of compound is:



8. Which of the following reactions will not result in the formation of carbon-carbon bond?



9. Reduction of aldehydes and ketones into hydrocarbon using zinc amalgam and conc. HCl is called.....
- (a) Dow process (b) Cope reduction  
(c) Wolf-Kishner reduction (d) Clemmensen's Reduction
10. Which of the following compound do not undergo aldol condensation?
- (a)  $\text{CH}_3\text{CH}_2\text{CHO}$  (b)  $\text{CH}_3\text{CHO}$   
(c)  $\text{CH}_3\text{COCH}_3$  (d) 
11. The treatment of following compound with NaOH yields.....as major product.
- 
- (i) Phenol (ii) Sodium phenoxide  
(iii) Sodium benzoate (iv) Benzophenone  
(a) I,II (b) III only  
(c) I,III (d) IV only
12. Which of the following will undergo Cannizzaro reaction?
- (a) Acetophenone (b) Propanone  
(c) 2,2-Dimethylpropanal  
(d) Both acetophenone and 2,2-Dimethylpropanal
13. Which of the following neither gives Fehling's test nor iodoform test?
- (a) Benzaldehyde (b) Ethanal  
(c) Propanone (d) Acetophenone
14. Which reagent(s) is/are best for following conversion?  
 $\text{CH}_3\text{-CH=CH-CN} \rightarrow \text{CH}_3\text{-CH=CH}_2\text{-CHO}$
- (a)  $\text{H}_2, \text{Pd-BaSO}_4$  (b) DiBAL-H  
(c)  $\text{H}_2/\text{Ni}$  (d)  $\text{NaBH}_4$
15. Benzophenone can be obtained by the reaction of -
- (a) Benzoyl chloride + Benzene + anhy.  $\text{AlCl}_3$   
(b) Benzoyl chloride + Diphenyl anion  
(c) Benzoyl chloride + Phenyl magnesium chloride  
(d) Benzene + carbon monoxide +  $\text{ZnCl}_2$

16. Which of the following compounds will give butanone on oxidation with alkaline  $\text{KMnO}_4$  solution?
- (a) Butan-1-ol (b) Butan-2-ol  
(c) Both of these (d) None of these
17. Reaction of alkene **X** with  $\text{O}_3$  followed by  $\text{H}_2\text{O}_2/\text{Zn}$  forms Propanone and Ethanal. Alkene **X** is-
- (a) Pent-3-ene (b) Pent-2-ene  
(c) 2-Methylbut-2-ene (d) 2,2-Dimethylprop-1-ene
18. Which of the following on oxidation followed by hydrolysis give acetic acid ?
- (a) Actaldehyde cyanohydrin (b) Acetone cyanohydrin  
(c) Formaldehyde cyanohydrin (d) None of these
19. When 2-Hydroxy benzoic acid is distilled with Zinc dust it gives
- (a) Phenol (b) Benzoic Acid  
(c) Benzaldehyde (d) A polymeric product
20.  $[\text{Ag}(\text{NH}_3)_2]\text{OH}$  forms silver when it reacts with:
- (a)  $\text{HCOOH}$  (b)  $\text{CH}_3\text{COOH}$   
(c)  $\text{CH}_3\text{COCH}_3$  (d)  $\text{CH}_3\text{OH}$

## II FILL IN THE BLANKS

- Tollens' reagent is chemically.....
- Reaction of phthalic acid with ammonia followed by strong heating gives.....
- Acetyl chloride may be converted into acetaldehyde by .....reduction.
- Groups like nitro- and chloro- .....acidity of aromatic carboxylic acids.
- Phenol and benzoic acid can be distinguished by the reaction with.....
- On reaction of propanal with Fehling's reagent red precipitate of ..... is formed.
- Reaction of Grignard reagent with .....results into formation of primary alcohols.
- Lower aldehydes are soluble in water due to.....interactions with water molecules.
- DIBAL-H is used to reduce nitriles or esters to corresponding.....
- Reaction of ethyl magnesium bromide with carbon dioxide followed by acidification gives .....

### III ASSERTION REASON TYPE QUESTIONS

The question given below consist of an Assertion and Reason. Use the following key to choose the appropriate answer.

- (a) Assertion and reason both are correct and reason is the correct explanation of the assertion.
- (b) Assertion and reason both are correct statements but reason is not correct explanation of assertion.
- (c) Assertion is correct statement but reason is wrong statement.
- (d) Assertion is wrong statement but reason is correct statement.

1. **ASSERTION:** Reaction of Pentanoic acid with  $\text{Cl}_2$ / Red P followed by water gives 2-Chloropentanoic acid.  
**REASON:** The substitution takes places at  $\alpha$ -carbon.
2. **ASSERTION:** Carboxylic acids are higher boiling than aldehydes and ketones of comparable molar masses.  
**REASON:** Due to strong intermolecular H-bonding in carboxylic acids.
3. **ASSERTION:** Nitration of benzoic acid gives *m*-nitrobenzoic acid.  
**REASON:** Carboxyl group increases electron density on ring.
4. **ASSERTION:** Benzaldehyde undergoes Cannizzaro reaction.  
**REASON:** It contains one  $\alpha$ -hydrogen.
5. **ASSERTION :** Formaldehyde is a planar molecule.  
**REASON :** It contains  $\text{sp}^2$  hybridised carbon atom.
6. **ASSERTION :** Compounds containing -CHO group are easily oxidised to corresponding carboxylic acids.  
**REASON :** Carboxylic acids can be reduced to alcohols by treatment with  $\text{LiAlH}_4$ .
7. **ASSERTION :** The  $\alpha$ -hydrogen atom in carbonyl compounds is less acidic.  
**REASON :** The anion formed after the loss of  $\alpha$ -hydrogen atom is resonance stabilised.
8. **ASSERTION :** Aromatic aldehydes and formaldehyde undergo Cannizaro reaction.  
**REASON :** Aromatic aldehydes are almost as reactive as formaldehyde.
9. **ASSERTION :** Aldehydes and ketones, both react with Tollens' reagent to form silvery mirror.  
**REASON :** Both, aldehydes and ketones contain a carbonyl group.

10. **ASSERTION:** Ketones are oxidised under drastic conditions.

**REASON:** Oxidation of ketones gives carboxylic acids having carbons lesser than parent molecule.


#### IV ONE WORD TYPE QUESTIONS

- Name the product formed by addition of one equivalent of monohydric alcohol to aldehydes.
- Name the product formed by the reaction of benzene with CO, HCl in presence of anhyd.  $\text{AlCl}_3$
- Which reagent is used to convert carboxylic acid to corresponding alcohol?
- Which reaction is carried out to reduce the number of carbons from carboxylic acids?
- Which ester will be formed by the reaction of methanol and propanoic acid?
- Write the major product formed by the reaction of benzaldehyde and acetophenone.
- Which reagent will be best to convert ketone to corresponding alcohol in presence of carboxylic acid?
- Which reagent converts carboxylic acids into corresponding anhydrides?
- Name the carboxylic acid formed by reaction of cyclohexene with  $\text{KMnO}_4\text{-H}_2\text{SO}_4$  and heating.
- Which out of each pair is expected to be stronger acid ?  
(a)  $\text{CH}_3\text{COOH}$  or  $\text{HCOOH}$   
(b)  $\text{CH}_2(\text{Cl})\text{COOH}$  or  $\text{CH}_2(\text{Br})\text{COOH}$
- Name the test only to distinguish between Pentan-2-one and Pentan-3-one and Pentan-3-one.
- Predict the products when cyclohexanecarbaldehyde reacts with Zinc amalgam and HCl.
- Write the catalyst used in Rosenmund's reduction
- Name the reagent used in following reaction  

$$\text{CH}_3-\overset{\text{O}}{\underset{\text{||}}{\text{C}}}-\text{CH}_3 \xrightarrow{\quad ? \quad} \text{CH}_3-\overset{\text{OH}}{\underset{\text{CH}_3}{\underset{|}{\text{C}}}}-\text{CH}_3$$
- Out of  $\text{CH}_3\text{CHO}$  and  $\text{CH}_3\text{COCH}_3$  which one is more reactive towards HCN.


**VERY SHORT ANSWER TYPE QUESTIONS (1 Marks)**

1. Arrange the following compounds in increasing order of their acidic strengths:



Ans.  $(\text{CH}_3)_2\text{CHCOOH} < \text{CH}_3\text{CH}(\text{Br})\text{CH}_2\text{COOH} < \text{CH}_3\text{CH}_2\text{CH}(\text{Br})\text{COOH}$

2. Draw the structure of the compound whose IUPAC name is 4-chloropentan-2-one.




3. Which type of aldehyde can undergo Cannizzaro reaction?

Ans. Aromatic and aliphatic aldehydes which do not contain  $\alpha$ -hydrogen.

4. Name the aldehyde which does not give Fehling's test.

Ans. Benzaldehyde.

5. Arrange the following in order of their increasing reactivity towards HCN:




Ans.  $\text{C}_2\text{H}_5\text{COCH}_3 < \text{CH}_3\text{COCH}_3 < \text{CH}_3\text{CHO} < \text{HCHO}$

6. Arrange the following compounds in increasing order of their boiling point:



Ans.  $\text{CH}_3\text{CH}_2\text{CH}_3 < \text{CH}_3\text{OCH}_3 < \text{CH}_3\text{CHO} < \text{CH}_3\text{CH}_2\text{OH}$


8. How is acetone obtained from ethanol?



9. Why do aldehydes and ketones have lower boiling point than alcohols?

Ans. Due to presence of associated molecules with H-bonding in alcohols.

10. Write reaction between acetyl chloride and dimethyl cadmium.




11. What happens when  $\text{CH}_3\text{CHO}$  is treated with  $\text{K}_2\text{Cr}_2\text{O}_7$  in presence of  $\text{H}_2\text{SO}_4$ ?





12. Write IUPAC name of following compound:



Ans. 3,7-Dimethylocta-2,6-dien-1-al

13. Give balanced equation and name of products when  $\text{CH}_3\text{COOH}$  is treated with  $\text{PCl}_5$ ?

Ans.  $\text{CH}_3\text{COOH} + \text{PCl}_5 \rightarrow \text{CH}_3\text{COCl} + \text{POCl}_3 + \text{HCl}$


14. What product is obtained when ethyl benzene is oxidized with alkaline  $\text{KMnO}_4$ ?

Ans. Benzoic acid ( $\text{C}_6\text{H}_5\text{COOH}$ ) is obtained.

15.  $\text{CH}_3\text{CHO}$  is more reactive than  $\text{CH}_3\text{COCH}_3$  towards reaction with  $\text{HCN}$ . Give reason.


Ans. Due to  $-\text{CH}_3$  group, which shows +I effect and steric hindrance caused by two  $-\text{CH}_3$  groups.

16. Write IUPAC names of the following compound:



Ans. But-2-ene-1,4-dioic acid

17. Write the IUPAC name of following molecule:



Ans. Methyl-2-methylbenzoate.

18. Why does benzoic acid does not undergo Friedel-Crafts reaction?

Ans.  $-\text{COOH}$  group in  $\text{C}_6\text{H}_5\text{COOH}$  is an electron withdrawing group, which deactivates the benzene ring, hence electrophilic substitution becomes difficult.


19. Benzaldehyde gives a positive test with Tollens' reagent but not with Fehling's and Benedict solutions. Why?

Ans. It is due to stronger oxidising nature of Tollens' reagent as compared to Fehling's and Benedict's solution and cannot oxidise benzaldehyde to benzoic acid. In general, all these three can oxidise aliphatic aldehydes.

20. Write the chemical equation for Wolf Kishner Reduction.

Ans.  $\text{CH}_3\text{CHO} \xrightarrow{\text{NH}_2\text{NH}_2} \text{CH}_3\text{-C=N-NH}_2 \xrightarrow[\text{Ethylene glycol}]{\text{KOH}} \text{CH}_3\text{-CH}_3 + \text{N}_2$


21. Name alkene which on ozonolysis give acetone as a product



22. Give reason why hydrazones of RCHO and ketone are not produced in strongly acidic medium?

Ans. In acidic medium,  $\ddot{\text{N}}\text{H}_2\ddot{\text{N}}\text{H}_2$  get protonated and will not act as  $\text{Nu}^\ominus$ .

23. Complete the reaction



24. Name the aldehyde, which any exist in gaseous state.

Ans.  $\text{HCHO}$  (Formaldehyde)



25. Give reason:- During esterification between acid and alcohol, water or the ether should be removed as soon as it is formed.

Ans.  $\text{Acid} + \text{alcohol} \rightleftharpoons \text{ether} + \text{water}$





To make the reaction fast, by removing product as per Le-chatlier principle

**SHORT ANSWER TYPE QUESTIONS (2 or 3 Marks)**

1. How will you convert:




2. Complete the following:



3. An organic compound 'X' has molecular formula  $C_6H_{10}O$ . It does not reduce Fehling's solution but forms a bisulphite compound. It also gives positive Iodoform test. What are possible structure of 'X'? Explain.

Ans. 'X' gives positive test with Iodoform. It is Methyl ketone.



are possible structures of the compound.

4. Give the chemical test to distinguish between:



Ans. (i)  $CH_3CHO$  produce silver mirror with Tollens' reagent.




(ii)  $CH_3CHO$  gives brick red ppt.



Brick red ppt.

5. Is benzaldehyde more reactive or less reactive in nucleophilic addition reactions than propanal? Explain your answer.

Ans. Carbon atom of carbonyl in  $\text{C}_6\text{H}_5\text{CHO}$  is less reactive than that of propanal.  $\text{C}_6\text{H}_5\text{CHO}$  is less polar due to resonance.



6. Which acid of each pair shown here would you expect to be stronger?

(i)  $\text{CH}_3\text{CO}_2\text{H}$  or  $\text{CH}_2\text{FCO}_2\text{H}$

(ii)  $\text{CH}_2\text{FCO}_2\text{H}$  or  $\text{CH}_3\text{ClCO}_2\text{H}$

(iii)  $\text{CH}_2\text{FCH}_2\text{CH}_2\text{CO}_2\text{H}$  or  $\text{CH}_3\text{CHFCH}_2\text{CO}_2\text{H}$

(iv)  $\text{F}_3\text{C}-\text{C}_6\text{H}_4-\text{COOH}$  or  $\text{H}_3\text{C}-\text{C}_6\text{H}_4-\text{COOH}$

Ans. (i)  $\text{FCH}_2\text{COOH}$


(ii)  $\text{CH}_2\text{FCOOH}$

(iii)  $\begin{array}{c} \text{H} \\ | \\ \text{H}_3\text{C}-\text{C}-\text{CH}_2\text{COOH} \\ | \\ \text{F} \end{array}$



(iv)  $\text{CF}_3-\text{C}_6\text{H}_4-\text{COOH}$

7. Carboxylic acids do not give reactions of aldehydes and ketones why?

Ans. It is due to resonance



8. Write IUPAC name of the following :



Ans. (i) 3,5-dimethylphenylethanoate.

(ii) 5-Bromo-3-chloro-2-iodobenzoic acid.

9. Distinguish between :


(i)  $C_2H_5OH$  and  $CH_3CHO$

(ii)  $C_6H_5COCH_3$  and  $C_6H_5CH_2CHO$



Ans. (i)  $CH_3CHO$  gives silvery mirror with Tollens' reagent while  $C_2H_5OH$  does not.

(ii) Acetophenone will give yellow ppt. of iodoform while  $C_6H_5CH_2CHO$  will not.


10. Complete the following reactions by identifying A, B and C :



Ans.



11. Write the structures of A,B,C,D and E in the following reactions:



Ans. A =  $CH_3COCH_3$

B =  $C_6H_5CH_2CH_2$

C =  $C_6H_5COOH$

D =  $C_6H_5COONa$


E =  $CHI_3$

12. Aldehydes usually do not form stable hydrates but chloral normally exists as chloral hydrate. Give reason.

Ans. In case of aldehyde reaction is reversible.

$$\text{R}-\text{C}(=\text{O})-\text{H} + \text{H}_2\text{O} \rightleftharpoons \text{R}-\text{C}(\text{OH})_2-\text{H}$$


In case of  $\text{CCl}_3\text{CHO}$ , Cl atoms increase positive charge on carbonyl carbon. Therefore, weak nucleophiles like water readily added to the carbonyl group.



13. Give possible explanation for the following:

- Cyclohexanone forms cyanohydrins in good yield but 2,2,6-trimethylcyclohexanone does not.
- There are two  $-\text{NH}_2$  groups in semicarbazide. However, only one is involved in formation of semicarbazone.

Ans. (i) Due to steric hindrance for  $\text{CN}^-$  at  $\text{C}=\text{O}$  due to 3-methyl groups at  $\alpha$ -position.




- Only one  $-\text{NH}_2$  group attached to  $\text{C}=\text{O}$  is involved in resonance. As result electron density on these  $-\text{NH}_2$  group decreases and hence does not act as nucleophile.



14. Convert the following in not more than two steps:

- Benzoic acid to Benzaldehyde
- Propanone to Propene





15. Write the reactions involved in the following reactions:

(i) Clemmensen reduction

(ii) Cannizzoro reaction

Ans.





16. Convert the following





(i) Ethylbenzene to Benzoic acid





(ii) Ethanal to But-2-enal

Ans.





17. Predict the organic products of the following reactions:






18. Complete the following as missing starting material, reagent or products:







1,2-Dicyclohexylethene




Potassium benzoate

19. Mention the reactions involved for the following conversion.









20. Give reason for the following:

- Carboxylic acids do not give characteristic reactions of carbonyl groups.
- Treatment of  $\text{C}_6\text{H}_5\text{CHO}$  with  $\text{HCN}$  gives a mixture of two isomers which cannot be separated even by fractional distillation.
- Sodium bisulphite is used for purification of ketones and aldehydes.


Ans.



- Due to two optical isomers fractional distillation is not possible.



- Due to formation of addition compound of aldehydes and ketones with  $\text{NaHSO}_3$  whereas impurities do not.



21. Write tests to distinguish between:


- $\text{CH}_3\text{CHO}$  and  $\text{C}_6\text{H}_5\text{CHO}$
- $\text{C}_6\text{H}_5\text{-OH}$  and  $\text{CH}_3\text{COOH}$
- Pentanal and Pentan-2-one

- Ans. (i)  $\text{CH}_3\text{CHO}$  gives brick red ppt. with Fehling while  $\text{C}_6\text{H}_5\text{CHO}$  not.  
 (ii) Phenol does not give brisk effervescence but  $\text{CH}_3\text{COOH}$  gives this test with  $\text{NaHCO}_3$ .  
 (iii) Pentanal forms silver mirror but Pentan-2-one does not.

22. Convert the following:

- (i) Benzaldehyde to Acetophenone  
 (ii) Malonic acid to Acetic acid  
 (iii) Acetaldehyde to Butan-2-ol






Ans.



### LONG ANSWER TYPE QUESTIONS (5 Marks)

1. Write chemical reaction to affect the following transformations:
- Butan-1-ol to Butanoic acid
  - Benzyl alcohol to Phenylethanoic acid
  - 3-Nitrobromobenzene to 3-Nitrobenzoic acid
  - 4-Methylacetophenone to Terephthalic acid
  - Cyclohexane to Hexane-1,6 dioic acid
  - Butanal to Butanoic acid



Ans.







2. Draw the structure of the following derivatives:

- 2,4-Dinitrophenylhydrazone of  $\text{C}_6\text{H}_5\text{CHO}$
- Cyclopropanone oxime
- Acetaldehyde Dimethylacetal
- Semicarbazone to Cyclobutanone
- Ethylene ketal of Hexan-3-one
- Methylhemiacetal of formaldehyde


Ans.





3. An aromatic compound 'A' (Molecular formula  $C_8H_8O$ ) gives positive 2, 4-DNP test. It gives a yellow precipitate of compound 'B' on treatment with iodine and sodium hydroxide solution. Compound 'A' does not give Tollens' or Fehling's test. On drastic oxidation with potassium permanganate it forms a carboxylic acid 'C' (Molecular formula  $C_7H_6O_2$ ), which is also formed along with the yellow compound in the above reaction. Identify A, B and C and write all the reactions involved.

Ans:



4. Give reason for the following:

- $\text{C}_6\text{H}_5\text{COOH}$  is weaker than formic acid.
- $\text{HCOOH}$  and  $\text{CH}_3\text{COOH}$  differentiated by Tollens' reagent.
- $\text{R-COOH}$  do not give characteristic reaction with  $>\text{C}=\text{O}$ .
- Carboxylic acids are stronger acids than phenols.
- Acid amides are weakly basic in nature.

Ans. (i) In  $\text{C}_6\text{H}_5\text{COOH}$ ,  $\text{C}_6\text{H}_5$ -destablize carboxylate ion due to its +R effect.

(ii) Presence of -CHO group.


(iii)  $>\text{C}=\text{O}$  group is sterically hindered in carboxylic acid.

(iv) Dispersal of negative charge on carboxylate ions than phenoxide ion.


(v) Acid amides are basic due to lone pair of electrons on nitrogen atom.

5. An organic compound 'A' ( $\text{C}_3\text{H}_6\text{O}$ ) is resistant to oxidation but forms compound 'B' ( $\text{C}_3\text{H}_8\text{O}$ ) on reduction. 'B' reacts with  $\text{HBr}$  to form the compound 'C'. 'C' with  $\text{Mg}$  forms Grignard's reagent 'D' which reacts with 'A' to form a product which on hydrolysis gives 'E'. Identify 'A' to 'E'.









Ans. 'A' must be ketone.



6. A ketone [A] which undergoes haloform reaction gives compound [B] on reduction. [B] on heating with  $\text{H}_2\text{SO}_4$  gives compound [C] which forms monozonide [D] by reacting with ozone. [D] on hydrolysis in the presence of Zinc dust gives acetaldehyde. Identify [A], [B], [C], [D]. Write the reaction involved.




7. Complete the following reaction




8. An organic compound [A] with molecular formula  $C_5H_8O_2$  is reduced to n-pentane on treatment with Zn-mg/HCl. The compound [A] forms a dioxime with hydroxyle amine and gives a positive iodo form test and Tollen's test. Identify the compound [A] and deduce its structure.

Ans.



- (9). Complete the missing products



- Ans. (a)  $\text{A}=\text{CH}_3\text{OH}$ ,  $\text{B}=\text{CH}_3\text{Cl}$ ,  $\text{C}=\text{CH}_3\text{CN}$ ,  $\text{D}=\text{CH}_3\text{COOH}$   
 (b)  $\text{A}=\text{CH}_3\text{CHO}$ ,  $\text{B}=\text{CH}_3\text{COOH}$ ,  $\text{C}=\text{CH}_3\text{COCl}$   
 (c)  $\text{A}=\text{CH}_3\text{CH}_2\text{-Br}$ ,  $\text{B}=\text{CH}_3\text{CH}_2\text{-CN}$ ,  $\text{C}=\text{CH}_3\text{CH}_2\text{COOH}$ ,  
 $\text{D}=\text{CH}_3\text{CH}_2\text{CH}_2\text{OH}$

10. [A], [B] and [C] are three non-cyclic functional isomer of a carbonyl compound with molecular formula  $C_4H_8O$ . Isomer [A] and [C] gives positive Tollen's test whereas Isomer [B] does not give Tollen's test but give positive iodoform test. Isomer [A], [B] on reduction with  $\text{Zn}[\text{Hg}]/\text{conc. HCl}$  give same product [D]. Identify A, B, C, D.

- Ans.  $\text{A}=\text{CH}_3\text{CH}_2\text{CH}_2\text{CHO}$  Butan-1-al  
 $\text{B}=\text{CH}-\overset{\overset{\text{O}}{\parallel}}{\text{C}}-\text{CH}_2-\text{CHO}$  Butan-2-one  
 $\text{C}=\text{CH}_3-\underset{\underset{\text{CH}_3}{|}}{\text{C}}-\text{CHO}$  2-Methylpropanal  
 $\text{D}=\text{CH}_3-\text{CH}_2-\text{CH}_2-\text{CH}_3$  Butane

**CASE-STUDY BASED QUESTIONS****1. Read the passage and answer the following questions:**

Carboxylic acids are compounds with excellent chemical and physical properties, the most particular characteristics of this type of organic compounds, is their high solubility in polar solvents, as water, or alcohols, methanol, ethanol, etc. Chemical structure contains a carbonyl function ( $\text{-C=O}$ ) and an hydroxyl group ( $\text{OH}$ ), these groups interact easily with polar compounds, forming bridges of H, obtaining high boiling points. The carbonyl group ( $\text{C=O}$ ) is considered a one of the most functional groups involved in many important reactions. The carboxylic acids are the most important functional group that present  $\text{C=O}$ .


This type of organic compounds can be obtained by different routes, some carboxylic acids, such as citric acid, lactic acid or fumaric acid are produced from by fermentation, most of these type of carboxylic acids are applied in the food industry. Historically, some carboxylic acids were produced by sugar fermentation. Synthetics route, there are different synthesis reactions such as reactions of oxidation from alcohols in the presence of strong oxidants such as  $\text{KMnO}_4$ , oxidation of aromatic compounds among other routes. For example, citric acid is a carboxylic acid, can be obtained by different routes, synthetic, enzymatic and naturally occurring, is considered harmless and cheap, used in the food industry, because is non-toxic, has a thermal stability to the  $175^\circ\text{C}$ . Bian et al., in 2017, reported the use of citric acid impregnated in porous material for the synthesis of Ni particles. They showed, that the presence of citric acid, is important in the dispersion of the Ni particles when are incorporate in porous materials, thus inhibiting the agglomeration.

Derivatives of carboxylic acid, as alkyl halides, esters, and amides, present different and important application in diverse areas. In the case of esters, these are obtained from the reaction between carboxylic acids and alcohols in presence of an acid catalyst usually  $\text{H}_2\text{SO}_4$  with heat, this type of reaction is known as esterification. In the case of the amides, it is obtained in the presence of an amine, may be primary and secondary, with a carboxylic acid, in this reaction also can be used a catalyst and heat to accelerate the reaction.



**Reference :** Aide Sienz-Galindo, Lluvia I. Lopez-Lopez, Fabiola N. de la Cruz- Duran, Adali O.Castafieda-Facio, Leticia A. Ramirez-Mendoza, Karla C. Cordova-Cisneros and Denisse de Loera-Carrera (March 15th 2018). **Applications of Carboxylic Acids in Organic Synthesis, Nanotechnology and Polymers, Carboxylic Acid-Key Role in Life Sciences**, Georgiana Ileana Badea and Gabriel Lucian Radu, *IntechOpen*, DOI: 10.5772/intechopen.74654.

(A) Identify A and B in following reaction:



(B) Assertion: Carboxylic acids are highly acidic.

Reason: Carboxylate ion is resonance stabilised.

- (a) Assertion and reason both are correct and reason is the correct explanation of the assertion.
  - (b) Assertion and reason both are correct statements but reason is not correct explanation of assertion.
  - (c) Assertion is correct statement but reason is wrong statement.
  - (d) Assertion is wrong statement but reason is correct statement.
- (C) What happens when phthalic acid reacts with ammonia followed by strong heating?
- (D) How acetyl chloride may be converted to ethanoic acid?

**2. Read the passage and answer the following questions:**

'Tetrahydrofuran (THF) is a bulk chemical, which can be obtained from various feedstocks including biomass. In addition, the C5 carboxylic acids are much more expensive than THF. Therefore, we adopted THF as model ether to study the catalytic system (Table 1). The reaction could be efficiently accelerated by  $\text{IrI}_4$  catalyst and LiI promoter in AcOH solvent at 170 °C, and the yield of C5 carboxylic acids reached 70% after 16 h (entry1). The products contained two isomers, i.e., pentanoic acid and 2-methylbutanoic acid, and their molar ratio was 58:42. A little C6 carboxylic acids were also formed in the reaction. The rest of the THF substrate was converted to butane.

In addition, trace of methane was also detected. We also tried different Ir catalyst precursors, such as  $\text{Ir}(\text{CO})(\text{PPh}_3)_2\text{Cl}$ ,  $\text{Ir}(\text{CO})_2(\text{acac})$ , and  $\text{IrCl}_3$ , the results indicated that they were not as efficient as  $\text{IrI}_4$ . We set the reaction time at 8 h and tested other catalytic systems. The  $\text{IrI}_4$  catalyst was essential to the reaction because no target product was observed without it. The Rh catalyst was effective for synthesis of carboxylic acids via olefin and/or alcohol hydrocarboxylation with  $\text{CO}_2$  and  $\text{H}_2$ . Whereas in this work no product was obtained when  $\text{RhI}_3$



(A) How ethers can be distinguished from carboxylic acid?


(B) Propanol reaction with.....and.....respectively gives.

3. **Read the passage and answer the following questions:**


The acetal is the most common protecting group for aldehydes and 1,3-dioxolanes are the most commonly encountered type of acetal, usually prepared by reaction of the aldehyde with ethylene glycol with azeotropic removal of water (eq1). Regeneration of the carbonyl is normally out with aqueous acid.



We have been concerned with the general problem of converting dioxolanes into Carboxylic acids without employing acid to first remove the protective group (eq 2). The non-acidic alternative to eq 2 would allow the introduction of acid groups into a molecule containing.



various acid-sensitive functionalities. Our solution to this problem is outlined in eq 3. Prugh and McCarthy in 1966 showed that cyclic acetals are converted into bromo esters when treated with N-bromosuccinimide (NBS). Indeed, a



variety of dioxolanes give good yields of the corresponding 2-bromoethyl esters when refluxed with NBS in  $\text{CCl}_4$ . For example, 3-phenyl-1,3-dioxolane gives a 98% yield of 2-bromoethyl benzoate (88% after distillation).

The transformation of eq 3 is completed by a zinc-induced elimination which yields the acid upon workup. Despite the precedent for this second step, a variety of reaction conditions failed to give any acid from 2-bromoethyl benzoate.

Zinc in refluxing THF gave no reaction. Even zinc which had activated with copper sulfate was ineffective and ultraactive zinc from the potassium metal or sodium naphthalenide reduction of zinc chloride also failed to promote elimination. Zinc in refluxing methanol or ethanol gives 42-46% benzoic acid plus 47-52% of transesterification product. Ester interchange can be avoided by using zinc in refluxing THF to give a 44% yield of benzoic acid and a recovery of starting material. Addition of catalytic sodium iodide improves the yield of benzoic acid from this reaction to with only of starting material recovered.

**Reference :** Lawrence C. Anderson, Harold W. Pinnick, **Preparation of carboxylic acids from protected aldehydes** *J. Org. Chem.* **1978**, 43, 17, 3417-3418  
<https://doi.org/10.1021/io00411a044>

- (A) The formation of acetal from aldehyde is an example of ..... reaction.
- (B) Which of the following reagent(s) can be used to convert butan-1-ol to butanoic acid?
  - (a) 1.  $\text{KMnO}_4$  —  $\text{KOH}$  2.  $\text{H}^+$
  - (b)  $\text{CrO}_3$  —  $\text{H}_2\text{SO}_4$
  - (c) Both i) and ii)
  - (d) None of these
- (C) Write the structure of the product formed when propanal reacts with methanol.
- (D) What happens when propanal reacts with ethyl magnesium iodide followed by reaction with dilute acid?

4. **Read the passage and answer the following questions:**

Nucleophilic additions to a carbonyl group leading to tetrahedral species which are products or intermediates in a mechanistic sequence occupy a central place in biochemistry as well as in organic chemistry. It has been shown recently that the structural pathway for the nucleophilic addition of an amino group to carbonyl can be mapped on the basis of crystal structure data. However, from structural data alone no direct information about the energy variation along the reaction pathway can be obtained. Furthermore, it was not entirely clear to what extent steric requirements of substituents on the nucleophile and the carbonyl group, as well as crystal packing effects, influence the arrangement of the reactive centers. An attempt to fill these gaps we have carried out calculations on the reaction path of the simple model system.



corresponding to nucleophilic addition of hydride anion to formaldehyde to produce methanolate anion. A calculation has also been made for the system consisting of an ammonia and a formaldehyde molecule at a distance of 20 nm.


**Reference :** H. B. Bürgi, J. M. Lehn, G. Wipf, **Ab initio study of nucleophilic addition to a carbonyl group**, *J. Am. Chem. Soc.* 1974, 96, 6, 1956-1957, Publication Date: March 1, 1974 <https://doi.org/10.1021/ja00813a062>

(A) Which of the following nucleophilic reaction generates chiral carbon?

- (a) Benzaldehyde + KOH
- (b) Benzaldehyde + ammonia
- (c) Propanone + KOH
- (d) Propanone + ammonia

(B) Write the major product on reaction of acetophenone with  $\text{CH}_3\text{-NH}_2$ ?

(C) Complete the reaction:



(D) Which will undergo reaction with 2,4-dinitrophenylhydrazine at fastest rate?

- (a) Acetophenone
- (b) Propanone
- (c) Benzaldehyde
- (d) Propanal

## ANSWERS

### I MULTIPLE CHOICE QUESTIONS

1. (b) 2. (d) 3. (b). 4. (b) 5. (b) 6. (a) 7. (d) 8. (a) 9. (d) 10. (d) 11. (c) 12. (c)  
13. (a) 14. (b). 15. (a) 16. (b) 17. (c) 18. (a) 19. (b) 20. (a)

### II FILL IN THE BLANKS

1. Ammoniacal silver nitrate 2. Phthalimide 3. Rosenmund  
4. Increases 5.  $\text{NaHCO}_3$  6.  $\text{Cu}_2\text{O}$   
7. Methanal 8. H-bonding. 9. Aldehydes  
10. Propanoic acid

### III ASSERTION REASON TYPE QUESTIONS

- III. 1. (a) 2. (a) 3. (c) 4. (c) 5. (a) 6. (b) 7. (d) 8. (c) 9. (d) 10. (b)

### IV ONE WORD ANSWER QUESTIONS


1. Hemiacetal 2. Benzaldehyde 3.  $\text{LiAlH}_4$  or  $\text{B}_2\text{H}_6$   
4. Decarboxylation 5. Methyl propanoate  
6. 1,3-Diphenylprop-2-en-1-one or Benzalacetophenone  
7.  $\text{NaBH}_4$  8.  $\text{P}_2\text{O}_5$  or  $\text{H}^+/\Delta$  9. Hexane-1,6-dioic acid  
10. a)  $\text{HCOOH}$ , (b)  $\text{CH}_2(\text{Cl})\text{COOH}$  11. Iodoform Test  
12. Methycyclohexane 13.  $\text{Pd/BaSO}_4$  14.  $\text{CH}_3\text{MgBr/H}_2\text{O/H}^+$   
15.  $\text{CH}_3\text{CHO}$

### CASE STUDY TYPE QUESTIONS

#### PASSAGE : 1

(A)  $\text{X} = \text{C}_6\text{H}_5\text{COCl}$   $\text{Y} = (\text{C}_6\text{H}_5\text{CO})_2\text{O}$

(B) (a)



#### PASSAGE : 2

- (A) With  $\text{NaHCO}_3$  carboxylic acids give brisk effervescence, whereas ethers cannot.  
(B) PCC/PdC

**PASSAGE :3**

- (A) Nucleophilic addition reaction  
 (B) (c)  
 (C)  $\text{CH}_3\text{CH}_2\text{CH}(\text{OCH}_3)_2$   
 (D)  $(\text{C}_2\text{H}_5)_2\text{CHOH}$

**PASSAGE :4**

- (A) (c)  
 (B)  $\text{C}_6\text{H}_5-\underset{\text{CH}_3}{\text{C}}=\text{N}-\text{CH}_3$   
 (C)  $\text{CH}_3 > \underset{\text{CH}_3}{\text{C}} \begin{matrix} \text{O}-\text{CH}_2 \\ \text{O}-\text{CH}_2 \end{matrix}$   
 (D) (d)

## UNIT TEST-1

## CHAPTER-11

Maximum Marks : 20

Time : 1 Hrs.

1. Identify the correct product in following reaction:



- (a) CH3CH=CHCHO (b) CH3CH=CHCOOH  
 (c) HOCH2CH=CHCH2OH (d) CH3CH2CH2CHO

2. Which of the following reagent(s) is most suitable for following conversion?

Propanone  $\rightarrow$  Propane

- (a) PCC (b)  $\text{LiAlH}_4$   
 (c) (i).  $\text{KMnO}_4$  -KOH (ii).  $\text{H}^+$  (d)  $\text{Zn(Hg)/HCl}$   
 3. The highest  $\text{pK}_a$  value is observed in-  
 (a) Phenol (b) Benzoic acid  
 (c) 4-Nitrobenzoic acid (d) Ethanoic acid  
 4. Which of the following undergoes nucleophilic addition reaction at fastest rate?  
 (a) Benzaldehyde (b) Acetophenone  
 (c) Methanal (d) Ethanal  
 5. Cannizzaro reaction is not shown by-

- (a) CC1(C)CCCCC1=O (b) c1ccccc1C=O  
 (c) HCHO (d) CC=O


6. Give a chemical test to distinguish between following compounds:

- (a) Benzaldehyde and Acetophenone  
 (b) Benzoic acid and Benzophenone

7. How butanoic acid can be synthesised using appropriate:

- (a) Grignard reagent  
 (b) Amide

8. Arrange following in ascending order of given properties:
- Ethanal, Ethanol, Methoxymethane, Propane (boiling point)
  - Propanal, Benzaldehyde, Acetophenone (reactivity towards nucleophilic addition reaction)
9. Complete the reaction sequence by writing structures of A-E. Also name the reaction involved in the conversion of A to B.




10. Explain following:
- $\alpha$ -hydrogens in aldehydes and ketones are acidic in nature.
  - There are two  $\text{-NH}_2$  groups in semicarbazide. However only one involves in the formation of semicarbazone.
  - Propanone is less reactive than propanal towards nucleophilic addition reactions.
11. How will you carry out following conversions?
- Bromobenzene to 1-Phenylethanol
  - Benzoic acid to *m*-Nitrobenzyl alcohol
  - Propanone to Propene




**UNIT TEST- 2**  
**CHAPTER-11**

**Maximum Marks : 20****Time : 1 Hrs.**

1. Give the IUPAC of



2. Which acid will be more acidic




3. Write product of  $\text{CH}_3\text{CHO}$  with hydrazine.
4. Which will have higher boiling point  $\text{CH}_3\text{CHO}$  or  $\text{CH}_3\text{COOH}$ .
5. How many mole of hydrazine will be used with one mole of Phthaldehyde
6. Write chemical distinguish test to separate following.
- (a) Pentan-2-one and Pentan-3-one
  - (b) Benzophenone and Benzoic acid
7. Write short not on
- (a) Hell-Volhard-Zelinsky Reaction
  - (b) Etard Reaction
8. How will you synthesise Acetone from propene.
9. Arrange the following compound in increasing order of their properties.
- (a)  $\text{CH}_3\text{COOH}$ ,  $\text{C}_6\text{H}_5\text{COCH}_3$ ,  $\text{CH}_3\text{CHO}$  [Reactivity towards No addition]
  - (b)  $\text{Cl-CH}_2\text{-COOH}$ ,  $\text{F-CH}_2\text{-COOH}$ ,  $\text{CH}_3\text{-CH}_2\text{-COOH}$  (acidic character)
10. Give reason.
- (a) Cyclohexanone forms cyanohydrin in good yield but 2,2,6-trimethylcyclohexanone does not.
  - (b) There are two- $\text{NH}_2$  group in semicarbazide howere only one is involved in the formation of semicarbazone.
11. (a) Convert the following
- (i) Benzaldehyde to 3-Phenylpropan-1-ol
  - (ii) Benzole acid to m-Nitrobenzyl alcohol
- (b) An organic compound with molecular formula  $\text{C}_5\text{H}_{10}\text{O}$  from 2,4-DNP derivatives, reduce Tollen's reagent and undergoes cannizarro's reaction. on vigrous oxidation it gives 1,2- Benzene-dicarboxylic acid. Identify the compound.

## Points to Remember

## Geometry

## Pyramidal shape




## Nomenclature






**Common Names:-** Amine is used as suffix after alkyl group e.g.  $\text{CH}_3\text{CH}_2\text{NH}_2$  is ethylamine

**IUPAC names:-** e is replaced by - amine e.g.  $\text{CH}_3\text{CH}_2\text{NH}_2$  is ethanamine.

## Classification and Nomenclature of Amines

## Classification










**Method of Preparation****1. Reduction of Nitro compounds****2. Ammonolysis Alkyl Halides****3. Reduction of nitriles****4. Reduction of amide****5. Gabriel phthalimide synthesis****6. Hoffmann bromamide degeneration reaction.**

**Reactions****(i) Basic character amines**

.React with acid to form salts  $\text{R-NH}_2 + \text{HX} \rightleftharpoons \text{R-NH}_3^+ \text{X}^-$  (salt)

.React with base to regenerate parent amines

**(iii) Carbylamine reaction:****(iv) With nitrous acid****(v) With Benzene sulfonyl chloride****(vi) Electrophilic Substitution**

### OBJECTIVE TYPE QUESTIONS

#### I MULTIPLE CHOICE QUESTIONS



1. In the nitration of benzene using conc.  $\text{H}_2\text{SO}_4$  and conc.  $\text{HNO}_3$  the species which initiates the reaction is:

(a)  $\text{NO}^+$  (b)  $\text{NO}_2^+$   
(c)  $\text{NO}_2^-$  (d)  $\text{NO}_3^-$




2. The correct IUPAC name of  $\text{CH}_2=\text{CH}-\text{CH}_2\text{NHCH}_3$

(a) Allylmethylamine (b) 2-Aminopent-4-ene  
(c) 4-Aminopent-1-ene (d) N-Methylprop-2-enamine

3. Which is the weakest base?





4. The correct order of basic strength for the following compound is:



(a)  $\text{ii} < \text{iii} < \text{i}$  (b)  $\text{iii} < \text{i} < \text{ii}$   
(c)  $\text{iii} < \text{ii} < \text{i}$  (d)  $\text{ii} < \text{i} < \text{iii}$





5. The structure of 'C' in following reaction sequence would be -










6. Which of the following statement about primary amine is false?
- Aryl amines react with nitrous acid to produce phenol
  - Alkyl amines are stronger base than ammonia
  - Alkyl amines are stronger base than aryl amines
  - Alkyl amines react with nitrous acid to produce alcohol
7. Which of the following is most stable diazonium salt?
- $\text{CH}_3\text{N}_2^+\text{X}^-$
  - $\text{C}_6\text{H}_5\text{N}_2^+\text{X}^-$
  - $\text{CH}_3\text{CH}_2\text{N}_2^+\text{X}^-$
  - $\text{C}_6\text{H}_5\text{CH}_2\text{N}_2^+\text{X}^-$
8. Method by which aniline can not be prepared is:
- Reduction of nitrobenzene with  $\text{H}_2/\text{Pd}$  in ethanol.
  - Potassium salt of phthalimide treated with chlorobenzene
  - Hydrolysis of phenyl isocyanide with acidic solution
  - Degradation of benzamide with bromine in alkaline medium solution.
9. In the chemical reaction:
- $$\text{CH}_3\text{CH}_2\text{NH}_2 + \text{CHCl}_3 + 3\text{KOH} \rightarrow \text{'A'} + \text{'B'} + 3\text{H}_2\text{O}$$
- The compound 'A' and 'B' are respectively:
- $\text{CH}_3\text{CH}_2\text{CONH}_2$  and  $3\text{KCl}$
  - $\text{CH}_3\text{CH}_2\text{NC}$  and  $\text{K}_2\text{CO}_3$
  - $\text{CH}_3\text{CH}_2\text{NC}$  and  $3\text{KCl}$
  - $\text{CH}_3\text{CH}_2\text{CN}$  and  $3\text{KCl}$


10. An amide (A) reacts with bromine in aqueous NaOH and forms amine containing three carbons. Identify (A):
- (a) 2-Methylpropanamide                      (b) Propanamine  
(c) Butanamide                                  (d) None of these
11. Which of the following compound will give significant amount of *meta* product during mononitration reaction?




12. The final product C in the following sequence of reaction is:



13. In the reaction, the structure of product A is:



14. Which of the following reactions from benzylamine:



15. Bromobenzene can be prepared from benzene diazonium chloride by its treatment with-

- (a) Cu/HBr
- (b) Br<sub>2</sub>, HBr
- (c) CuBr/HBr
- (d) Br<sub>2</sub>/CCl<sub>4</sub>

16. Acetamide and Ethylamine can be distinguish by reacting with

- (a) aq. HCl and heat
- (b) aq. NaOH and heat
- (c) Acidified KMnO<sub>4</sub>
- (d) Bromine Water



17. The order of reactivity of halides with amines is
  - (a)  $RI > RBr > RCl$
  - (b)  $RBr > RI > RCl$
  - (c)  $RCl > RBr > RI$
  - (d)  $RBr > RCl > RI$
18. Which of the following does not affect the  $K_b$  of an organic base?
  - (a) +I or -I effect
  - (b) solvation effect
  - (c) density
  - (d) Steric hinderance
19. Which of the following on reduction with  $LiAlH_4$  yields a secondary amine?
  - (a) Methyl isocyanide
  - (b) Acetamide
  - (c) Methyl cyanide
  - (d) Nitroethane
20. In diazatisation of aniline with  $NaNO_2$  and  $HCl$  acid the excess of acid is used primarily to
  - (a) Suppress the concentration of free aniline
  - (b) suppress the hydrolysis to phenol
  - (c) ensure a stoichiometric amount of  $HNO_2$  acid
  - (d) neutralise the base liberated.

## II FILL IN THE BLANKS

1. Reaction of nitrobenzene with  $Fe + HCl$  results into the formation of.....
2. Aromatic amines are.....bases while aliphatic amines are.....bases than ammonia.
3. Gabriel phthalimide synthesis is used for synthesis of .....amines.
4. Benzenesulphonyl chloride is also known as.....reagent.
5. Butanamide on reaction with  $LiAlH_4$  forms.....
6. To reduce activation of aniline it is deactivated by.....reaction.
7. Nitration of aniline forms *para* and ..... isomers of nitroaniline as major products.
8. Benzene diazonium chloride may be converted into phenol by reaction with .....at 283 K.
9. Primary amines are soluble in water due to.....
10. Aryl ..... and ..... can not be prepared by direct halogenation.

### III ASSERTION REASON TYPE QUESTIONS

In the following questions a statement of assertion followed by a statement of reason is given. Choose the correct answer out of the following choices.


- (a) Both assertion and reason are correct statements and reason is correct explanation of assertion.
  - (b) Both assertion and reason are correct statements but reason is not correct explanation of assertion.
  - (c) Assertion is correct statement but reason is wrong statement.
  - (d) Assertion is wrong statement but reason is correct statement.
1. **ASSERTION :** Acylation of amines gives a monosubstituted product whereas alkylation of amines gives polysubstituted product.  
**REASON :** Acyl group sterically hinders the approach of further acyl groups.
  2. **ASSERTION :** Hoffmann's bromamide reaction results into formation of primary amines.  
**REASON :** Primary amines are more basic than secondary amines.
  3. **ASSERTION :** N-Ethylbenzene sulphonamide is soluble in alkali.  
**REASON :** Hydrogen attached to nitrogen in sulphonamide is strongly acidic.
  4. **ASSERTION :** N, N-Diethylbenzene sulphonamide is insoluble in alkali.  
**REASON :** Sulphonyl group attached to nitrogen atom is strong electron withdrawing group.
  5. **ASSERTION :** Only a small amount of HCl is required in the reduction of nitro compounds with iron scrap and HCl in the presence of steam.  
**REASON :** FeCl<sub>2</sub> formed gets hydrolysed to release HCl during the reaction.
  6. **ASSERTION :** Aromatic 1° amines can not be prepared by Gabriel phthalimide synthesis.  
**REASON :** Aryl halides undergo nucleophilic substitution with anion formed by phthalimide.
  7. **ASSERTION :** Acetanilide is less basic than aniline.  
**REASON :** Acetylation of aniline results in decrease of electron density on nitrogen.
  8. **ASSERTION:** n-Propylamine has higher boiling point than Trimethylamine.  
**REASON:** Among n-Propylamine molecules, there is hydrogen bonding but there is not hydrogen bonding in trimethylamine.
  9. **ASSERTION:** Aniline does not undergo Friedel Crafts reaction.  
**REASON:** Friedel Crafts reaction is an electrophilic substitution reaction.
  10. **ASSERTION:** Ethylamine is more basic than aniline.  
**REASON:** Due to +I effect of ethyl group electron density on nitrogen increases.

**IV ONE WORD TYPE QUESTIONS**

1. Name the reaction in which primary amines reacts with  $\text{CHCl}_3$  and  $\text{KOH}$  forming foul smelling substance.
2. Write the IUPAC name of white precipitate formed by reaction of aniline with bromine water.
3. Write the product formed by the Hoffmann bromamide degradation of benzamide.
4. Is  $\text{CH}_3\text{CONH}_2$  weaker/stronger base than  $\text{CH}_3\text{CH}_2\text{NH}_2$ ?
5. Which type of electrophilic substitution reaction is not possible with aniline?
6. Name the product formed when benzene diazonium chloride reacts with  $\text{H}_3\text{PO}_2$ .
7. Name the reaction which is used to convert diazonium salt into corresponding halide by reacting with  $\text{Cu(I)}$  halides.
8. Which type of reaction involves in the cleavage of  $\text{C-X}$  bond in ammonolysis?
9. What is the pH during coupling reaction between phenol and benzenediazonium salt.
10. Write the reagent which can be used to convert nitrobenzene to aniline.
11. How many structural isomers are possible for  $\text{C}_4\text{H}_{11}\text{N}$  compound.
12. Name the reagent for the conversion of amide to amine containing same number of carbon atoms.
13. Name a synthetic amino compound which is used as an anesthetic in dentistry.
14. What kind of substitution is involved in Gabriel phthalimide synthesis.
15. Name the solvent which we used in acetylation of aniline using acetic anhydride.

**VERY SHORT ANSWER TYPE QUESTIONS (1 Marks)**

1. Convert *m*-dinitrobenzene to *m*-nitroaniline.



2. Write IUPAC name of  $\text{CH}_3 - \underset{\text{CH}_3}{\text{N}} - \overset{\text{CH}_2 - \text{CH}_3}{\underset{\text{CH}_3}{\text{C}}} - \text{CH}_2 - \text{CH}_3$

Ans. 3-Methyl-N,N-dimethylpentanamine

3. Give one use of quaternary ammonium salts.

Ans. It is used as detergents, e.g.,  $[\text{CH}_3(\text{CH}_2)_{15}\text{N}(\text{CH}_3)_2]^+\text{Cl}^-$

4. Mention the chemical formula of Hinsberg's reagent.

Ans. Benzene sulphonyl chloride,  $\text{C}_6\text{H}_5\text{-SO}_2\text{Cl}$

5. Why aniline dissolves in HCl?

Ans.  $\text{C}_6\text{H}_5\text{NH}_2 + \text{HCl} \rightarrow [\text{C}_6\text{H}_5\text{NH}_3]^+\text{Cl}^-$

It dissolves due to its basic nature.

6. How will you test the presence of primary amine?

Ans. By carbylamine test.



7. What is vapour phase nitration?


Ans.  $\text{CH}_4 + \text{HNO}_3 \rightarrow \text{CH}_3\text{NO}_2$

(High temperature and nitration in vapour phase only)

8. Direct nitration of aniline is not carried out. Explain.


Ans. ( $\text{H}_2\text{SO}_4 + \text{HNO}_3$ ) easily oxidizes aniline into tarry complex product due to high electron-density on the benzene ring of aniline.

9. What happens when aniline is treated with bromine?




10. Write a chemical equation to illustrate the ammonolysis.

Ans. For alcohols:




For alkyl halides:




11. Prepare/convert nitrobenzene into aniline.

Ans.




12. Convert  $\text{C}_6\text{H}_5\text{COOH}$  to  $\text{C}_6\text{H}_5\text{NH}_2$



13. Write isomerism exhibited by different amines.

Ans. Chain, position, metamerism, functional.

14. Arrange the following compounds in increasing order of solubility in water:



15. What is the role of  $\text{HNO}_3$  in the nitrating mixture used for nitration of benzene.

Ans.  $\text{HNO}_3$  provides  $\text{NO}_2^+$  electrophile.


16. Why  $\text{C}_6\text{H}_5\text{N}_2\text{Cl}$  not stored and is used immediately after its preparation?

Ans. It is highly unstable

17. What is the best reagent to convert  $\text{C}_6\text{H}_5\text{CONH}_2$  into aniline?


Ans.  $\text{Br}_2 + \text{NaOH}$

18. Write of IUPAC of



Ans. 2,4-Dinitro Chlorobenzene

19. Write the structure of N-Methyl-N-ethylethanamine.



20. Rearrange the following compound in an increasing order of their basic strength.

Aniline, p-nitroaniline, P- toluidine.

Ans. p-Nitroaniline < Aniline < p-Toluidine

21. Ammonolysis of alkyl halide does not give amine in pure state why.

Ans. Because 2° and 3°, amines are also formed

22. A poisonous gas is formed by the reaction of  $\text{R-NH}_2$  with  $\text{CHCl}_3$ , KOH. Give the name of the test.

Ans. Carbylamine test

23. Out of Pentan-1-ol and pent-1-amine which is more soluble.

Ans. pentan-1-ol

24. Identify A and B



Ans. A-  $\text{CH}_3\text{-CONH}_2$       B-  $\text{CH}_3\text{-NH}_2$

25. Write the name of test to distinguish between aniline and N-methyl aniline.



Ans. Carbylamine test

### SHORT ANSWER TYPE QUESTIONS (2 or 3 Marks)

1. How will you convert following :

(a) 3-Methylaniline to 3-Nitrotoluene


(b) Aniline to 1,3,5-Tribromobenzene



2. A compound 'A' having molecular formula  $C_3H_7ON$  reacts with  $Br_2$  in presence of  $NaOH$  to give compound 'B'. This compound 'B' reacts with  $HNO_2$  to form alcohol and  $N_2$  gas. Identify compound 'A' and 'B' and write the reactions involved.




3. Account for following:
- Amino group in aniline is *o*- and *p*- directing in aromatic electrophilic substitution reactions but aniline on nitration gives a substantial amount of *m*-nitroaniline.
  - Aniline does not undergoes Friedel Crafts reaction.
- Ans. (i) It is because aniline is protonated to form anilinium cation, in which  $-NH_2$  group is *meta*-directing.
- (ii) It is because aniline is basic lewis can form adduct with  $AlCl_3$  which deactivates the ring.
4. How will you synthesise ethanamine by Gabriel phthalimide synthesis?



5. Write short notes on following:
- Coupling reaction
  - Ammonolysis






6. Account for the following :

- (b) (a) Electrophilic substitution in aromatic amines takes place more readily than benzene.  
 (b) Nitro compounds have higher boiling points than hydrocarbons having almost same molecular mass.


Ans. (a) -  $\text{NH}_2$  is electron releasing group so electrophilic substitution takes place faster.  
 (b) Nitro compounds are more polar than hydrocarbons therefore have more van der Waals forces of attraction.

7. Write the structure of reagents/organic compounds 'A' to 'E' :




8. What happens when :

- (a) An alkyl halide reacts with  $\text{AgNO}_2$  and product is reduced.  
 (b) An alkyl halide is treated with  $\text{AgCN}$  and product is hydrolysed.  
 (c) Methyl magnesium bromide is treated with cyanogen chloride.






9. Write reaction for benzoylation of aniline.



10. Draw structure of the following compounds:

- (a) N-Isopropylaniline  
(b) t-Butylamine




11. Give reasons:

- (a) Electrophilic substitution in aromatic amines takes place more readily than benzene  
(b)  $\text{CH}_3\text{CONH}_2$  is weaker base than  $\text{CH}_3\text{CH}_2\text{NH}_2$ .


Ans: (a)  $-\text{NH}_2$  group in aromatic amines shows +R effect and increases electron density on ring and facilitating the attack of electrophile.

- (b) In ethanamide the electron pair on nitrogen is in resonance with carbonyl group so electrons are less readily available than ethanamide.


12. An organic aromatic compound 'A' with the molecular formula  $\text{C}_6\text{H}_7\text{N}$  is sparingly soluble in water. 'A' on treatment with dil. HCl gives a water soluble compound 'B'. 'A' also reacts with chloroform in presence of alcoholic KOH to form an obnoxious smelling compound 'C'. 'A' reacts with benzene sulphonyl chloride to form alkali soluble compound 'D'. 'A' reacts with  $\text{NaNO}_2$  and HCl to form a compound 'E' which on reaction with phenol forms an orange dye. Elucidate the structures of the organic compounds from 'A' to 'F'.




A




B




C



D



E




F

13 Write chemical distinguish test between following pair of compound.

(i) Aniline and N-Methylaniline

(ii)  $(\text{CH}_3)_2\text{NH}$  and  $(\text{CH}_3)_3\text{N}$

Ans. (i) Aniline +  $\text{CHCl}_3 + \text{KOH} \rightarrow$  Foul smell



N-Methyl aniline +  $\text{CHCl}_3 + \text{KOH} \rightarrow$  No reaction

(ii)  $(\text{CH}_3)_2\text{NH} + \text{HNO}_2$  Yellow oily compound

$(\text{CH}_3)_3\text{N} + \text{HNO}_2$  Salt soluble in water



14. Write the main product when benzene, reacts with following reagent

(i)  $\text{HBF}_4/\Delta$


(ii)  $\text{Cu/HBr}$

(iii)  $\text{H}_2\text{O}$

Ans. (i)  $\text{N}_2^+\text{Cl}^-$




F





(ii)



Br



(iii)




OH

15. Write the method to prepare a pure sample of primary amine having one more carbon than the alkyl halide used ?




16. How will you prepare methyl orange from sulphanilic acid ?






17. Why acetamide is weaker base than ethyl amine

Ans: Due to presence of electron withdrawing carbonyl group in acetamide




18. Complete the following reactions:



Ans. (i)  $\text{R-CH}_2\text{-NH}_2$

(ii)  $\text{C}_6\text{H}_6$

(iii)




19. Give reason why Trimethylamine reacts with  $\text{BF}_3$  while triphenylamine does not.

Ans: In trimethylamine  $(\text{Me})_3\text{N}$ , the methyl group show +I effect, hence increase the reactivity towards Lewis acid ( $\text{BF}_3$ ). While in  $(\text{Ph})_3\text{N}$ , Phenyls group ( $\text{C}_6\text{H}_5$ ) show -I effect which decrease the  $e^-$  density over N atom. hence reactivity decreases towards Lewis acid ( $\text{BF}_3$ )

20. Justify with suitable reason: Why is para nitroaniline is weaker base than m-nitro aniline?


Ans. The  $e^-$  over N in both case are in resonance in ring.



## LONG ANSWER TYPE QUESTIONS (5 Marks)

1. Arrange the following :

(a) In decreasing order of  $pK_b$  value




(b) In increasing order of basic strength:

(i) Aniline, *p*-Nitroaniline and *p*-Toluidine

(ii)  $C_6H_5NH_2$ ,  $C_6H_5NHCH_3$ ,  $C_6H_5CH_2NH_2$


(c) In decreasing order of basic strength :



(d) Decreasing order of basic strength in gas phase :



(e) Increasing order of boiling point :



Ans : (a)  $C_6H_5NH_2 > C_6H_5NHCH_3 > C_2H_5NH_2 > (C_2H_5)_2NH$

(b) (i) *p*-Nitroaniline < Aniline < *p*-Toluidine



(c)  $(CH_3)_2NH_2 > CH_3NH_2 > CH_3N(CH_3)_2 > C_6H_5NH_2$

(d)  $(C_2H_5)_3N > (C_2H_5)NH > C_2H_5NH_2 > NH_3$

(e)  $(CH_3)_2NH < C_2H_5NH_2 < C_2H_5OH$




2. How will you convert the following compound:

(i) Ethanoic acid into Methanamine

(ii) Hexanenitrile into 1- Aminopentane

(iii) Methanol into Ethanoic acid


(iv) Ethanamine into Methanamine




3. Write short note on the following :-

- (a) Carbylamine reaction
- (b) Diazotization
- (c) Hoffmann bromide reaction
- (d) Coupling reaction
- (e) Ammonolysis

(i) **Carbylamine reaction** : When primary amine (aromatic or aliphatic) warmed with chloroform and alc. KOH, isocyanides are formed which can be identified by their offensive smell. This test is used to identify the presence of primary amine or chloroform.



(b) **Diazotization** : When primary aromatic amine is treated with  $\text{NaNO}_2$  and  $\text{HCl}$  at 273-278K, diazonium salt is obtained. This reaction is known as diazotization.




Benzenediazonium chloride is a very important synthetic compound, which can be changed into haloarenes, phenol, cyanobenzene, benzene etc.

- (c) **Hoffmann's bromide reaction** : When any primary amide (aliphatic or aromatic) is treated with bromine and alkali, it gives the amine with one less carbon atom.




This reaction is used to reduce one carbon atom from a compound.

- (d) **Coupling reaction** : When benzenediazonium chloride is treated with phenols or aromatic amines, azo dyes are produced in which diazo (-N=N-) group is retained. Coupling reactions generally take place at *p*-position of phenol or aromatic amines.




- (e) **Ammonolysis** : Reaction of alkyl halides with ammonia is known as ammonolysis. Ammonolysis generally gives the mixture of 1°, 2°, 3° amines and quaternary ammonium salt.



4. Complete the following reaction :-

- $\text{C}_6\text{H}_5\text{NH}_2 + \text{H}_2\text{SO}_4(\text{conc.}) \rightarrow$
- $\text{C}_6\text{H}_5\text{N}_2\text{Cl} + \text{C}_2\text{H}_5\text{OH} \rightarrow$
- $\text{C}_6\text{H}_5\text{NH}_2 + (\text{CH}_3\text{CO})_2\text{O} \rightarrow$
- $\text{C}_6\text{H}_5\text{N}_2\text{Cl} + \text{H}_3\text{PO}_2 + \text{H}_2\text{O} \rightarrow$
- $\text{C}_6\text{H}_5\text{NH}_2 + \text{CHCl}_3 + 3\text{KOH}(\text{alc.}) \rightarrow$


Ans: (i)



- (ii)  $\text{C}_6\text{H}_6 + \text{N}_2 + \text{HCl} + \text{CH}_3\text{CHO}$   
 (iii)  $\text{C}_6\text{H}_5\text{NHCOCH}_3 + \text{CH}_3\text{COOH}$   
 (iv)  $\text{C}_6\text{H}_6 + \text{H}_3\text{PO}_3 + \text{HCl} + \text{N}_2$   
 (v)  $\text{C}_6\text{H}_5\text{NC} + 3\text{KCl} + 3\text{H}_2\text{O}$


5. Accomplish the following conversion :-

- (I)  $\text{C}_6\text{H}_5\text{NO}_2 \rightarrow \text{C}_6\text{H}_5\text{COOH}$   
 (ii) Benzene  $\rightarrow m$ -Bromophenol  
 (iii)  $\text{C}_6\text{H}_5\text{COOH} \rightarrow \text{C}_6\text{H}_5\text{NH}_2$   
 (iv) Aniline  $\rightarrow 2,4,6$ -Tribromoaniline  
 (v) Benzylchloride  $\rightarrow 2$ -Phenylethanamine






7. Write A, B and C in the given reaction sequences;




- Ans. (i)  $\text{C}_6\text{H}_5\text{CN}$ ,  $\text{C}_6\text{H}_5\text{COOH}$ ,  $\text{C}_6\text{H}_5\text{CONH}_2$   
 (ii)  $\text{CH}_3\text{CH}_2\text{CN}$ ,  $\text{CH}_3\text{CH}_2\text{NH}_2$ ,  $\text{CH}_3\text{CH}_2\text{OH}$   
 (iii)  $\text{C}_6\text{H}_5\text{NH}_2$ ,  $\text{C}_6\text{H}_5\text{N}_2^+\text{Cl}^-$ ,  $\text{C}_6\text{H}_5\text{OH}$   
 (iv)  $\text{CH}_3\text{CONH}_2$ ,  $\text{CH}_3\text{NH}_2$ ,  $\text{CH}_3\text{OH}$   
 (v)  $\text{CH}_3\text{CH}_2\text{CN}$ ,  $\text{CH}_3\text{CH}_2\text{-CO-NH}_2$ ,  $\text{CH}_3\text{-CH}_2\text{-NH}_2$

8. An organic compound 'A' on treatment with aqueous ammonia and heating forms compound 'B' which on heating with  $\text{Br}_2$  and  $\text{KOH}$  forms a compound 'C' of molecular formula  $\text{C}_6\text{H}_7\text{N}$ . Write the structure and IUPAC of compound A, B and C.



9. Write the structure of compound A to E in the following sequence of reaction.




Ans.

A=




Benzene

B=



Nitrobenzene

C=



Aniline

D=




Phenylisocyanide

E=



N-methylaniline


10. Write the structure A, B, C, D, E in the given reaction.




Ans.

A=  $\text{C}_6\text{H}_5\text{NHCOCH}_3$ 

B=




C=



D=



E=



### CASE-STUDY BASED QUESTIONS

1. **Read the passage and answer the following questions:**

Friedel-Crafts (FC) reaction is an important method to incorporate carbon skeletons into aromatic system. Great successes have been achieved for the hydroarylation of neutral arenes (such as toluene, anisole, and their homologues). Because the FC reactions typically require Lewis acid catalysts, for arenes containing nitrogen atom, the substrate scope of FC reactions are quite limited due to the coordination between amine and Lewis acid catalyst, except indole and pyrrole. Being profited from the extremely weak basic properties, acid-catalyzed additions of indole and pyrrole to alkenes have obtained great achievements. However, the hydroarylation of alkaline arenes to alkenes still remains many challenges. Some researches have shown the possibility of hydroarylation between the parent anilines  $C_6H_5NH_2$  and alkenes. However, the reaction of arenes with stronger basicity (such as N,N-dimethylaniline and N,N-diethylaniline) still is a big problem, due to their ability to coordinate with Lewis acid catalyst which can lead to deactivation of the aromatic ring. Furthermore, alkaline arenes can trap the proton in the C-H activation process and the reaction will be terminated as result.


Recently, Bertrand et al. reported an anti-Bredt cyclic diaminocarbene which showed increased  $\lambda$ -accepting character without diminishing its  $\sigma$ -donor property. We found that Gold(I) compound derived from this new carbene can be used as effective catalyst for the FC reaction between alkenes and N,N-dialkylanilines. Now, these new FC reactions are receiving more and more research interests. As we known, most of the electrophilic substitution reactions followed the Markovnikov rule. For the FC reaction of alkenes, the reactions following the Markovnikov rule should form branched product. Only several examples were reported on the formation of linear product by anti-Markovnikov rule. For the FC reactions between alkenes and N, N-dialkylanilines catalyzed by carbene Gold(I), both Markovnikov and anti-Markovnikov hydroarylations were observed and all these reactions gave high para-selectivity products. The selectivity to the branched or linear product was highly depended on the structure of alkenes.

**Reference :** Wu, H., Zhao, T. & Hu, X. **Friedel-Crafts Reaction of N,N-Dimethylaniline with Alkenes Catalyzed by Cyclic Diaminocarbene-Gold(I) Complex.** *Sci Rep* 8, 11449 (2018). <https://doi.org/10.1038/s41598-018-29854-0>

- (A) Why aniline does not undergoes Friedel-Crafts reaction?
- (B) Write reaction to convert aniline into N,N-dimethylaniline.
- (C) How *p*-Nitroaniline can be synthesised from Aniline?
- (D) Why -NH<sub>2</sub> group in aniline is *ortho*-, *para*-directing?

2. **Read the passage and answer the following questions:**

The development of base metal catalysts for industrially relevant amination and hydrogenation reactions by applying abundant and atom economical reagents continues to be important for the cost-effective and sustainable synthesis of amines which represent highly essential chemicals. In particular, the synthesis of primary amines is of central importance because these compounds serve as key precursors and central intermediates to produce value-added fine and bulk chemicals as well as pharmaceuticals, agrochemicals and materials. Here we report a Ni-triphos complex as the first Ni-based homogeneous catalyst for both reductive amination of carbonyl compounds with ammonia and hydrogenation of nitroarenes to prepare all kinds of primary amines. Remarkably, this Ni-complex enabled the synthesis of functionalized and structurally diverse benzylic, heterocyclic and aliphatic linear and branched primary amines as well as aromatic primary amines starting from inexpensive and easily accessible carbonyl compounds (aldehydes and ketones) and nitroarenes using ammonia and molecular hydrogen. This Ni-catalyzed reductive amination methodology has been applied for the amination of more complex pharmaceuticals and steroid derivatives. Detailed DFT computations have been performed for the Ni-triphos based reductive amination reaction, and they revealed that the overall reaction has an inner-sphere mechanism with H<sub>2</sub> metathesis as the rate determining step.



**Reference :** Kathiravan Murugesan, Zhihong Wei, Vishwas G. Chandrashekhar, Haijun Jiao, Matthias

Beller, Rajenahally V. Jagadeesh **General and selective synthesis of primary amines using Ni-based homogeneous catalysts**\_Chem. Sci., 2020,11, 4332-4339

(A) Convert nitrobenzene to chlorobenzene.

(B) What happens when butanone reacts with ammonia according to scheme given above.

(C) Mention one method, other than mentioned here for conversion of nitrobenzene to aniline.

3. **Read the passage and answer the following questions:**

For a group of nitro-substituted anilines and diphenylamines, a plot of PICHA (ordinate) against  $\text{pK}_{\text{BH}}^+$  gave a straight line of slope 0.6. That is, the acidities of the amines seemed to be affected to a smaller extent by ring substitution than were their basicities.



This result was somewhat surprising because, although resonance with the nitro group occurs in both the neutral amine and the anion, and hence affects both equilibria, it involves a very important charge delocalization in the amide anion.



It has been suggested that the nitro substituent has an anomalous effect in acid mixtures (2), where some of the  $\text{pK}_{\text{BH}}^+$  values had been determined. In the course of extending the H— scale to very basic values, we had the opportunity to use amines containing substituents other than nitro (3) and to determine if the unusual effect of substituents on acidity vis-a-vis basicity was due to the presence of nitro groups. In accompanying papers the  $\text{pK}_{\text{BH}}^+$  values (4) and the  $\text{pK}_{\text{HA}}$  values of these aromatic amines are reported. (The basicities of amines are described, as is customary, in terms of the acidity of their conjugate acids.)

**Reference :** Ross Stewart and Douglas Dolman. **A comparison of the acidity and basicity of aromatic amines.** *Canadian Journal of Chemistry*. 45(9): 925-928.  
<https://doi.org/10.1139/v67-156>

- (A) Which of the following has highest  $\text{P}_{\text{Ka}}$  value?
- (a) Aniline
  - (b) *p*-Nitroaniline
  - (c) *m*-Nitroaniline
  - (d) *o*-Nitroaniline
- (B) Why basic strength of aliphatic amines are higher than ammonia?
- (C) What will be the correct order of basic strength of following:  $\text{EtNH}_2$ ,  $(\text{Et})_2\text{NH}$ ,  $(\text{Et})_3\text{N}$
- (D) What happens when aniline reacts with sulphuric acid?

## ANSWERS

### I MULTIPLE CHOICE QUESTIONS

1. b 2. d 3. a 4. d 5. c 6. a 7. b 8. c 9. c 10. a  
11. c 12. d 13. a 14. a 15. c 16. b 17. a 18. c 19. a 20. c

### II FILL IN THE BLANKS

- |                      |                       |
|----------------------|-----------------------|
| 1. Aniline           | 2. weaker, stronger   |
| 3. Primary aliphatic | 4. Hinsberg           |
| 5. Butanamine        | 6. Acetylation        |
| 7. <i>meta</i>       | 8. Water.             |
| 9. H-bonding         | 9. Fluorides, Iodides |

### III ASSERTION REASON TYPE QUESTIONS

1. c 2. c 3. a 4. b 5. a 6. c 7. a 8. a 9. b 10. a


### IV ONE WORD TYPE QUESTIONS

- |                            |                          |
|----------------------------|--------------------------|
| 1. Carbylamine reaction    | 2. 2,4,6-tribromoaniline |
| 3. Aniline                 | 4. weaker                |
| 5. Friedel-Crafts reaction | 6. Benzene               |
| 7. Sandmeyer               | 8. $S_N2$                |
| 9. pH-9 to 10              | 10. $Sn + HCl$           |
| 11. 8                      | 12. $LiAlH_4$            |
| 13. Novacain               | 14. $SN^1/SN^2$          |
| 15. Pyridine               |                          |

### CASE STUDY BASED QUESTIONS

PASSAGE : 1


(A) Aniline forms salt with anhydrous  $AlCl_3$ .



(C) Refer NCERT Page - 403

(D) Due to +R effect of  $-NH_2$  Sp.

PASSAGE : 2



PASSAGE : 3

- (A) (d)
- (B) Due to +I effect of -R gps present in aliphatic amines
- (C)  $(\text{Et})_2\text{NH}$ ,  $(\text{Et})_3\text{N}$ ,  $\text{EtNH}_2$



## UNIT TEST-1

## CHAPTER-13

Maximum Marks : 20

Time Allowed : 1 Hour

1. Benzamide reacts with  $\text{Br}_2$  and  $\text{NaOH}$  forming..... 1
2. The correct order of basicity of amines is: 1
  - (a) Ammonia > Ethanamine > Aniline
  - (b) Ethanamine > Ammonia > Aniline
  - (c) Ammonia > Aniline > Ethanamine
  - (d) Ethanamine > Aniline > Ammonia
3. When aniline reacts with  $\text{NaNO}_2/\text{HCl}$  then reaction with  $\text{CuCN}$  followed by acidic hydrolysis. What will be the final product of the reaction? 1
  - (a) Nitrobenzene
  - (b) Benzaldehyde
  - (c) Benzoic acid
  - (d) Phenol
4. Assertion: Aniline forms 2,4,6-Tribromoaniline on reaction with bromine water.  
Reason:  $-\text{NH}_2$  is *ortho, para*- directing. 1
  - (a) Both assertion and reason are correct statements and reason is correct explanation of assertion.
  - (b) Both assertion and reason are correct statements but reason is not correct explanation of assertion.
  - (c) Assertion is correct statement but reason is wrong statement.
  - (d) Assertion is wrong statement but reason is correct statement.
5. Assertion: Primary amines cannot be synthesised by Gabriel phthalimide synthesis.  
Reason: Due to steric hindrance caused by bulky group substitution is not possible.
- (a) Both assertion and reason are correct statements and reason is correct explanation of assertion. 1
  - (b) Both assertion and reason are correct statements but reason is not correct explanation of assertion.
  - (c) Assertion is correct statement but reason is wrong statement.
  - (d) Assertion is wrong statement but reason is correct statement.
6. How will you synthesise Butanamine by Gabriel phthalimide synthesis? 2
7. Write chemical test distinguish to between following pair of compounds: 2
  - (a) Aniline and Benzylamine


- (b) Methylamine and Dimethylamine
8. How will you convert the following compound: 2
- (a) Ethanoic acid to methanamine
- (b) Aniline to *p*-romoaniline
9. An aromatic compound 'A' on treatment with aqueous ammonia and heating forms compound 'B' which on heating with Br<sub>2</sub> and KOH forms a compound 'C' of molecular formula C<sub>6</sub>H<sub>13</sub>N. Write the structures and IUPAC names of compounds A, B and C. 3
10. Write short notes on following: 3
- (a) Benzoylation reaction
- (b) Hoffmann bromamide degradation reaction
- (c) Carbylamine reaction
11. Explain with suitable reason: 3
- (a) Acetylation of aniline reduces its activation effect.
- (b) CH<sub>3</sub>NH<sub>2</sub> is more basic than CH<sub>3</sub>CONH<sub>2</sub>.
- (c) Nitration of aniline gives significant amount of *meta*-Nitroaniline in addition to *o*- and *p*-Nitroaniline.

**UNIT TEST-II**  
**CHAPTER-13**

**Maximum Marks : 20****Time Allowed : 1 Hr**

1. Give the IUPAC of following compound.

1



2. Give reason:- aniline does not undergo friedel craft alkylation.
3. How will you convert Benzene into aniline
4. What is Hinsberg reagent.
5. Out of butan-1-ol and but-1-amine which one has higher boiling point.
6. Write short note on following:  
(i) Gabriel phthalimide synthesis  
(ii) diazotisation
7. Write chemical distinguish test between following pair.  
(a) Aniline and benzylamine  
(b) Ethane amine and N, N-dimethylethanamine.
8. Complete the following reaction.  
(i)  $\xrightarrow{(\text{CH}_3\text{CO})_2\text{O}}$  A  $\xrightarrow{\text{Br}_2}$  B  
(ii)  $\xrightarrow[0-5^\circ\text{C}]{\text{HNO}_2}$  A  $\xrightarrow{\text{H}_3\text{PO}_2}$  B
9. Arrange the following as per mentioned properties.  
(a) Ethane amine, N-ethylethaneamine, N-N-diethylethane amine (Basic strength in aqueous medium)  
(b) Aniline, p-Nitroaniline, p-Tolidine (increasing basic character)
10. Convert the following compounds:  
(a) Benzamide to toluene  
(b) Aniline to 2,4,6 - Tribromofluorobenzene
11. Two isomeric compound 'A' and 'B' having molecular formula  $\text{C}_4\text{H}_{11}\text{N}$ , both lose  $\text{N}_2$  on treatment with  $\text{HNO}_2$  and gives compound 'C' and 'D' respectively. 'C' is resistant to oxidation but immediately respond to Lucas reagent. Whereas 'D' respond to Lucas reagent after 5 minutes and gives a positive iodoform test. Identify A, B, C, D.


# UNIT 14


# BIOMOLECULES

## Points to Remember

# Biomolecules

These are the macromolecules essential for survival of life, e.g. carbohydrates, proteins, etc.





1. **Carbohydrates** : These are optically active polyhydroxy aldehydes or ketones or the compounds which produce these on hydrolysis.

2. **Classification** :

(i) **Monosaccharides** : Those carbohydrates which cannot be hydrolysed into further simpler carbohydrates. e.g, glucose, fructose, galactose etc.

(ii) **Disaccharides** : Those carbohydrates which produces two monosaccharides on hydrolysis. e.g, sucrose, maltose and lactose.

(iii) **Oligosaccharides** : Those carbohydrates which give two to ten monosaccharides on hydrolysis.

(iv) **Polysaccharides** : Those carbohydrates which on hydrolysis give large number of monosaccharides on hydrolysis. e.g, starch, cellulose, glycogen.

3. **Sugar** : Carbohydrates which are sweet in taste.


(i) **Reducing sugars** : Those which reduce Fehling's or Tollens' reagent due to availability of free aldehydic groups. e.g, glucose, fructose, galactose.

(ii) **Non-reducing sugars** : Those which do not reduce Fehling's or Tollens' reagent. They do not have free aldehydic group. e.g, sucrose.


4. **Glucose** : It is a monosaccharide with molecular formula  $C_6H_{12}O_6$ .

5. **Preparation** :

(i) **From sucrose** :




(ii) **From starch** :



6. **Structure** :

**Fischer structure** :


(+) glucose has 'D' configuration as shown :



D-(+)-glucose

'D'- means – OH group on first chiral 'C' from the bottom is on right hand and (+) means it is dextrorotatory *i.e.*, it rotates plane polarized light towards right.

### Reactions of glucose :



### Objections against open chain structure of glucose

The open chain structure was unable to explain the following reactions :

- (a) It does not give the 2,4-DNP test, Schiff's test and does not form the hydrogensulphite addition product with  $\text{NaHSO}_3$ .
- (b) The pentacetate of glucose does not react with  $\text{NH}_2\text{OH}$ , indicating the absence of free aldehydic group.
- (c) Glucose exist in 2 different crystalline forms  $\alpha$  and  $\beta$  forms. These are called anomers. They differ in optical rotation, they also differ in melting point.

After which a close chain (cyclic) structure of glucose was proposed by Haworth.

\* Anomers are isomers which have a different configuration at C-1 functional group

7. **Glycosidic linkage** : The linkage between two monosaccharide units through oxygen is called the glycosidic linkage.
8. **Proteins** : These are macro molecules made up of amino acids joined by amide linkage ( $-\text{CONH}-$ ) is called as peptide linkage. These are required for growth and development of the body.
9. **Amino acids** : These contain an amino ( $-\text{NH}_2$ ) and an acidic ( $-\text{COOH}$ ) group and are therefore amphoteric in nature. In solution, they exist in the form of twitter ion (a dipolar ion).
10. **Native state of protein** : The parental state or the natural state in which the protein is found.
11. **Denaturation of protein** : Destruction of the native state of protein is denaturation. It can be brought by physical and chemical methods. The  $2^\circ$  and  $3^\circ$  structures are destroyed, only  $1^\circ$  structure is retained.

**Enzymes** : These are biocatalyst and generally globular proteins *e.g.*, invertase, zymase, phenylalanine hydroxylase, urease etc.

### Main characteristics of enzymes :

- (i) It speed up the biological reaction upto million times.



- (ii) It is highly specific and work on lock and key theory.
- (iii) It is highly sensitive to pH and temperature.
12. **Nucleic acids :** These are biomolecules which are long chain polymers of nucleotides. They are of two types :
- (i) **Deoxyribonucleic acid (DNA)**
- (ii) **Ribonucleic acid (RNA)**
13. Nucleoside = Base + Sugar  
Nucleotide = Base + Sugar + Phosphoric acid

### OBJECTIVE TYPE QUESTIONS

#### I MULTIPLE CHOICE QUESTIONS

- Which of the following acids is a vitamin?
  - aspartic acid
  - ascorbic acid
  - asipic acid
  - saccharic acid
- Non-reducing sugar out of following is -
  - glucose
  - sucrose
  - maltose
  - lactose
- In a protein molecule amino acids are linked together by:
  - peptide bond
  - coordinate bond
  - glycosidic bond
  - phosphodiester bond
- One strand of DNA has the sequence. ATGCTT, the sequence of complementary strand would be:
  - TCCGAA
  - TACGTA
  - TACGAA
  - TAGCTA
- Which of the following vitamin is water soluble?
  - vitamin C
  - vitamin D
  - vitamin K
  - vitamin E
- In both DNA and RNA, base and phosphate ester linkage are at -
  - $C_5'$  and  $C_2'$  respectively of sugar molecule
  - $C_2'$  and  $C_5'$  respectively of sugar molecule
  - $C_3'$  and  $C_5'$  respectively of sugar molecule
  - $C_5'$  and  $C_1'$  respectively of sugar molecule

7. The two functional groups present in a typical carbohydrates are:  
(a)  $\text{-OH}$  and  $\text{-COOH}$  (b)  $\text{-CHO}$  and  $\text{-COOH}$   
(c)  $\text{>C=O}$  and  $\text{-OH}$  (d)  $\text{-CHO}$  and  $\text{-COCl}$
8. The presence or absence of hydroxyl group on which carbon atom of sugar differentiates RNA and DNA.  
(a) 1st (b) 2nd  
(c) 3rd (d) 4th
9. The carbohydrate known as invert sugar is -  
(a) lactose (b) sucrose  
(c) maltose (d) glucose
10. Pick the disaccharide from following:  
(a) maltose (b) cellulose  
(c) maltase (d) starch
11. Which one of the following is not an aldose?  
(a) Glucose (b) Ribose  
(c) Fructose (d) Galactose
12. Biomolecule containing transition metal is-  
(a) Vitamin C (b) Chlorophyll  
(c) Haemoglobin (d) RNA
13. Which of the following does not have glycosidic linkage?  
(a) Maltose (b) Amylose  
(c) Galactose (d) Sucrose
14. Fibrous proteins are present in:  
(a) Haemoglobin (b) Albumin  
(c) Collagen (d) Insulin
15. Hydrolysis of lactose with dilute acid yields  
(a) equimolar mixture of D-glucose and D-fructose  
(b) equimolar mixture of D-glucose and D-galactose  
(c) equimolar mixture of D-galactose and D-fructose  
(d) equimolar mixture of D-galactose and D-sucrose

16. Match the carbohydrate in Column I with its characteristic given in Column II

**Column-I**

(A) Lactose

(B) Starch

(C) Sucrose

(D) Fructose

(a) A - s, B - r, C - p, D - q

(b) A - p, B - q, C - r, D - s

(c) A - r, B - s, C - p, D - q

(d) A - s, B - r, C - q, D - p

**Column-II**

(p) Ketohexose

(q) Disaccharide

(r) Polysaccharide

(s) on hydrolysis gives  $\beta$ -D-glucose and  $\beta$ -D-galactose

17. Match the carbohydrate in Column I with its characteristic given in Column II

**Column-I**

(A) Keratin

(B) Haemoglobin

(C) Riboflavin

(D) Glycine

(a) A-p, B-q, C-s, D-r

(b) A-q, B-p, C-s, D-r

(c) A-q, B-p, C-r, D-s

(d) A-s, B-r, C-q, D-p

**Column-II**

(p) protein

(q)  $\beta$ -pleated protein(r)  $\alpha$ -amino acid

(s) Water soluble vitamin

18. The no. of chiral carbon present in  $\beta$ -D-(+)-glucose is:

(a) 2

(b) 4

(c) 5

(d) 1

19. Which of the following base is not present in RNA?

(a) Adenine

(b) Uracil

(c) Cytosine

(d) Thymine

20. Hormone produced under stress which stimulates glycogenolysis in the liver of human being ?

(a) thyroxine

(b) Insulin

(c) Adrenaline

(d) Estradiol

**II FILL IN THE BLANKS**

- The disease beri-beri is caused due to lack of .....
- Scurvy is caused due to deficiency of .....
- .....(carbohydrate) is not digested by human beings but digested by herbivorous animals.
- .....on hydrolysis gives D-glucose and D-galactose.

5. Amylose and amylopectin are the two components of.....
6. Carbohydrates which yields a large number of monosaccharide units on hydrolysis are called .....
7. Carbohydrates which reduce Tollens' reagent are called .....
8. Deficiency of ..... leads to xerophthalmia and night blindness.
9. ....contains pentose sugar, and base whereas.....contains pentose sugar, base as well as phosphate group.
10. The pair of stereoisomers which differ only in the configuration of the hydroxyl group at C-1 are called .....

### III ASSERTION-REASON TYPE QUESTIONS

In each of the following questions, a statement of Assertion (A) is given followed by a corresponding statement of Reason (R) just below it. Of the statements, mark the correct answer as

- (a) Both assertion and reason are correct, and reason is the correct explanation of the assertion.
  - (b) Both assertion and reason are correct, but reason is not the correct explanation of the assertion.
  - (c) Assertion is correct, but reason is incorrect.
  - (d) Assertion is incorrect but reason is correct.
1. **ASSERTION :** A solution of sucrose in water is dextrorotatory but on hydrolysis in presence of little HCl it becomes laevorotatory.  
**REASON :** Sucrose on hydrolysis gives unequal amount of glucose and fructose as a result sign of rotation changes.
  2. **ASSERTION :** Fructose does not contain aldehyde group but still reduce Tollens' reagent.  
**REASON :** In the presence of base, fructose undergoes rearrangement to form glucose and mannose.
  3. **ASSERTION :** D-(+)-Glucose is dextrorotatory in nature.  
**REASON :** 'D' represents its dextrorotatory nature.
  4. **ASSERTION :** Vitamin D can be stored in our body.  
**REASON :** Vitamin D is fat soluble vitamin.

5. **ASSERTION :** All naturally occurring  $\alpha$ -amino acids except glycine are optically active.  
**REASON :** Most naturally occurring amino acids have D-configuration.
6. **ASSERTION :** In presence of enzyme, substrate molecule can be attacked by the reagent effectively.  
**REASON :** Active sites of enzymes hold the substrate, molecule in a suitable position.
7. **ASSERTION:** Sucrose is a non-reducing sugar.  
**REASON :** It has glycosidic linkage.
8. **ASSERTION:** Vitamin C has to be continuously supplied through diet.  
**REASON:** Vitamin C is a water soluble vitamin.
9. **ASSERTION :** Cellulose is not digested by human beings.  
**REASON :** Cellulose is a polymer of  $\beta$ -D-glucose.
10. **ASSERTION:** Non-essential amino acids are not necessary for protein synthesis.  
**REASON:** Non-essential amino acids are produced in the human body.

#### **IV ONE WORD ANSWER TYPE QUESTIONS**

1. Name the component of starch which is water soluble.
2. Write the product formed when glucose is treated with HI.
3. What are the products of hydrolysis of maltose?
4. Name the purines present in DNA.
5. Write the name of linkage joining two amino acids.
6. The deficiency of which vitamin causes the disease pernicious anaemia.
7. Name the base that is found in nucleotide of RNA only.
8. Name the vitamin whose deficiency is responsible for poor coagulation of blood.
9. Write the product formed on reaction of D-glucose with  $\text{Br}_2$  water.
10. Name the polysaccharide which is stored in the liver of animals.

**VERY SHORT ANSWER TYPE QUESTIONS (1Marks)**

**Q. 1. What structural feature is required for a carbohydrate to behave as reducing sugar ?**

**Ans.** The carbonyl group of any one monosaccharide present in carbohydrate must be free.

**Q. 2. Give the significance of (+) sign in the name D-(+)-glucose.**

**Ans.** (+) sign indicates dextro-rotatory nature of glucose.

**Q. 3. Glucose is an aldose sugar but it does not react with sodium hydrogen sulphite. Give reason.**

**Ans.** The – CHO group reacts with – OH group at C-5 to form a cyclic hemiacetal.

**Q. 4. Why is sucrose called invert sugar ?**

**Ans.** When sucrose is hydrolysed by water, the optical rotation of solution changes from positive to negative.

**Q. 5. Name the amino acid which is not optically active.**

**Ans.** Glycine.

**Q. 6. Give reason : Amylase present in the saliva becomes inactive in the stomach.**

**Ans.** HCl present in stomach decreases the pH.

**Q. 7. Name the interactions responsible for the stability of  $\alpha$ -helical structure of proteins**

**Ans.** Hydrogen bonding.

**Q. 8. Which nucleic acid is responsible for carrying out protein synthesis in the cell ?**

**Ans.** RNA (Ribonucleic acid)


**Q. 9. When RNA is hydrolysed, there is no relationship among quantities of different bases obtained. What does this fact suggest about structures of RNA ?**

**Ans.** RNA is single stranded.

**Q. 10. What type of linkage holds together the monomers of DNA and RNA ?**

**Ans.** Phosphodiester linkage.

**Q. 11. Give the Howarth projection of D-glucopyranose.**



**Q. 12. Where does the water present in the egg go after boiling the egg ?**

**Ans.** On boiling, during denaturation process water gets absorbed in denaturated proteins.

**Q. 13. Name a protein which is insoluble in water.**

**Ans.** Keratin.

**Q. 14. Mention two important functions of carbohydrates in plants.**

**Ans.** Major energy source, storage molecules like starch in plants.

**Q. 15. Name the different types of RNA molecules found in cells of organisms.**

**Ans.** tRNA, mRNA, rRNA.

**Q. 16. Why are carbohydrates generally optically active ?**

**Ans.** Because they contain one or more chiral carbon.

**Q. 17. During curdling of milk, what happens to sugar present in it ?**

**Ans.** Lactose changes to lactic acid.

**Q. 18. The two strands in DNA are not identical but complementary. Explain.**

**Ans.** Base pairing rule is followed; A = T and G  $\equiv$  C. (Hydrogen bonding between complementary bases)

**Q. 19. If one strand of DNA has the sequence 5'-G-G-A-C-T-A-C-T-3', what is the sequence of bases in the complementary strand ?**

**Ans.** 3'-C-C-T-G-A-T-G-A-5'

**Q. 20. What are monosaccharides ?**

**Ans.** Sugars which cannot be hydrolysed to give simpler units or compounds.

**Q. 21. What is the difference between native protein and denatured protein ?**

**Ans.** Proteins found in a biological system with unique 3D-structure and biological activity is called native protein. When native protein is subjected to physical and chemical change, protein loses its biological activity and is called denatured protein.

**SHORT ANSWER TYPE QUESTIONS (2 Marks)**

**Q. 1. Define the following terms in relation to proteins :**

- (i) **Peptide linkage**                      (ii) **Denaturation**

**Ans. (i) Peptide linkage :** A link between two amino acids with loss of water – CO – NH – peptide linkage.

- (ii) A process that changes the three dimensional structure of native protein is called denaturation of protein. It results into breaking of hydrogen bonds and disulphide linkages. Thus, a completely denatured protein has a shape of random coil.

**Q. 2. List the reactions of glucose which cannot be explained by its open chain structure.**

- Ans. (i)** Despite having the aldehyde group, glucose does not give 2, 4 DNP test or Schiff's test.
- (ii) It does not form hydrogensulphite addition product with  $\text{NaHSO}_3$ .
- (iii) The penta acetate of glucose does not react with hydroxylamine indicating the absence of free – CHO group.

**Q. 3. Explain what is meant by :**

- (i) **Biocatalyst**                      (ii) **Glycosidic linkage**

**Ans. (i)** Biocatalysts are the catalysts which increases the rate of metabolism biochemical reactions.

- (ii) The linkage between the monosaccharide units through oxygen is called glycosidic linkage.



**Q. 4. Explain the following terms :**

**(i) Invert sugar**

**(ii) Polypeptides**

**Ans. (i)** An equimolar mixture of glucose and fructose produced on hydrolysis of sucrose is called invert sugar. It is called so because sucrose is dextrorotatory whereas its hydrolysis product is laevorotatory.

**(ii)** Polypeptides are polymers of amino acids containing less than 100 amino acids. For example, oxytocin, vasopressin, etc.

**Q. 5. Name the product of hydrolysis of sucrose. Why is sucrose not a reducing sugar ?**

**Ans.** On hydrolysis, sucrose gives equimolar mixture of D-(+)-glucose and D-(-)-fructose. Sucrose is not a reducing sugar as glucose and fructose are linked through their reducing centres in structure of sucrose.

**Q. 6. Explain nucleotides and nucleosides.**

**Ans.** A nucleoside contains only two basic components of nucleic acids *i.e.*, pentose sugar and nitrogenous base.

A nucleotide contains all the three basic components of nucleic acids *i.e.*, a phosphoric acid group, pentose sugar and nitrogenous base.

**Q. 7. Describe what do you understand by primary structure and secondary structure of proteins.**

**Ans. Primary structure of proteins :** The protein in which amino acids are linked with each other in a specific sequence is said to be the primary structure of that protein.


**Secondary structure of proteins :** It refers to the shape in which a long polypeptide chain can exist *i.e.*,  $\alpha$ -helix and  $\beta$ -pleated structure.

**Q. 8. What is essentially the difference between  $\alpha$ -form of glucose and  $\beta$ -form of glucose ? Explain.**

**Ans.**  $\alpha$ -form of glucose and  $\beta$ -form of glucose differ only in the configuration of the hydroxyl group at C<sub>1</sub> in cyclic structure of glucose/hemiacetal form of glucose.

**Q. 9. What are anomers ? Give the structures of two anomers of glucose.**

**Ans.** Monosaccharides which differs in configuration at functional group C-atom ( $C_1$  and  $C_2$ ), *e.g.*,  $\alpha$ -glucose and  $\beta$ -glucose.




**Q. 10. Write the hydrolysed product of :**

- (i) Maltose                      (ii) Cellulose

**Ans.** (i)  $\alpha$ -D-glucose                      (ii)  $\beta$ -D-glucose

**Q. 11. (i) Acetylation of glucose with acetic anhydride gives glucose penta-acetate. Write the structure of penta acetate.**

(ii) Explain why glucose penta acetate does not react with hydroxylamine ?





(ii) The molecule of glucose penta acetate has a cyclic structure in which  $-\text{CHO}$  is involved in ring formation.

**Q. 12. Write the products of oxidation of glucose with :**

(i) Bromine water

(ii) Nitric acid





**Q. 13. State two main differences between globular and fibrous proteins.**

Ans.	Globular protein	Fibrous protein
(i)	They form a $\alpha$ -helix structure.	(i) They have $\beta$ -pleated structure.
(ii)	They are water soluble.	(ii) They are water insoluble.

**Q. 14. What are essential and non-essential amino acid ? Give two examples of each type.**

**Ans.** Essential amino acids are those which are not produced in our body and required to be supplied from outside, *e.g.*, valine, leucine.

Non-essential amino acids are those which are produced by our body, *e.g.*, glycine, alanine.

**Q. 15. Coagulation of egg white on boiling is an example of denaturation of protein. Explain it in terms of structural changes.**

**Ans.** Protein albumin present in egg white gets denatured *i.e.*, 2° & 3° structures are destroyed and 1° structure is retained.

**Q. 16. Describe two important functions of nucleic acids.**

**Ans.** (i) DNA is responsible for transfer of heredity information from one generation to another.

(ii) RNA is responsible for protein synthesis.

**Q.17.(i) What type of linkage is responsible for the formation of proteins ?**

**(ii) Write the product formed when glucose is treated with HI.**

**Ans.** (ii) Peptide linkage.

(iii) n-hexane.

**Q.18. Differentiate between the following :**

- (i) Secondary and tertiary structure of protein**
- (ii)  $\alpha$ -helix and  $\beta$ -pleated sheet structure of protein**

**Ans. (i) Secondary structure** is responsible for the shape of protein  $\alpha$ -helix and  $\beta$ -pleated sheets in which polypeptide chains have peptide bonds.

**Tertiary structure** represents overall folding of polypeptide chain and give rise to the fibrous or globular molecular shape.

- (ii)  $\alpha$ -helix structure :** The peptide chains coiled up to form right handed helix involving H-bonding (Intramolecular).

**$\beta$ -pleated sheets :** The peptide chains lie side by side together by intermolecular hydrogen bonding.


**Q.19.(i) Name the four bases present in DNA.**

- (ii) Which of them is not present in RNA ?**

- (iii) Give the structure of a nucleotide of DNA.**

**Ans. (i)** Adenine, Guanine, Thymine, Cytosine.

- (ii) Thymine.**



**Q.20. Glucose or sucrose are soluble in water but cyclohexane and benzene are insoluble in water. Explain.**

**Ans.** Glucose contain 5 – OH groups and sucrose contain eight – OH groups, because of this they form intermolecular hydrogen bonding, so they are soluble in water. But benzene and cyclohexane doesn't contain – OH groups hence doesn't form intermolecular hydrogen bonding, so they are not soluble in water.

### CBSE STUDY BASED QUESTIONS

**1. Read the passage given below and answer the following questions:**

Living systems are made up of various complex biomolecules like carbohydrates, proteins, nucleic acids, lipids, etc. Proteins and carbohydrates are essential constituents of our food. Carbohydrates are the main source of energy that is ingested by the human body. Brain mainly utilizes the glucose. Red blood cells also use glucose only. Fiber in the diet is not digested by human body due to lack of cellulase enzyme. Glucose is the major energy source in the body. Glycogen is the storage form of glucose and glycogen is stored in skeletal muscles and liver. If glucose intake exceeds than it is utilized in the body it is converted into fat. Riboses are utilized in formation of deoxyribonucleic acid. Carbohydrates are polyhydroxy alcohol with potentially active carbonyl group which may be aldehyde or keto group. Carbohydrates can be classified on the basis of carbon atom present in the carbohydrates. Carbohydrates are classified into four types monosaccharides, disaccharides, oligosaccharides, polysaccharides. Monosaccharides cannot be hydrolyzed further into simpler form. They may contain 3-7 carbon atoms but monosaccharides containing 5-6 carbon atoms are more abundant in nature. All monosaccharides reduce Tollens' reagent as well as Fehling's solution and hence are called reducing sugars. Pentoses and hexoses have cyclic structures, furanose and pyranose. Disaccharides give two monosaccharides on hydrolysis. Polysaccharides may be homopolysaccharides and heteropolysaccharides. Plants produce carbohydrates by photosynthesis. In most animals, carbohydrates are the quickly accessible reservoir of energy. The main function of carbohydrates is to provide energy, but they also play an important role in the structure and function of the body organs and nerve cells.

**The following questions are multiple choice questions. Choose the most appropriate answer:**

- (A) Which of the following statements is **not** true about glucose?
- (a) It is an aldohexose.
  - (b) On heating with HI it forms n-hexane.
  - (c) It is present in furanose form.
  - (d) It does not give 2, 4-DNP test.
- (B) The  $\alpha$ - and  $\beta$ -forms of glucose are
- (a) isomers of D(+) glucose and L(−) glucose respectively
  - (b) anomers of glucose
  - (c) isomers which differ in the configuration of C-2
  - (d) isomers which differ in the configuration of C-5
- (C) The monosaccharide constituents of lactose are:
- (a)  $\alpha$ -D-glucose and  $\beta$ -D-fructose
  - (b)  $\alpha$ -D-glucose only
  - (c)  $\beta$ -D-glucose only
  - (d)  $\beta$ -D-glucose and  $\beta$ -D-galactose
- (D) Glycogen is a branched chain polymer of  $\alpha$ -D-glucose units in which chain is formed by C1-C4 glycosidic linkage whereas branching occurs by the formation of C1–C6 glycosidic linkage. Structure of glycogen is similar to \_\_\_\_\_.
- (a) Amylose
  - (b) Amylopectin
  - (c) Cellulose
  - (d) Glucose

**2. Read the passage given below and answer the following questions:**

Proteins are very important biomolecules of living systems.  $\alpha$ -Amino acids are the building blocks of proteins. About 20  $\alpha$ -amino acids have been isolated by the hydrolysis of proteins. Ten amino acids which the body cannot synthesize are called essential amino acids. The remaining ten are called non-essential amino acids. Proteins are complex nitrogenous polymers of amino acids connected through peptide bonds. Protein is very important in sports performance as it can boost glycogen storage, reduce muscle soreness and promote muscle repair. For those who are active regularly, there may be benefit from consuming a portion of protein at each mealtime and spreading protein intake throughout the day. Protein intake that exceeds the recommended daily allowance is widely accepted

for both endurance and power athletes. The various techniques utilized to rate protein will be discussed. Traditionally, sources of dietary protein are seen as either being of animal or vegetable origin. Animal sources provide a complete source of protein (i.e. containing all essential amino acids), whereas vegetable sources generally lack one or more of the essential amino acids. Animal sources of dietary protein, despite providing a complete protein and numerous vitamins and minerals, have some health professionals concerned about the amount of saturated fat common in these foods compared to vegetable sources. The advent of processing techniques has shifted some of this attention and ignited the sports supplement marketplace with derivative products such as whey, casein and soy. Individually, these products vary in quality and applicability to certain populations.

**The following questions are multiple choice questions. Choose the most appropriate answer:**

- (A) Correct statement about amino acids is-
- (a) All amino acids are optically active
  - (b) All amino acids except glycine are optically active.
  - (c) All amino acids except glutamic acid are optically active.
  - (d) All amino acids except lysine are optically active.
- (B) Proteins are found to have two different types of secondary structures viz.  $\alpha$ -helix and  $\beta$ -pleated sheet structure.  $\alpha$ -helix structure of protein is stabilised by:
- (a) Peptide bonds
  - (b) van der Waals forces
  - (c) Hydrogen bonds
  - (d) Dipole-dipole interactions
- (C) Example of Globular proteins is -
- (a) Myosin
  - (b) Albumin
  - (c) Collagen
  - (d) Fibroin
- (D) Which of the statements about denaturation given below are correct?
- (1) Denaturation of proteins causes loss of secondary and tertiary structures of the protein.
  - (2) Denaturation leads to the conversion of double strand of DNA into single strand.
  - (3) Denaturation affects primary structure which gets distorted.
- (a) (2) and (3)
  - (b) (1) and (3)
  - (c) (1) and (2)
  - (d) (1), (2) and (3)

**3. Read the passage given below and answer the following questions:**

The particles in the nucleus of cell, responsible for heredity, are called chromosomes which are made up of proteins and another type of biomolecules called nucleic acids. Nucleic acids are long chain polymers of nucleotides. Nucleotides are low molecular weight intracellular compounds that play major roles in physiological and biological functions, They act as precursors for nucleic acid synthesis and are also fundamental for intermediary metabolism. The two types of nucleic acids found in the chromosomes of cells of mammals are called 'deoxyribonucleic acid' and 'ribonucleic acid'. They are usually abbreviated as DNA and RNA respectively. As they are found in the nucleus of cells, they are called nucleic acids. Nucleotides and nucleic acids turn over rapidly, especially in growing tissues or those undergoing constant cell renewal. Tissues that grow have a net formation of new DNA and a rapid turnover of RNA. Nucleotides consists of a nitrogenous base (purine or pyrimidine), a pentose (ribose or deoxyribose), and one or more phosphate groups. The nitrogenous bases are derived from two parent heterocyclic molecules. The major purines found in living organisms are adenine and guanine, while cytosine, thymine, and uracil are the major pyrimidine bases. Nitrogenous bases can be formed from amino acid precursors or reutilized after their release from nucleic acid breakdown via the salvage pathway. The purine ring carbon atoms formed from the dispensable amino acids glycine, glutamic acid, and aspartame. The carbon atoms pyrimidines are derived from carbamoyl phosphate and aspartame. It has been concluded that there are about six billion base pairs in the DNA of a single human cell.

**The following questions are multiple choice questions. Choose the most appropriate answer:**

- (A) Dinucleotide is obtained by joining two nucleotides together by phosphodiester linkage. Between which carbon atoms of pentose sugars of nucleotides are these linkages present?
- |               |               |
|---------------|---------------|
| (a) 5' and 3' | (b) 1' and 5' |
| (c) 5' and 5' | (d) 3' and 3' |
- (B) In DNA, the complementary bases are:
- Uracil and adenine; cytosine and guanine
  - Adenine and thymine; guanine and cytosine.
  - Adenine and thymine; guanine and uracil
  - Adenine and guanine; thymine and cytosine.



- (C). The correct statement regarding RNA and DNA is:
- (a) The sugar component RNA is arabinose and sugar in DNA is ribose
  - (b) The sugar component in RNA is 2rdeoxyribose and the sugar component in DNA is arabinose.
  - (c) The sugar component in RNA is arabinose and the sugar component in DNA is 2'-deoxyribose.
  - (d) The sugar component in RNA is ribose and sugar component in DNA is 2' deoxyribose
- (D). Which one of the following is not present in RNA ?
- (a) Uracil
  - (b) Ribose
  - (c) Thymine
  - (d) Phosphate

**4. Read the passage given below and answer the following questions:**

Adenosine triphosphate (ATP) is the energy-carrying molecule found in the cells of all living things. ATP captures chemical energy obtained from the breakdown of food molecules and releases it to fuel other cellular processes. ATP is a nucleotide that consists of three main structures: the nitrogenous base, adenine; the sugar, ribose; and a chain of three phosphate groups bound to ribose. The phosphate tail of ATP is the actual power source which the cell taps. Available energy is contained in the bonds between the phosphates and is released when they are broken, which occurs through the addition of a water molecule (a process called hydrolysis). Usually only the outer phosphate is removed from ATP to yield energy; when this occurs ATP is converted to adenosine diphosphate (ADP), the form of the nucleotide having only two phosphates. The importance of ATP (adenosine triphosphate) as the main source of chemical energy in living matter and its involvement in cellular processes has long been recognized. The primary mechanism whereby higher organisms, including humans, generate ATP is through mitochondrial oxidative phosphorylation. For the majority of organs, the main metabolic fuel is glucose, which in the presence of oxygen undergoes complete combustion to  $\text{CO}_2$  and  $\text{H}_2\text{O}$ :



The free energy ( $\Delta G$ ) liberated in this exergonic ( $\Delta G$  is negative) reaction is partially trapped as ATP in two consecutive processes: glycolysis (cytosol) and oxidative phosphorylation (mitochondria). The first produces 2 mol of ATP per mol of glucose, and the second 36 mol of ATP per mol of glucose. Thus, oxidative phosphorylation yields 17-18 times as much useful energy in the form of ATP as can be obtained from the same amount of glucose by glycolysis alone. The efficiency of glucose metabolism is the ratio of amount of energy produced when 1 mol of glucose oxidised in cell to the enthalpy of combustion of glucose. The energy lost in the process is in the form of heat. This heat is responsible for keeping warm.

**Reference :** Erecinska', M., & Silvet, I. k..(1989). TP and Brain Function Journal of Cerebral Blood Flow & Metabolism, 9(1), 2-19.

<https://doi.org/10.10381jcbfm.1989.2> and

<https://www.britannica.com/science/adenosine-triphosphate>

- (A) Cellular oxidation of glucose is a:
- (a) spontaneous and endothermic process
  - (b) non spontaneous and exothermic process
  - (c) non spontaneous and endothermic process
  - (d) spontaneous and exothermic process
- (B) What is the efficiency of glucose metabolism if 1 mole of glucose gives, 38ATP energy?(Given: The enthalpy of combustion of glucose is 686 kcal, 1ATP= 7.3kcal)
- (a) 100%
  - (b) 38%
  - (c) 62%
  - (d) 80%
- (C) Which of the following statement is true?
- (a) ATP is a nucleoside made up of nitrogenous base adenine and ribose sugar.
  - (b) ATP consists the nitrogenous base, adenine and the sugar, deoxyribose.
  - (c) ATP is a nucleotide which contains a chain of three phosphate groups bound to ribose sugar.
  - (d) The nitrogenous base of ATP is the actual power source.
- (D) Nearly 95% of the energy released during cellular respiration is due to:
- (a) glycolysis occurring in cytosol
  - (b) oxidative phosphorylation.
  - (c) glycolysis occurring in mitochondria
  - (d) oxidative phosphorylation occurring in mitochondria
- (E) Which of the following statements is correct?
- (a) ATP is a nucleotide which has three phosphate groups while ADP is a nucleoside which has three phosphate groups.
  - (b) ADP contains a nitrogenous bases adenine, ribose sugar and two phosphate groups bound to rihose.
  - (c) ADP is the main source of chemical energy in living matter.
  - (d) ATP and ADP are nucleosides which differ in number of phosphate groups.

## 5. Read the passage and answer the following questions:

## EVIDENCE FOR THE FIBROUS NATURE OF DNA

The basic chemical formula of DNA is now well established. It consists of a very long chain, the backbone of which is made up of alternate sugar and phosphate groups, joined together in regular 3' 5' phosphate di-ester linkages. To each sugar is attached a nitrogenous base, only four different kinds of which are commonly found in DNA. Two of these---adenine and guanine--- are purines, and the other two thymine and cytosine---are pyrimidines. A fifth base, 5-methyl cytosine, occurs in smaller amounts in certain organisms, and a sixth, 5-hydroxy-methyl-cytosine, is found instead of cytosine in the T even phages. It should be noted that the chain is unbranched, a consequence of the regular internucleotide linkage. On the other hand the sequence of the different nucleotides is, as far as can be ascertained, completely irregular. Thus, DNA has some features which are regular, and some which are irregular. A similar conception of the DNA molecule as a long thin fiber is obtained from physicochemical analysis involving sedimentation, diffusion, light scattering, and viscosity measurements. These techniques indicate that DNA is a very asymmetrical structure approximately 20 Å wide and many thousands of angstroms long. Estimates of its molecular weight currently center between  $5 \times 10^6$  and  $10^7$  (approximately  $3 \times 10^4$  nucleotides). Surprisingly each of these measurements tend to suggest that the DN is relatively rigid, a puzzling finding in view of the large number of single bonds (5 per nucleotide) in the phosphate-sugar backbone. Recently these indirect inferences have been confirmed by electron microscopy.

**Reference:** Watson, J. D., & Crick, F. H. (1953, January). The structure of DNA In Cold spring Harbor symposia on quantitative biology (Vol. 18, pp. 123-131) Cold Spring Harbor Laboratory Press.

(A) Purines present in DNA are:

- (a) adenine and thymine
- (b) guanine and thymine
- (c) cytosine and thymine
- (d) adenine and guanine

(B) DNA molecule has \_\_\_\_\_ internucleotide linkage and Purines sequence of the different nucleotides

- (a) regular, regular
- (b) regular, irregular
- (c) irregular, regular
- (d) irregular, irregular

(C) DNA has a \_\_\_\_\_ backbone.

- (a) phosphate -purine
- (b) pyrimidines- sugar
- (c) phosphate- sugar
- (d) purine- pyrimidine

(D) Out of the four different kinds of nitrogenous bases which are commonly found in DNA, \_\_\_\_\_ has been replaced in some organisms.

- (a) adenine
- (b) guanine
- (c) cytosine
- (d) thymine

## ANSWERS

### **I MULTIPLE CHOICE QUESTIONS**

- I 1. (b) 2. (b) 3. (a) 4. (c) 5. (a) 6. (c) 7. (c) 8. (b) 9. (b) 10. (a) 11. (c) 12. (c) 13. (c) 14. (c) 15. (b) 16. (a)-(q, s), (b)-(r) (c)-(q), (d)-(p) 17. (a)-(p, q), (b)-(p), (c)-(s), (d)-(r) 18. b 19. (d) 20. (c)

### **II FILL IN THE BLANKS**

1. Amylose 2. n-Hexane 3.  $\alpha$ -D-glucose 4. Adenine and guanine  
5. Peptide linkage. 6. Vitamin B<sub>12</sub> 7. Uracil 8. Vitamin K. 9. Gluconic acid  
10. Glycogen.

### **III ASSERTION REASON TYPE QUESTIONS**

1. Vitamin B 2. Vitamin C 3. Cellulose 4. Lactose 5. Starch 6. Polysaccharides  
7. Reducing sugars 8. vitamin A 9. Nucleoside, nucleotide 10. Anomers

### **IV ONE WORD ANSWER TYPE QUESTIONS**

1. (c) 2. (a) 3. (c) 4. (a) 5. (c) 6. (a) 7. (b) 8. (a) 9. (b) 10. (d)

### **CASE STUDY BASED QUESTIONS**

1. A. (c) B. (b) C. (d) D. (b)  
2. A. (b) B. (c) C. (b) D. (c)  
3. A. (b) B. (b) C. (d) D. (c)  
4. A. (d) B. (b) C. (c) D. (d) E. (b)  
5. A. (d) B. (b) C. (c) D. (c)

**UNIT TEST-I**  
**Biomolecules**

**Maximum Marks : 20****Time Allowed : 1 Hours**

1. Name polysaccharide which is stored in the liver of animals. (1)
2. Name the enantiomer of D-glucose. (1)
3. Why is sucrose called invert sugar? (1)
4. Name the building blocks of proteins. (1)
5. Give the structure of simplest optically active amino acid. (1)
6. What are anomers? Give the structures of two anomers of glucose. (2)
7. Write the hydrolysed products of (2)  
(i) maltose (ii) cellulose
8. What are vitamins? How are they classified? (2)
9. What do you understand by following: (3)  
(i) denaturation of protein  
(ii) specificity of an enzyme
10. Differentiate between the following (3)  
(i) secondary and tertiary structure of protein.  
(ii)  $\alpha$ -helix and  $\beta$ -pleated sheet structure of protein.  
(iii) fibrous and globular proteins.
11. (i) Name four bases present in DNA. (3)  
(ii) Which of them is not present in RNA?  
(iii) Give the structure of a nucleotide of DNA.

**UNIT TEST-II****Biomolecules****Maximum Marks : 20****Time Allowed : 1 Hours**

1. Name following: (1)
  - (i) Nitrogenous Base present in RNA but not in DNA.
  - (ii) Optically inactive amino acid
2. Explain- Amino acids shows amphoteric behaviour. (1)
3. Give two examples of disachharides. (1)
4. The linkage responsible for stability of primary structure of protein is (1)
5. Name the diseases associated with the deficiency of vitamin A and vitamin B<sub>12</sub>. (1)
6. Write any three reactions which cannot be explained by linear structure of D-glucose. Write cyclic structure of glucose. (2)
7. Discuss the tertiary and quaternary structure of proteins. Name the intermolecular forces responsible for stability of these structures. (2)
8. Explain- denaturation of proteins. Discuss its effect on the primary, secondary, tertiary and quaternary structures of proteins. (2)
9. Differentiate following pairs giving examples: (3)
  - (i) Reducing and non-reducing sugars
  - ii) Essential and non-essential amino acids
  - (iii) Fibrous and globular proteins
10. Write the reactions of D-glucose with following: (3)
  - (i) HI/ $\Delta$
  - (ii) Bromine water
  - (iii) Acetic anhydride
11. Give reasons for following: (3)
  - (i) Amino acids are soluble in water.
  - (ii) Tryptophan is required in diet regularly but glutamic acid does not.
  - (iii) DNA is more stable than RNA.

**CBSE SAMPLE PAPER**  
**CHEMISTRY THEORY**  
**(043)**

**MM:70****Time: 3 hours****General Instructions:****Read the following instructions carefully.**


- a) There are 35 questions in this question paper with internal choice.
- b) SECTION A consists of 18 multiple-choice questions carrying 1 mark each.
- c) SECTION B consists of 7 very short answer questions carrying 2 marks each.
- d) SECTION C consists of 5 short answer questions carrying 3 marks each.
- e) SECTION D consists of 2 case-based questions carrying 4 marks each.
- f) SECTION E consists of 3 long answer questions carrying 5 marks each.
- g) All questions are compulsory.
- h) Use of log tables and calculators is not allowed

**SECTION A**

The following questions are multiple-choice questions with one correct answer. Each question carries 1 mark. There is no internal choice in this section.

1. The major product of acid catalysed dehydration of 1-methylcyclohexanol is:
  - a. 1-methylcyclohexane
  - b. 1-methylcyclohexene
  - c. 1-cyclohexylmethanol
  - d. 1-methylenecyclohexane
2. Which one of the following compounds is more reactive towards  $\text{S}_{\text{N}}1$  reaction?
  - a.  $\text{CH}_2=\text{CHCH}_2\text{Br}$
  - b.  $\text{C}_6\text{H}_5\text{CH}_2\text{Br}$
  - c.  $\text{C}_6\text{H}_5\text{CH}(\text{C}_6\text{H}_5)\text{Br}$
  - d.  $\text{C}_6\text{H}_5\text{CH}(\text{CH}_3)\text{Br}$
3.  $\text{KMnO}_4$  is coloured due to:
  - a. d-d transitions
  - b. charge transfer from ligand to metal
  - c. unpaired electrons in d orbital of Mn
  - d. charge transfer from metal to ligand

4. Which radioactive isotope would have the longer half-life  $^{15}\text{O}$  or  $^{19}\text{O}$ ? (Given rate constants for  $^{15}\text{O}$  and  $^{19}\text{O}$  are  $5.63 \times 10^{-3} \text{ s}^{-1}$  and  $k = 2.38 \times 10^{-2} \text{ s}^{-1}$  respectively.)
- $^{15}\text{O}$
  - $^{19}\text{O}$
  - Both will have the same half-life
  - None of the above, information given is insufficient
5. The molar conductivity of  $\text{CH}_3\text{COOH}$  at infinite dilution is  $390 \text{ Scm}^2/\text{mol}$ . Using the graph and given information, the molar conductivity of  $\text{CH}_3\text{COOK}$  will be:



- $100 \text{ Scm}^2/\text{mol}$
- $115 \text{ Scm}^2/\text{mol}$
- $150 \text{ Scm}^2/\text{mol}$
- $125 \text{ Scm}^2/\text{mol}$


**\*FOR VISUALLY CHALLENGED LEARNERS**

5. What is the molar conductance at infinite dilution for sodium chloride if the molar conductance at infinite dilution of  $\text{Na}^+$  and  $\text{Cl}^-$  ions are  $51.12 \times 10^{-4} \text{ Scm}^2/\text{mol}$  and  $73.54 \times 10^{-4} \text{ Scm}^2/\text{mol}$  respectively?
- $124.66 \text{ Scm}^2/\text{mol}$
  - $22.42 \text{ Scm}^2/\text{mol}$
  - $198.20 \text{ Scm}^2/\text{mol}$
  - $175.78 \text{ Scm}^2/\text{mol}$



6. For the reaction,  $A + 2B \rightarrow AB_2$ , the order w.r.t. reactant A is 2 and w.r.t. reactant B. What will be change in rate of reaction if the concentration of A is doubled and B is halved?
- increases four times
  - decreases four times
  - increases two times
  - no change
7. Arrange the following in the increasing order of their boiling points:  
A: Butanamine, B: N,N-Dimethylethanamine, C: N-Ethylethanamine
- $C < B < A$
  - $A < B < C$
  - $A < C < B$
  - $B < C < A$
8. The CFSE of  $[\text{CoCl}_6]^{3-}$  is  $18000 \text{ cm}^{-1}$  the CFSE for  $[\text{CoCl}_4]^-$  will be:
- $18000 \text{ cm}^{-1}$
  - $8000 \text{ cm}^{-1}$
  - $2000 \text{ cm}^{-1}$
  - $16000 \text{ cm}^{-1}$
9. What would be the major product of the following reaction?  
 $\text{C}_6\text{H}_5\text{-CH}_2\text{-OC}_6\text{H}_5 + \text{HBr} \rightarrow \text{A} + \text{B}$
- $\text{A} = \text{C}_6\text{H}_5\text{CH}_2\text{OH}$ ,  $\text{B} = \text{C}_6\text{H}_6$
  - $\text{A} = \text{C}_6\text{H}_5\text{CH}_2\text{OH}$ ,  $\text{B} = \text{C}_6\text{H}_5\text{Br}$
  - $\text{A} = \text{C}_6\text{H}_5\text{CH}_3$ ,  $\text{B} = \text{C}_6\text{H}_5\text{Br}$
  - $\text{A} = \text{C}_6\text{H}_5\text{CH}_2\text{Br}$ ,  $\text{B} = \text{C}_6\text{H}_5\text{OH}$
10. Which of the following statements is not correct for amines?
- Most alkyl amines are more basic than ammonia solution.
  - $\text{pK}_b$  value of ethylamine is lower than benzylamine.
  - $\text{CH}_3\text{NH}_2$  on reaction with nitrous acid releases  $\text{NO}_2$  gas.
  - Hinsberg's reagent reacts with secondary amines to form sulphonamides.
11. Which of the following tests/ reactions is given by aldehydes as well as ketones?
- Fehling's test
  - Tollen's test
  - 2,4 DNP test
  - Cannizzaro reaction

12. Arrhenius equation can be represented graphically as follows:



The (i) intercept and (ii) slope of the graph are:

- |                |               |
|----------------|---------------|
| a. (i) $\ln A$ | (ii) $E_a/R$  |
| b. (i) $A$     | (ii) $E_a$    |
| c. (i) $\ln A$ | (ii) $-E_a/R$ |
| d. (i) $A$     | (ii) $-E_a$   |

**\*FOR VISUALLY CHALLENGED LEARNERS**

- \*12. The unit of rate constant for the reaction



which has rate  $= k[A]^2[B]$  is:

- |   |
|---|
| a. $\text{mol L}^{-1} \text{s}^{-1}$          |
| b. $\text{s}^{-1}$                            |
| c. $\text{mol L}^{-1}$                        |
| d. $\text{mol}^{-2} \text{L}^2 \text{s}^{-1}$ |
13. The number of ions formed on dissolving one molecule of  $\text{FeSO}_4 \cdot (\text{NH}_4)_2 \text{SO}_4 \cdot 6\text{H}_2\text{O}$  in water is:
- |      |
|------|
| a. 3 |
| b. 4 |
| c. 5 |
| d. 6 |
14. The oxidation of toluene to benzaldehyde by chromyl chloride is called
- |                            |
|----------------------------|
| a. Etard reaction          |
| b. Riemer-Tiemann reaction |
| c. Stephen's reaction      |
| d. Cannizzaro's reaction   |
15. Given below are two statements labelled as Assertion (A) and Reason (R)

**Assertion (A):** An ether is more volatile than an alcohol of comparable molecular mass.

**Reason (R):** Ethers are polar in nature.

Select the most appropriate answer from the options given below:

- a. Both A and R are true and R is the correct explanation of A
- b. Both A and R are true but R is not the correct explanation of A.
- c. A is true but R is false.
- d. A is false but R is true.

16. Given below are two statements labelled as Assertion (A) and Reason (R)

**Assertion (A):** Proteins are found to have two different types of secondary structures viz alpha-helix and beta-pleated sheet structure.

**Reason (R):** The secondary structure of proteins is stabilized by hydrogen bonding.

Select the most appropriate answer from the options given below:

- a. Both A and R are true and R is the correct explanation of A
- b. Both A and R are true but R is not the correct explanation of A.
- c. A is true but R is false.
- d. A is false but R is true.

17. Given below are two statements labelled as Assertion (A) and Reason (R)

**Assertion :** Magnetic moment values of actinides are lesser than the theoretically predicted values.

**Reason :** Actinide elements are strongly paramagnetic.

Select the most appropriate answer from the options given below:

- a. Both A and R are true and R is the correct explanation of A
- b. Both A and R are true but R is not the correct explanation of A.
- c. A is true but R is false.
- d. A is false but R is true.

18. Given below are two statements labelled as Assertion (A) and Reason (R)

**Assertion (A):** Tertiary amines are more basic than corresponding secondary and primary amines in gaseous state.

**Reason (R):** Tertiary amines have three alkyl groups which cause +I effect. Select the most appropriate answer from the options given below:

- a. Both A and R are true and R is the correct explanation of A
- b. Both A and R are true but R is not the correct explanation of A.
- c. A is true but R is false.
- d. A is false but R is true.

**SECTION B**

This section contains 7 questions with internal choice in two questions. The following questions are very short answer type and carry 2 marks each.

19. A first-order reaction takes 69.3 min for 50% completion. What is the time needed for 80% of the reaction to get completed?  
(Given:  $\log 5 = 0.6990$ ,  $\log 8 = 0.9030$ ,  $\log 2 = 0.3010$ )
20. Account for the following:
  - a. There are 5 OH groups in glucose
  - b. Glucose is a reducing sugar

**OR**

What happens when D – glucose is treated with the following reagents

- a. Bromine water
- b.  $\text{HNO}_3$
21. Give reason for the following:
  - a. During the electrophilic substitution reaction of haloarenes, para substituted derivative is the major product.
  - b. The product formed during  $\text{S}_{\text{N}}1$  reaction is a racemic mixture.

**OR**

- a. Name the suitable alcohol and reagent, from which 2-Chloro-2-methyl propane can be prepared.
- b. Out of the Chloromethane and Fluoromethane, which one has higher dipole moment and why?
22. The formula  $\text{Co}(\text{NH}_3)_5\text{CO}_3\text{Cl}$  could represent a carbonate or a chloride. Write the structures and names of possible isomers.
23. Corrosion is an electrochemical phenomenon. The oxygen in moist air reacts as follows:  

$$\text{O}_2(\text{g}) + 2\text{H}_2\text{O}(\text{l}) + 4\text{e}^- \rightarrow 4\text{OH}^-(\text{aq}).$$

Write down the possible reactions for corrosion of zinc occurring at anode, cathode, and overall reaction to form a white layer of zinc hydroxide.

24. Explain how and why will the rate of reaction for a given reaction be affected when
  - a. a catalyst is added
  - b. the temperature at which the reaction was taking place is decreased
25. Write the reaction and IUPAC name of the product formed when 2-Methylpropanal (isobutyraldehyde) is treated with ethyl magnesium bromide followed by hydrolysis.

## SECTION C

This section contains 5 questions with internal choice in two questions. The following questions are short answer type and carry 3 marks each.

26. Write the equations for the following reaction:
  - a. Salicylic acid is treated with acetic anhydride in the presence of conc.  $\text{H}_2\text{SO}_4$
  - b. Tert butyl chloride is treated with sodium ethoxide.
  - c. Phenol is treated with chloroform in the presence of NaOH
27. Using Valence bond theory, explain the following in relation to the paramagnetic complex  $[\text{Mn}(\text{CN})_6]^{3-}$ 
  - a. type of hybridization
  - b. magnetic moment value
  - c. type of complex – inner, outer orbital complex
28. Answer the following questions:
  - a. State Henry's law and explain why are the tanks used by scuba divers filled with air diluted with helium (11.7% helium, 56.2% nitrogen and 32.1% oxygen)?
  - b. Assume that argon exerts a partial pressure of 6 bar. Calculate the solubility of argon gas in water. (Given Henry's law constant for argon dissolved in water,  $K_{\text{H}} = 40 \text{ kbar}$ )
29. Give reasons for any 3 of the following observations:
  - a. Aniline is acetylated before nitration reaction.
  - b.  $\text{pK}_{\text{b}}$  of aniline is lower than the m-nitroaniline.
  - c. Primary amine on treatment with benzenesulphonyl chloride forms a product which is soluble in NaOH however secondary amine gives product which is insoluble in NaOH.
  - d. Aniline does not react with methyl chloride in the presence of anhydrous  $\text{AlCl}_3$  catalyst.
30.
  - a. Identify the major product formed when 2-cyclohexylchloroethane undergoes a dehydrohalogenation reaction. Name the reagent which is used to carry out the reaction.
  - b. Why are haloalkanes more reactive towards nucleophilic substitution reactions than haloarenes and vinylic halides?

## OR

- a. Name the possible alkenes which will yield 1-chloro-1-methylcyclohexane on their reaction with HCl. Write the reactions involved.
- b. Allyl chloride is hydrolysed more readily than n-propyl chloride. Why?

### SECTION D

The following questions are case-based questions. Each question has an internal choice and carries 4 (1+1+2) marks each. Read the passage carefully and answer the questions that follow.

31. Strengthening the Foundation: Chargaff Formulates His "Rules"

Many people believe that James Watson and Francis Crick discovered DNA in the 1950s. In reality, this is not the case. Rather, DNA was first identified in the late 1860s by Swiss chemist Friedrich Miescher. Then, in the decades following Miescher's discovery, other scientists--notably, Phoebus Levene and Erwin Chargaff--carried out a series of research efforts that revealed additional details about the DNA molecule, including its primary chemical components and the ways in which they joined with one another. Without the scientific foundation provided by these pioneers, Watson and Crick may never have reached their groundbreaking conclusion of 1953: that the DNA molecule exists in the form of a three-dimensional double helix.

Chargaff, an Austrian biochemist, as his first step in this DNA research, set out to see whether there were any differences in DNA among different species. After developing a new paper chromatography method for separating and identifying small amounts of organic material, Chargaff reached two major conclusions:

- (i) the nucleotide composition of DNA varies among species.
- (ii) Almost all DNA, no matter what organism or tissue type it comes from maintains certain properties, even as its composition varies. In particular, the amount of adenine (A) is similar to the amount of thymine (T), and the amount of guanine (G) approximates the amount of cytosine (C). In other words, the total amount of purines (A + G) and the total amount of pyrimidines (C + T) are usually nearly equal. This conclusion is now known as "Chargaff's rule."

Chargaff's rule is not obeyed in some viruses. These either have single- stranded DNA or RNA as their genetic material.

**Answer the following questions:**

- a. A segment of DNA has 100 adenine and 150 cytosine bases. What is the total number of nucleotides present in this segment of DNA?
- b. A sample of hair and blood was found at two sites. Scientists claim that the samples belong to same species. How did the scientists arrive at this conclusion?
- c. The sample of a virus was tested and it was found to contain 20% adenine, 20% thymine, 20 % guanine and the rest cytosine. Is the genetic material of this virus (a) DNA- double helix (b) DNA-single helix (c) RNA? What do you infer from this data?

**OR**

How can Chargaff's rule be used to infer that the genetic material of an organism is double-helix or single-helix?

32. Henna is investigating the melting point of different salt solutions.

She makes a salt solution using 10 mL of water with a known mass of NaCl salt. She puts the salt solution into a freezer and leaves it to freeze.

She takes the frozen salt solution out of the freezer and measures the temperature when the frozen salt solution melts.

She repeats each experiment.

S.No.	Mass of the salt used in g	Melting point $^{\circ}\text{C}$	
		Reading Set 1	Reading Set 2
1	0.3	-1.9	-1.9
2	0.4	-2.5	-2.6
3	0.5	-3.0	-5.5
4	0.6	-3.8	-3.8
5	0.8	-5.1	-5.0
6	1.0	-6.4	-6.3

**Assuming the melting point of pure water as  $0^{\circ}\text{C}$ , answer the following questions:**

- One temperature in the second set of results does not fit the pattern. Which temperature is that? Justify your answer.
- Why did Henna collect two sets of results?
- In place of NaCl, if Henna had used glucose, what would have been the melting point of the solution with 0.6 g glucose in it?

**OR**

What is the predicted melting point if 1.2 g of salt is added to 10 mL of water? Justify your answer.


**SECTION E**

The following questions are long answer type and carry 5 marks each. Two questions have an internal choice.

33.
  - Why does the cell voltage of a mercury cell remain constant during its lifetime?
  - Write the reaction occurring at anode and cathode and the products of electrolysis of aq KCl.
  - What is the pH of HCl solution when the hydrogen gas electrode shows a potential of  $-0.59\text{ V}$  at standard temperature and pressure?

**OR**

- Molar conductivity of substance "A" is  $5.9 \times 10^3 \text{ S/m}$  and "B" is  $1 \times 10^{-16} \text{ S/m}$ . Which of the two is most likely to be copper metal and why?
- What is the quantity of electricity in Coulombs required to produce 4.8 g of Mg from molten  $\text{MgCl}_2$ ? How much Ca will be produced if the same amount of electricity was passed through molten  $\text{CaCl}_2$ ? (Atomic mass of Mg = 24 u, atomic mass of Ca = 40 u).
- What is the standard free energy change for the following reaction at room temperature? Is the reaction spontaneous?



- A hydrocarbon (A) with molecular formula  $\text{C}_5\text{H}_{10}$  on ozonolysis gives two products (B) and (C). Both (B) and (C) give a yellow precipitate when heated with iodine in presence of NaOH while only (B) give a silver mirror on reaction with Tollen's reagent.
  - Identify (A), (B) and (C).
  - Write the reaction of B with Tollen's reagent
  - Write the equation for iodoform test for C
  - Write down the equation for aldol condensation reaction of B and C.

**OR**

An organic compound (A) with molecular formula  $\text{C}_2\text{Cl}_3\text{O}_2\text{H}$  is obtained when (B) reacts with Red P and  $\text{Cl}_2$ . The organic compound (B) can be obtained on the reaction of methyl magnesium chloride with dry ice followed by acid hydrolysis.

- Identify A and B
  - Write down the reaction for the formation of A from B. What is this reaction called?
  - Give any one method by which organic compound B can be prepared from its corresponding acid chloride.
  - Which will be the more acidic compound (A) or (B)? Why?
  - Write down the reaction to prepare methane from the compound (B).
- Answer the following:
    - Why are all copper halides known except that copper iodide?
    - Why is the  $E^\circ_{(\text{V}^{3+}/\text{V}^{2+})}$  value for vanadium comparatively low?
    - Why HCl should not be used for potassium permanganate titrations?



## MARKING SCHEME

## SECTION A

Q1 to 18 each correct answer 1 mark

1. b. 1-methylcyclohexene  
According to Saytzeff rule i.e highly substituted alkene is major product. Here dehydration reaction takes place, alkene is formed due to the removal of a water molecule.
2. c.  $\text{C}_6\text{H}_5\text{CH}(\text{C}_6\text{H}_5)\text{Br}$   $\text{C}_6\text{H}_5\text{CH}(\text{C}_6\text{H}_5)^+$  carbocation formed is more stable
3. b. charge transfer from ligand to metal  
The Mn atom in  $\text{KMnO}_4$  has +7 oxidation state with electron configuration  $[\text{Ar}]3d^0 4s^0$ . Since no unpaired electrons are present, d-d transitions are not possible. The molecule should, therefore, be colourless.  
Its intense purple due to  $\text{L} \rightarrow \text{M}$  (ligand to metal) charge transfer  $2p(\text{L})$  of O to  $3d(\text{M})$  of Mn.
4. a.  $^{15}\text{O}$   
The rate constant for the decay of O-15 is less than that for O-19. Therefore, the rate of decay of O-15 will be slower and will have a longer half life.
5. b.  $115 \text{ Scm}^2/\text{mol}$   $\Delta^\circ \text{CH}_3\text{COOK} = \Delta^\circ \text{CH}_3\text{COOH} + \Delta^\circ \text{OKCl} - \Delta^\circ \text{HCl} = 390 + 150 - 425 = 115 \text{ Scm}^2/\text{mol}$
- 5\* (For visually challenged learners)  
a.  $124.66 \times 10^{-4} \text{ Sm}^2 \text{mol}^{-1}$   
Molar conductance of  $\text{NaCl} = \lambda^+ \text{Na} + \lambda^-_{\text{Cl}}$   
 $= 51.12 \times 10^{-4} + 73.54 \times 10^{-4} = 124.66 \times 10^{-4} \text{ Sm}^2 \text{mol}^{-1}$
6. a. increases 4 times  
 $\text{Rate} = [\text{A}]^2$   
If  $[\text{A}]$  is doubled then  $\text{Rate}' = [2\text{A}]^2 = 4[\text{A}]^2 = 4 \text{ Rate}$
7. d.  $\text{B} < \text{C} < \text{A}$   
In primary amine intermolecular association due to H-bonding is maximum while in tertiary it is minimum.
8. b.  $8000 \text{ cm}^{-1}$   
 $\Delta t = (4/9) \times 18000 \text{ cm}^{-1} = 8000 \text{ cm}^{-1}$
9. d.  $\text{A} = \text{C}_6\text{H}_5\text{CH}_2\text{Br}$ ,  $\text{B} = \text{C}_6\text{H}_5\text{OH}$ ,  
 $\text{C}_6\text{H}_5\text{CH}_2\text{OC}_6\text{H}_5$   $\text{H} + \text{C}_6\text{H}_5\text{CH}_2\text{OC}_6\text{H}_5$
10. c.  $\text{CH}_3\text{NH}_2$  on reaction with nitrous acid releases  $\text{NO}_2$  gas Wrong statement .  
The evolution of nitrogen gas takes place.


11. c. 2,4 DNP test  
Fehling's, Tollen's and Cannizzao reaction is shown by alcohols only.
12. c. (i)ln A (ii) - Ea/R
- 12\* (For visually challenged learners)  
d.  $\text{mol}^{-2} \text{L}^2 \text{s}^{-1}$  since the order of reaction is 3.
13. c. 5 l  
 $\text{Fe}^{2+}$ ,  $2 \text{SO}_4^{2-}$  and  $2 \text{NH}_4^+$  ions
14. A Etard reaction
15. b Both A and R are true but R is not the correct explanation of A. A and R are two different statements about ethers  
The correct reason is that hydrogen bonding does not exist amongst ether molecules.
16. b Both A and R are true but R is not the correct explanation of A.
17. b Both A and R are true but R is not the correct explanation of A. The magnetic moment is less as the 5f electrons of actinides are less effectively shielded which results in quenching of orbital contributions, they are strongly paramagnetic due to presence of unpaired electrons
18. a Both A and R are true and R is the correct explanation of A.

### SECTION B


19. Half life  $t_{1/2} = 0.693 / k$   
 $k = 0.693 / 69.3 = 1/100 = 0.01 \text{ min}^{-1}$  (1/2)  
 For first order reaction  

$$k = \frac{2.303}{t} \log \frac{[\text{Ro}]}{[\text{R0}]}$$
 (1)  

$$t = \frac{2.303}{0.01} \log \frac{100}{20}$$
  
 $t = 230.3 \log 5$  ( $\log 5 = 0.6990$ )  
 $t = 160.9 \text{ min}$  (1/2)
20. a. Acetylation of glucose with acetic anhydride gives glucose pentaacetate which confirms the presence of five -OH groups. Since it exists as a stable compound, five -OH groups should be attached to different carbon atoms (1)




- b. Glucose reduces Fehlings reagent (1)



OR

- a. (1)



- b. (1)
- $$\begin{array}{c}
 \text{CHO} \\
 | \\
 (\text{CHOH})_4 \\
 | \\
 \text{CH}_2\text{OH}
 \end{array}
 \xrightarrow{\text{Oxidation}}
 \begin{array}{c}
 \text{COOH} \\
 | \\
 (\text{CHOH})_4 \\
 | \\
 \text{COOH}
 \end{array}
 \xleftarrow{\text{Oxidation}}
 \begin{array}{c}
 \text{COOH} \\
 | \\
 (\text{CHOH})_4 \\
 | \\
 \text{CH}_2\text{OH}
 \end{array}$$
- Saccharic acid                      Gluconic acid

21. a. At the ortho position, higher steric hindrance is there, hence para isomer is usually predominate and is obtained in the major amount. (1)
- b. During the  $\text{S}_{\text{N}}1$  mechanism, intermediate carbocation formed is  $\text{sp}^2$  hybridized and planar in nature. This allows the attack of nucleophile from either side of the plane resulting in a racemic mixture. (1)


OR

- a. Tert butyl alcohol or 2-methyl propan-2-ol using Lucas reagent, mixture of concHCl and  $\text{ZnCl}_2$  the reaction will follow the  $\text{S}_{\text{N}}1$  pathway. (1)
- b. Chloromethane is having higher dipole moment. Due to smaller size of fluorine the dipole moment of fluoromethane is comparatively lesser. (1)
22.  $[\text{Co}(\text{NH}_3)_5\text{CO}_3]\text{Cl}$  and  $[\text{Co}(\text{NH}_3)_5\text{Cl}]\text{CO}_3$  (1/2+1/2)
- Pentaaminecarbonatocobalt(III)chloride (1/2)
- Pentaaminechloridocobalt(III)carbonate (1/2)
23. Anode:  $\text{Zn(s)} - \text{Zn}^{2+}(\text{aq}) + 2\text{e}^-$  (1/2)
- Cathode:  $\text{O}_2(\text{g}) + 2\text{H}_2\text{O(l)} + 4\text{e}^- \rightarrow 4\text{OH}^-(\text{aq})$ . (1/2)
- Overall:  $2\text{Zn(s)} + \text{O}_2(\text{g}) + 2\text{H}_2\text{O(l)} - 2\text{Zn}^{2+}(\text{aq}) + 4\text{OH}^-(\text{aq})$
- $2\text{Zn(s)} + \text{O}_2(\text{g}) + 2\text{H}_2\text{O(l)} - 2\text{Zn(OH)}_2(\text{ppt})$  (1)

24. The rate of reaction will increase. The catalyst decreases the activation energy of the reaction therefore the reaction becomes faster. (1/2+1/2)
- b. The rate of reaction will decrease. At lower temperatures the kinetic energy of molecules decreases thereby the collisions decrease resulting in a lowering of rate of reaction. (1/2+1/2)
25.  $(\text{CH}_3)_2\text{CHCHO} + \text{C}_2\text{H}_5\text{MgBr} \xrightarrow{\text{dry ether}} (\text{CH}_3)_2\text{CHCH}(\text{C}_2\text{H}_5)(\text{OMgBr})$   
 $(\text{CH}_3)_2\text{CHCH}(\text{C}_2\text{H}_5)(\text{OMgBr}) \xrightarrow{\text{H}^+/\text{H}_2\text{O}} (\text{CH}_3)_2\text{CHCH}(\text{C}_2\text{H}_5)(\text{OH})$  (1)  
 2-Methylpentan-3-ol (1)


## SECTION C

26. (i) Aspirin is formed



- (ii)  $(\text{CH}_3)_3\text{CCl} \xrightarrow{\text{sodium ethoxide}} (\text{CH}_3)_2\text{C}=\text{CH}_2$  (1)  
 2methylpropene

- (iii) o-hydroxybenzaldehyde will be formed



27.  $[\text{Mn}(\text{CN})_6]^{3-}$   
 $\text{Mn} = [\text{Ar}] 3d^5 4s^2$   
 $\text{Mn}^{3+} = [\text{Ar}] 3d^4$   
 Mn (ground state)



Mn in +3 state



Mn in  $[\text{Mn}(\text{CN})_6]^{3-}$



$d^2sp^3$  hybridisation

xx are electrons donated by ligand  $\text{CN}^-$

Type of hybridization –  $d^2sp^3$  (1)

Magnetic moment value –  $\sqrt{n(n+2)} = \sqrt{2(2+2)} = 2.87 \text{ BM}$

( $n$  = no. of unpaired electrons) (1)

Type of complex – inner orbital (1)

28. a. Henry's law: the partial pressure of the gas in vapour phase ( $p$ ) is proportional to the mole fraction of the gas ( $x$ ) in the solution. (1)

The pressure underwater is high, so the solubility of gases in blood increases. When the diver comes to surface the pressure decreases so does the solubility causing bubbles of nitrogen in blood, to avoid this situation and maintain the same partial pressure of nitrogen underwater too, the dilution is done. (1)

b.  $p = K_H \cdot x$

mole fraction of argon in water  $x = p/k = 6/40 \times 10^3 = 1.5 \times 10^{-4}$  (1)

29. (any 3)

a. Aniline is acetylated, before nitration reaction in order to avoid formation of tarry oxidation products and protecting the amino group, so that  $p$ -nitro derivative can be obtained as major product. (1)

b.  $pK_b$  of aniline is lower than the  $m$ -nitro aniline. The basic strength of aniline is more than  $m$ -nitroaniline.  $pK_b$  value is inversely proportional to basic strength. Presence of Electron withdrawing group decrease basic strength. (1)

c. Due to the presence of acidic hydrogen in the  $N$ -alkylbenzenesulphonamide formed by the treatment of primary amines. (1)


d. Aniline does not react with methylchloride in the presence of  $AlCl_3$  catalyst, because aniline is a base and  $AlCl_3$  is Lewis acid which lead to formation of salt. (1)

30. a. The major product formed when 2-cyclohexylchloroethane undergoes dehydrohalogenation reaction is 1-cyclohexylethene. The reagent which is used to carry out the reaction is ethanolic  $KOH$ . (1+1)


b. Haloalkanes are more reactive than haloarenes and vinylic halides because of the presence of partial double bond character  $C-X$  bond in haloarenes and vinylic halides. Hence they do not undergo nucleophilic reactions easily. (1)

OR

- a. Methylenecyclohexane



- 1-Methylcyclohexene



- b. Allyl chloride shows high reactivity as the carbocation formed in the first step is stabilised by resonance while no such stabilisation of carbocation exists in the case of n-propyl chloride. (1)

### SECTION D

31. a. A = 100 so T = 100  
C = 150 so G = 150  
Total nucleotides = 100 + 100 + 150 + 150 = 500 (1)
- b. They studied the nucleotide composition of DNA. It was the same so they concluded that the samples belong to same species. (1)
- c. A = T = 20%  
But G is not equal to C so double helix is ruled out. (1/2)  
The bases pairs are ATGC and not AUGC so it is not RNA (1/2)  
The virus is a single helix DNA virus (1)

### OR

According to Charagaff rule, all double helix DNA will have the same amount of A and T as well as C will be same amount as G. If this is not the case then the helix is single stranded. (2)

32. The melting point of ice is the freezing point of water. We can use the depression in freezing point property in this case.
- a. 3<sup>rd</sup> reading for 0.5 g there has to be an increase in depression of freezing point and therefore decrease in freezing point so also decrease in melting point when amount of salt is increased but the trend is not followed on this case. (1)
- b. two sets of reading help to avoid error in data collection and give more objective data. (1)
- c.  $\Delta T_f(\text{glucose}) = \frac{1 \times K_f \times 0.6 \times 1000}{180 \times 10}$  (1/2)  
 $\Delta T_f(\text{NaCl}) = \frac{2 \times K_f \times 0.6 \times 1000}{58.5 \times 10}$  (1/2)  
 $3.8 = \frac{2 \times K_f \times 0.6 \times 1000}{58.5 \times 10}$   
 Divide equation 1 by 2  
 $\frac{\Delta T_f(\text{glucose})}{3.8} = \frac{58.5}{2 \times 180}$  (1/2)  
 $\Delta T_f(\text{glucose}) = 0.62$  Freezing point or Melting point = - 0.62°C (1/2)

OR

depression in freezing point is directly proportional to molality (mass of solute when the amount of solvent remains same) (1)

0.3 g depression is  $1.9^{\circ}\text{C}$

0.6 g depression is  $3.8^{\circ}\text{C}$

1.2 g depression will be  $3.8 \times 2 = 7.6^{\circ}\text{C}$  (1)

## SECTION E

33. The cell potential remains constant during its life as the overall reaction does not involve any ion in solution whose concentration can change during its life time. (1)

b.  $\text{KCl (aq)} \rightarrow \text{K}^+ \text{(aq)} + \text{Cl}^- \text{(aq)}$

cathode:  $\text{H}_2\text{O (l)} + \text{e}^- \rightarrow \frac{1}{2} \text{H}_2 \text{(g)} + \text{OH}^- \text{(aq)}$  (1/2)

anode:  $\text{Cl}^- \text{(aq)} \rightarrow \frac{1}{2} \text{Cl}_2 \text{(aq)} + \text{e}^-$  (1/2)

net reaction:

$\text{KCl (aq)} + \text{H}_2\text{O (l)} \rightarrow \text{K}^+ \text{(aq)} + \text{OH}^- \text{(aq)} + \frac{1}{2} \text{H}_2 \text{(g)} + \frac{1}{2} \text{Cl}_2 \text{(g)}$  (1)

c. Given, potential of hydrogen gas electrode =  $-0.59 \text{ V}$

Electrode reaction:  $\text{H}^+ + \text{e}^- \rightarrow \frac{1}{2} \text{H}_2$

Applying Nernst equation,

$$E(\text{H}^+/\text{H}_2) = E^\circ(\text{H}^+/\text{H}_2) - \frac{0.059}{n} \log \frac{[\text{H}_2]^{1/2}}{[\text{H}^+]} \quad (1)$$

$$E^\circ(\text{H}^+/\text{H}_2) = 0 \text{ V}$$

$$E(\text{H}^+/\text{H}_2) = -0.59 \text{ V} \quad \text{---}$$

$$n = 1$$

$$[\text{H}_2] = 1 \text{ bar}$$

$$-0.59 = 0 - 0.059 (-\log [\text{H}^+]) \quad (1/2)$$

$$-0.59 = -0.059 \text{pH}$$

$$\therefore \text{pH} = 10 \quad (1/2)$$

OR


a. "A" is copper, metals are conductors thus have high value of conductivity. (1)

b.  $\text{Mg}^{2+} + 2\text{e}^- \rightarrow \text{Mg}$

1 mole of magnesium ions gains two moles of electrons or  $2F$  to form 1 mole of Mg

24 g Mg requires  $2F$  electricity


$$4.8 \text{ g Mg requires } 2 \times 4.8/24 = 0.4 F = 0.4 \times 96500 = 38600 \text{ C} \quad (1)$$



$2F$  electricity is required to produce 1 mole = 40 g Ca

0.4 F electricity will produce 8 g Ca (1)

c.  $F = 96500 \text{ C}$ ,  $n = 2$ ,



$$E^\circ_{\text{cell}} = E^\circ_{\text{cathode}} - E^\circ_{\text{anode}}$$

$$= 0.15 - (-0.14) = 0.29\text{V} \quad (1)$$

$$\Delta G^\circ = -nFE^\circ_{\text{cell}}$$

$$= -2 \times 96500 \times 0.29 = 55970 \text{ J/mol} \quad (1)$$


34. A is an alkene

B is an aldehyde with  $-\text{CH}_3$  group

C is a methyl ketone



A:  $\text{CH}(\text{CH}_3)=\text{C}(\text{CH}_3)_2$  B:  $\text{CH}_3\text{CHO}$  C:  $\text{O}=\text{C}(\text{CH}_3)_2$  (1.5 = 1/2 each)



↓ heat



↓ heat



**OR**

a. (A):  $\text{CCl}_3\text{COOH}$  (B):  $\text{CH}_3\text{COOH}$  (1)

b.  $\text{CH}_3\text{COOH}$  (i) Red P /  $\text{Cl}_2$   $\text{CCl}_3\text{COOH}$ , Hell Volhard Zelinsky reaction



c. A will be more acidic due to presence of 3 Cl groups (electron withdrawing groups) which increase acidity of carboxylic acid. (1)

e.  $\text{CH}_3\text{COOH}$  (i) NaOH, CaO (ii) heat  $\text{CH}_4 + \text{Na}_2\text{CO}_3$  (1)

35. a.  $\text{Cu}^{2+}$  oxidizes iodide ion to iodine. (1)

b. The low value for V is related to the stability of  $\text{V}^{2+}$  (half-filled t<sub>2g</sub> level) (1)

c. Permanganate titrations in presence of hydrochloric acid are unsatisfactory since hydrochloric acid is oxidised to chlorine.

d. The d orbital is full with ten electrons and shield the electrons present in the higher s-orbital to a greater extent resulting in increase in size.

e. The chromates and dichromates are interconvertible in aqueous solution depending upon pH of the solution. Increasing the pH (in basic solution) of dichromate ions a colour change from orange to yellow is observed as dichromate ions change to chromate ions.



## UNSOLVED SAMPLE PAPER

## SECTION A

MM: 70

Time : 3 hours

## General Instructions:

Read the following instructions carefully.

- There are 35 questions in this question paper with internal choice.
- SECTION A consists of 18 multiple-choice questions carrying 1 mark each.
- SECTION B consists of 7 very short answer questions carrying 2 marks each.
- SECTION C consists of 5 short answer questions carrying 3 marks each.
- SECTION D consists of 2 case-based questions carrying 4 marks each.
- SECTION E consists of 3 long answer questions carrying 5 marks each.
- All questions are compulsory.**
- Use of log tables and calculators is not allowed.**

## SECTION A

The following questions are multiple-choice questions with one correct answer.

Each questions carries 1 mark. There is no internal choice in this section.

- 50 mL of an aqueous solution of glucose  $C_6H_{12}O_6$  (Molar mass: 180 g/mol) contains  $6.02 \times 10^{22}$  molecules. The concentration of the solution will be  
(A) 0.1 M  
(B) 0.2 M  
(C) 1.0 M  
(D) 2.0 M
- If the standard electrode potential of an electrode is greater than zero then we can infer that its?  
(A) reduced form is more stable compared to hydrogen gas.  
(B) oxidised form is more stable compared to hydrogen gas.  
(C) reduced and oxidised forms are equally stable  
(D) reduced form is less stable than the hydrogen gas.
- Total number of unpaired electrons present in  $Co^{3+}$  (Atomic number=27) is  
(A) 2  
(B) 7  
(C) 4  
(D) 5
- The incorrect statement about interstitial compounds is ?  
(A) They are chemically reactive  
(B) They are very hard  
(C) They retain metallic conductivity.  
(D) They have high melting point.
- The correct IUPAC name of  $CH_3-\overset{\overset{OH}{|}}{\underset{\underset{OH}{|}}{C}}-CH_2CH_3$  is  
(A) tert-butyl alcohol  
(B) 2,2- Dimethylpropanol  
(C) 2-Methylbutan-2-ol  
(D) 3-Methylbutan-3-ol


6. If a salt bridge is removed between the half cells the voltage:
  - (A) drops to zero
  - (B) does not change
  - (C) increase gradually
  - (D) increase rapidly
7. Radio active decay is an example of
  - (A) first order
  - (B) zero order
  - (C) second order
  - (D) none of these
8. Half life period of a first order reaction is
  - (A) directly propostional to the initial concentration of the reactant.
  - (B) Independent of initial concentration of reactions
  - (C) half of the sate constant
  - (D) None of these
9. The number of moles of  $\text{KMnO}_4$  that will be needed to react with one mole of  $\text{Fe}^{2+}$  ions in acidic medium is.
  - (A) 5
  - (B) 2
  - (C) 3
  - (D) 1
10. The hydrization of the complex  $[\text{Ni}(\text{CN})_4]^{2-}$  is :
  - (A)  $\text{Sp}^3$
  - (B)  $\text{dsp}^2$
  - (C)  $\text{dsp}^3$
  - (D)  $\text{d}^2\text{sp}^3$
11. Which of the follwoing molecule is chiral ?
  - (A) 2-Bromobutane
  - (B) I-Bromobutane
  - (C) 2-Bromopropane
  - (D) 2-Bromopropan-2-ol
12. Which of the following haloalkanes react with aqueous KOH most early?
  - (A) 1-Bromobutane
  - (B) I-Bromobutane
  - (C) 2-Bromo-2-methyle propane
  - (D) 2-chrorobutane
13. Which of the follwoing compound do not undergo aldol condensation?
  - (A)  $\text{C}_6\text{H}_5\text{CHO}$
  - (B)  $\text{CH}_3\text{CHO}$
  - (C)  $\text{CH}_3\text{COCH}_3$
  - (D)  $\text{CH}_3\text{CH}_2\text{CHO}$
14. Which of the following is most stable diazonium salt?
  - (A)  $\text{CH}_3\text{N}_2^+\text{X}^-$
  - (B)  $\text{C}_6\text{H}_5\text{N}_2^+\text{X}^-$
  - (C)  $\text{CH}_3\text{CH}_2\text{N}_2^+\text{X}^-$
  - (D)  $\text{C}_6\text{H}_5\text{CH}_2\text{N}_2^+\text{X}^-$

For questions number 15 to 18 two statements are given - one labelled Assertion (A) and the other labelled Reason (R). Select the correct answer to these questions from the codes (i), (ii), (iii) and (iv) as given below:

- (i) Both assertion (A) and reason (B) are correct statements , and reason (R) is the correct explanation of the assertion (A)
  - (ii) Both assertion (A) and reason (B) are correct statements , but reason (R) is not the correct explanation of the assertion (A)
  - (iii) Assertion (A) is correct , but reason (B) is incorrect statement
  - (iv) Assertion (A) is incorrect, but reason (R) is correct statement.
15. Assertion (A) : Boiling points of alkyl halides decrease in the order  $\text{R-I} > \text{R-Br} > \text{R-Cl} > \text{R-F}$   
 Reason(B): Van der Walls forces decrease with increase in the size of halogen atom

16. Assertion (A) : Low spin tetrahedral complexes are rarely observed  
Reason(B): The orbital splitting energies are not sufficiently large to forcing pairing
17. Assertion (A) : Benzene and hexane form idedsolution  
Reason(B): Both benzene and hexane are hydrocarbhone
18. Assertion (A) : Increase in temperature increases sole of reaction  
Reason(B): More reactant molecules have energy greater than threshold energy.

### SECTION-B

19. What happens when:  
(i) Propene is treated with HBr in presence of paroxide  
(ii) Benzene is treated with  $\text{CH}_3\text{Cl}$  in presence of  $\text{AlCl}_3$
20. Complete the following reactions:  
(i)   $\text{OH} + \text{SOCl}_2 \longrightarrow$   
(ii)  $\text{CH}_3\text{-Br} + \text{NaI} \xrightarrow{\text{acetone}}$
21. (a) Aniline does not undergo friedal crafts reaction.  
(b) Amino group in amiline is o-and p-directing in aromatic all electrophilic substitution reaction but aniline on nitration give a substantial amount of m-nitro aniline.  
Arrange the following:  
(a) In decreasing order of  $\text{pK}_b$  value  
 $\text{CH}_3\text{CH}_2\text{NH}_2$ ,  $\text{C}_6\text{H}_5\text{NH}_2$ ,  $(\text{C}_2\text{H}_5)_2\text{NH}$ ,  $\text{C}_6\text{H}_5\text{NHCH}_3$   
(b) Increasing order of boiling point:  
 $\text{C}_2\text{H}_5\text{OH}$ ,  $(\text{CH}_3)_2\text{NH}$ ,  $\text{C}_2\text{H}_5\text{NH}_2$
22. Write 'A' and 'B' in the given reactions sequences  
(i)  $\text{C}_6\text{H}_5\text{N}_2\text{Cl} \xrightarrow{\text{CuCN}} \text{A} \xrightarrow{\text{H}_3\text{O}^+} \text{B}$   
Complete hydrolysis  
(ii)  $\text{CH}_3\text{CH}_2\text{Br} \xrightarrow{\text{KCN}} \text{A} \xrightarrow{\text{LiAlH}_4} \text{B}$
23. Write the products of glucose with  
(i)  $\text{Br}_2\text{-H}_2\text{O}$  (ii)  $\text{HI}$
24. For a 5% solution of the (Molar mass =  $60 \text{ g mol}^{-1}$ ) calculate the osmotic pressure at  $300 \text{ K}$  ( $R=0.0821 \text{ L K}^{-1} \text{ mol}^{-1}$ ).

25. Give the formula of the following components:-

- (a) Potassium tetrahydroxozincate(II)
- (b) Tris ethylenediaminecobalt(III) ion

### SECTION C

26. (a) Determine the unit of rate constant for first order reaction  
 (b) Show that time required for the completion of 99% the first order reaction is twice the 90% of completion.

### OR

- (a) What are pseudo first order reaction ? Give example
- (b) Rate constant  $k$  reaction varies with temperature 'T' according to the equation:

$$\log K = \log A - \frac{E_a}{2.303RT}$$

When a graph is plotted for  $\log K$  vs  $1/T$ , A straight line with a slope of 4250K is obtained. Calculate  $E_a$  for the reaction.

27. Write one chemical reaction to illustrate the following:
- (a) Williamson synthesis
  - (b) Kolbe's Reaction
  - (c) coupling Reaction
28. (a) Give the IUPAC name and electronic configuration of central metal atom in terms of  $t_{2g}$  and  $e_g$  of  $K_4[Mn(CN)_6]$ .  
 (b) What is meant by 'Chelate effect' ? Give an example.
29. An antifreeze solution is prepared by dissolving 31 g of ethylene glycol (Molar mass =  $62 \text{ g mol}^{-1}$ ) in 600 g of water. Calculate the freezing point of the solution. ( $K_f$  for water =  $1.86 \text{ K kg mol}^{-1}$ )
30. Account for the following:
- (i) Copper (I) compounds are colourless whereas Cu(II) compounds are coloured.
  - (ii) Zn, Cd and Hg are not considered as d-block elements.
  - (iii) d-block elements exhibit variable oxidation states.

## SECTION D

31. Read the given passage and answer the questions follow. 1x5=5

The d-block of the periodic table contains the elements of the group 3-12 and are known as transition elements. In general, the electronic on configuration of these elements is  $(n-1) d^{1-10} ns^{1-2}$ . The d-orbitals of the penultimate energy level in their atoms receive electrons giving rise to the three rows of the transition metals i.e., 3d, 4d and 5d series. However, Zn, Cd and Hg are not regarded as transition elements. Transition elements exhibit certain characteristic properties like variable oxidation stats, complex formation, formation of coloured ions and alloys, catalytic activity, etc. Transition metals are hard (except Zn, Cd and Hg) and have a high melting point.

1. Why are Zn, Cd and Hg non-transition elements?
2. Which transition metal of 3d series does not show variable oxidation states?
3. Why do transition metals and their compounds show catalytic activity?
4. Why is  $Cu^{2+}$  ion coloured while  $Zn^{2+}$  ion is colourless in aqueous solutions?

32. **Read the given passage and answer the questions number 1 to 6 that follow :-**

The substitution reaction of alkyl halide mainly occurs by  $S_N1$  or  $S_N2$  mechanism. Whatever mechanism alkyl halides follow for the ubstitution reaction to occur, the polarity of the carbon halogen bond is responsible for these substitution reactions. The rate of  $SN^1$  reactions are governed by the stability of carbocation whereas for  $SN^2$  reactions steric factor is the deciding factor, If the starting material is a chiral compound, we may end up with an inverted product or racemic mixture depending upon the type of mechanism followed by alkyl halide. Cleavage of ethers with HI is also governed by steric factor and stability of carbocation, which indicates that in organic chemistry, these two major factors help us in deciding the kind of product formed.

1. Predict the stereochemistry of the product formed if an optically active alkyl halide undergoes substitution reaction by  $SN^1$  mechanism.


2. Name the instrument used for measuring the angle by which the plane polarised light is rotated.
3. Predict the major product formed when 2-Bromopentane reacts with alcoholic KOH.
4. Give one use of  $\text{CHI}_3$ .

**OR**

Out of  $\text{CH}_3\text{-Cl}$  and  $\text{CH}_3\text{-I}$ , which is more reactive in  $\text{S}_\text{N}1$  reaction?

### SECTION E

33. (a) The standard Gibbs energy ( $\Delta_r G^\circ$ ) for the following cell reaction is  $300 \text{ kJ mol}^{-1}$   
 $\text{Zn(s)} + 2\text{Ag}^+(\text{aq}) \rightarrow \text{Zn}^{2+}(\text{aq}) + 2\text{Ag(s)}$   
 Calculate  $E^\circ_{\text{cell}}$  for the reaction. (Given:  $1 \text{ F} = 96500 \text{ mol}^{-1}$ )
- (b) Calculate  $\lambda_m^\circ$  for  $\text{MgCl}_2$  if  $\lambda^\circ$  values for  $\text{Mg}^{2+}$  ion and  $\text{Cl}^-$  ion are  $106 \text{ S cm}^2 \text{ mol}^{-1}$  and  $76.3 \text{ S cm}^2 \text{ mol}^{-1}$  respectively.
- (c) In the plot of molar conductivity ( $\Lambda_m$ ) Vs/ square root of concentration ( $C$ ), following curves are obtained for two electrolytes X and Y :



Answer the following:

- (i) Predict the nature of electrolyte X and Y
- (ii) What happens on extrapolation of  $\Lambda_m$  to concentration approaching zero for electrolytes X and Y ?


**OR**

- (a) When a steady current of 2A was passed through two electrolytic cells A and B containing electrolytes  $\text{ZnSO}_4$  and  $\text{CuSO}_4$  connected in series, 2g of Cu were deposited at the cathode of cell B. How long did the current flow ?  
 What mass of Zn was deposited at cathode of cell A ?  
 [Atomic mass:  $\text{Cu} = 63.5 \text{ g mol}^{-1}$ ,  $\text{Zn} = 65 \text{ g mol}^{-1}$ ;  $\text{F} = 96500 \text{ C mol}^{-1}$ ]
- (b) Explain the effect of dilution on the molar conduction of electrolytic solution.
34. (a) Describe the preparation of potassium dichromate from iron chromite ore.
- (b) How does acidified  $\text{KMnO}_4$  solution react with
  - (i)  $\text{Fe}^{2+}$
  - (ii)  $\text{I}^-$
  - (iii) Oxalic acid,
 Write the ionic equations involved for the reaction.

35. (a) An organic compound 'A' having molecular formula  $C_5H_{10}O$  gives negative tollen's test, forms n-pentane on Clemmensen reduction but doesn't give iodoform test. Identify 'A' and give all the reactions involved.
- (b) Carrying out the following conversions.
- Propanoic acid to 2-Bromopropanoic acid
  - Benzoyl chloride to benzaldehyde
  - How will you distinguish between benzaldehyde and acetaldehyde?

**OR**

- (a) Complete the following sequence of reactions?



- Identify (A) to (D)
- Give the IUPAC name of (A)
- How can you distinguish between:
  - Ethanol and Propanone
  - Benzoic acid and Phenol?