

Directorate of Education, GNCT of Delhi
Practice Paper (Session 2025 – 26)
Class XII
Mathematics (CODE: 041)

Time Allowed: 3 Hours

Maximum Marks: 80

General Instructions:

1. This question paper contains **FIVE** sections – A, B, C, D & E. Each part is compulsory.
However, there are internal choices in some questions.
2. Section A has 18 MCQ's and 02 Assertion-Reason based questions of 1 mark each.
3. Section B has 5 Very Short Answer (VSA)-type questions of 2 marks each.
4. Section C has 6 Short Answer (SA)-type questions of 3 marks each.
5. Section D has 4 Long Answer (LA)-type questions of 5 marks each.
6. Section E has 3 source based/case based/passage based/integrated units of assessment (4 marks each) with sub parts.

SECTION – A (Multiple Choice Questions) Each question carries 1 mark Each MCQ has four options with only one correct option, choose the correct option.		
1.	The domain of $f(x) = \cos^{-1}(2x)$ is : (a) $[-1,1]$ (b) $[-2,2]$ (c) $[\frac{-1}{2}, \frac{1}{2}]$ (d) $[\frac{-1}{4}, \frac{1}{4}]$	1
2.	Let $A = \{9, -17\}$, then the number of reflexive relations defined on A is : (a) 2 (b) 4 (c) 8 (d) 16	1
3.	If A is a square matrix of order 3 & $ 2A = 32$, then the value of $ \text{adj}A $ is (a) 16 (b) 36 (c) 64 (d) 216	1
4.	For which value of x, are the determinants $\begin{vmatrix} x & 3 \\ 5 & x \end{vmatrix}$ & $\begin{vmatrix} -5 & 1 \\ 0 & -2 \end{vmatrix}$ equal? (a) 0 (b) ± 5 (c) -5 only (d) 5 only	1
5.	If A is a row matrix of order $a \times b$, B is a Column Matrix of order $b \times c$ and C is a square matrix of order $3 \times b$, then $(2a + 3b + c) =$ (a) 5 (b) 7 (c) 10 (d) 12	1


6.	If A is a square matrix of order 3 with $A \neq 1$, then A.(adj A) is <u>Not</u> a/an (a) Diagonal Matrix (b) Scalar Matrix (c) Identity Matrix (d) Symmetric Matrix	1
7.	The value of the cofactor of the element of second row and third column in the matrix $\begin{pmatrix} 5 & 3 & 2 \\ 2 & -1 & 0 \\ 1 & 2 & 3 \end{pmatrix}$ is (a) -5 (b) 5 (c) -7 (d) 7	1
8.	If $x = \sin t$ and $y = \cos t$, then $\frac{d^2y}{dx^2}$ is : (a) $\sec^3 t$ (b) $-\sec^3 t$ (c) $\operatorname{cosec}^3 t$ (d) $-\operatorname{cosec}^3 t$	1
9.	If $(\cos x)^y = (\cos y)^x$, then $\frac{dy}{dx} =$ (a) $\frac{y \tan x + \log(\cos y)}{x \tan y - \log(\cos x)}$ (b) $\frac{y \tan x - \log(\cos y)}{x \tan y - \log(\cos x)}$ (c) $\frac{y \tan x + \log(\cos y)}{x \tan y + \log(\cos x)}$ (d) $\frac{x \tan y + \log(\cos x)}{y \tan x + \log(\cos y)}$	1
10.	Derivative of $\tan^{-1}\left(\frac{x}{\sqrt{1-x^2}}\right)$ with respect to $\sin^{-1}\left(2x\sqrt{1-x^2}\right)$ is (a) 0 (b) 0.5 (c) 1 (d) 2	1
11.	If the rate of change of volume of a sphere is twice the rate of change of its radius, then the surface area of the sphere is : (a) $\frac{1}{2}$ sq.unit (b) 1 sq. unit (c) 2 sq. unit (d) 4 sq. unit	1
12.	The area (in Sq. units) bounded by the parabola $y^2 = x$, y - axis & the line $y = 3$ is : (a) $\frac{1}{3}$ (b) 1 (c) 3 (d) 9	1

13.	The value of $4 \times \int_0^{\frac{\pi}{2}} \left(\frac{\sin^{0.5} x}{\sin^{0.5} x + \cos^{0.5} x} \right) dx$ is (a) $\frac{\pi}{4}$ (b) $\frac{\pi}{2}$ (c) π (d) 2π	1
14.	The general solution of the differential equation $\frac{dx}{dy} = \frac{x}{y}$ is (a) $xy = c$ (b) $x + y = c$ (c) $x - y = c$ (d) $\frac{x}{y} = c$	1
15.	The integrating factor for solving the differential equation $x \frac{dy}{dx} = x + y$ is (a) e^x (b) e^{-x} (c) $\log x$ (d) $\frac{1}{x}$	1
16.	In an LPP, corner points of the feasible region determined by the system of linear constraints are (1, 1), (4, 0) and (0, 4). If $Z = px + qy$, where $p, q > 0$ is to be minimized, the condition on p and q, so that the minimum of Z occurs at (4, 0) and (1, 1), will be : (a) $0.5p = q$ (b) $p = q$ (c) $2p = q$ (d) $3p = q$	1
17.	If the feasible region of a linear programming problem with objective function $Z = ax + by$, is bounded, then which of the following is correct ? (a) It will only have a maximum value. (b) It will only have a minimum value. (c) It will have both maximum & minimum value. (d) It will have neither maximum nor minimum value.	1

18.	If $\vec{a} = 1, \vec{b} = 2$ & $\vec{a} \cdot \vec{b} = 1$, then $\vec{a} + 2\vec{b} =$ (a) $2\sqrt{2}$ (b) 3 (c) $\sqrt{10}$ (d) $\sqrt{11}$	1
19.	ASSERTION-REASON BASED QUESTIONS (Q.19 & Q.20) In the following questions, a statement of assertion (A) is followed by a statement of Reason (R). Choose the correct answer out of the following choices. (a) Both A and R are true and R is the correct explanation of A. (b) Both A and R are true but R is not the correct explanation of A. (c) A is true but R is false. (d) A is false but R is true. <i>ASSERTION(A)</i>: If $\vec{a} \times \vec{b} ^2 + \vec{a} \cdot \vec{b} ^2 = 25$ & $\vec{a} = 5$ then $\vec{b}$ is a unit vector. <i>REASONING(R)</i>: $\sin^2 x + \cos^2 x = 1, \vec{a} \times \vec{b} = \vec{a} \vec{b} \sin \theta$ and $\vec{a} \cdot \vec{b} = \vec{a} \vec{b} \cos \theta$.	1
20.	<i>ASSERTION(A)</i>: $P\left(\frac{A}{B}\right) + P\left(\frac{A'}{B}\right) = 1$. <i>REASONING(R)</i>: A & A' are complementary events.	1
SECTION B This section comprises of very short answer type-questions (VSA) of 2 marks each		
21.	A drone is flying in 3D space. At any instant, its position vector is given by (4, 9, 2). It moves in a straight path such that every change in its coordinates satisfies the relation: $\frac{x-4}{3} = \frac{9-y}{2} = \frac{3z-6}{12}$ Determine the direction ratios & direction cosines of the path along which drone is moving.	2
22.	<i>Evaluate</i>: $I = \int \frac{dx}{(x^2 + 1)(x^2 + 4)}$.	2

23.	A company is designing a robotic arm that moves within a limited angular range. The control angle is defined by the function $f(x) = \cos^{-1}\left(\frac{3-2x}{7}\right).$ Determine the range of possible values of x for which the control angle $f(x)$ is defined. OR The greatest integer function $f(x) = [x]$ converts every real number to the nearest smaller integer. (a) Explain with examples, why this function cannot be one-one. (b) Can this function ever be onto if the co-domain is all real number $\mathbb{R}$? Justify logically.	2
24.	If $y = (x + \sqrt{x^2 + 1})^2$, then prove that $(x^2 - 1)\left(\frac{dy}{dx}\right)^2 = 4y^2$. OR Show that the function $f(x) = \frac{16 \sin x}{4 + \cos x} - x$ is strictly decreasing in $(\frac{\pi}{2}, \pi)$.	2
25.	Let $\vec{a} = 2\hat{i} + 3\hat{j} + 4\hat{k}$, $\vec{b} = 4\hat{i} - 3\hat{j} + 9\hat{k}$ & $\vec{c} = 3\hat{i} - 2\hat{j} + 6\hat{k}$. Find a vector $\vec{d}$ which is perpendicular to both $\vec{a}$ & $\vec{b}$ and $\vec{c} \cdot \vec{d} = 39$.	2
SECTION C (This section comprises of short answer type questions (SA) of 3 marks each)		
26.	If A & B are two independent events, then show that A' & B are also Independent Events. Hence, find the value of $P(A' \cap B)$ such that $P(A \cap B) = 0.06$, $P(A) = 0.2$ & $P(B) = 0.3$	3

27.	Solve the following Differential Equation when $y(0) = 0$: $\frac{dy}{dx} + (\sec^2 x) y = \tan x.(\sec^2 x)$ <i>OR</i> Find the general solution of the differential equation $x dy - y dx = \sqrt{x^2 + y^2} dx$	3
28.	The median of an equilateral triangle is increasing at the rate of $2\sqrt{3} \text{ cm / sec}$. Find the rate at which its side is increasing. <i>OR</i> Prove that: $\frac{d}{dx} \left(\frac{x}{2} \sqrt{a^2 - x^2} + \frac{a^2}{2} \sin^{-1} \left(\frac{x}{a} \right) \right) = \sqrt{a^2 - x^2}.$	3
29.	Find : $I = \int \frac{(x + \sin 2x) dx}{1 + \cos 2x}$ <i>OR</i> Evaluate : $I = \int_{-1}^1 x \sin \pi x dx.$	3
30.	Given that $\hat{a}, \hat{b}$ & $\hat{c}$ are unit vectors such that $\hat{a} \cdot \hat{b} = \hat{a} \cdot \hat{c} = 0$ and angle between $\hat{b}$ & $\hat{c}$ is $\frac{\pi}{6}$. (i) Show that $\hat{b} \times \hat{c} = \frac{1}{2}.$ (ii) Use this to infer what constant must multiply $(\hat{b} \times \hat{c})$ to make it a unit vector. (iii) Hence justify why $\hat{a} = \pm 2(\hat{b} \times \hat{c}).$	3

31.	Consider the linear programming problem, where the objective function $Z = 2x + 4y$ needs to be minimized subject to constraints $2x + y \geq 1000$ $x + 2y \geq 800$ $x \geq 0, y \geq 0$	3
SECTION D (This section comprises of long answer-type questions (LA) of 5 marks each)		
32.	If Matrix $A = \begin{pmatrix} 3 & 2 & 1 \\ 4 & 1 & 3 \\ 1 & 1 & 1 \end{pmatrix}$, find A^{-1} and hence solve the following system of linear equations: $3x + 2y + z = 2000$ $4x + y + 3z = 2500$ $x + y + z = 900$ OR Observe the following Graph and Using the concept of Matrices find the values of constants a, b & c. Hence write the expression for y in terms of x. 	5

33.	Using integration find the area bounded by the curve $x^2 + y^2 = 4$. OR A planet orbits its star in an elliptical path modeled by the equation : $9x^2 + 4y^2 = 36$ Astronomers want to calculate the total area enclosed by this orbit to understand the region in which the planet moves. Find the area of the elliptical orbit using integration.	5
34.	Find the intervals where $f(x) = \sin^4 x + \cos^4 x, x \in [0, \frac{\pi}{2}]$ is (a) Strictly increasing (b) Strictly Decreasing	5
35.	Two airplanes are flying along different straight-line trajectories, represented by $L_1: \frac{x-1}{1} = \frac{y-2}{-1} = \frac{z-1}{1} \text{ \& } L_2: \frac{x-2}{2} = \frac{y+1}{1} = \frac{z+1}{2}$ To avoid mid-air collisions, air traffic controllers need to determine the shortest distance between the two flight paths. Calculate this shortest distance to check if the aircraft are at a safe separation.	5



SECTION - E

This section comprises of 3 case-study/passage-based questions of 4 marks each with two sub-parts.

First two case study questions have three sub parts (A), (B) & (C) of marks 1, 1, 2 respectively.

The third case study question has two sub-parts of 2 marks each.

36. A garden designer is planning to create a flower bed in the shape of an isosceles trapezium. The three sides other than the base (two equal non – parallel sides and the smaller base) are all 10 cm each.
- To make the design look more appealing, she wants to maximize the area of the trapezium while maintaining these side lengths.
- On the basis of above information answer the following questions:



- (a) Let the length of the base of smaller triangular region is x cm.
Find the distance between the two parallel sides.
- (b) Express the area of the trapezium in terms of x .
- (c) (i) Using First derivative Approach, find the value of x for which the area is maximum and calculate this maximum area.
- OR
- (ii) Using Second derivative Approach, find the value of x for which the area is maximum and calculate this maximum area.

1

1

2

2

37. A jeweller is testing a random sampling method using three identical boxes — Box I, Box II, and Box III. Each box contains two coins, as shown below :

BOX	Gold coins	Silver coins
Box I	2	0
Box II	1	1
Box III	0	2

A person chooses one box at random and draws one coin without looking. The drawn coin turns out to be Gold.

On the basis of above information answer the following questions:



- (a) List all the possible outcomes (sample space) when a person randomly chooses a box and draws one coin. 1
- (b) Find the probability that the chosen coin is a gold coin. 1
- (c) (i) Using the result of (b), find the probability that the other coin in the box is also gold, given that a gold coin has been drawn. 2

OR

- (ii) Using the result of (b), find the probability that the other coin in the box is silver, given that a gold coin has been drawn. 2

38. A social networking app is testing a new feature that connects users based on their interactions. The app records the relation R on the set $A = \{\text{Aditi, Bhavna, Charu, Divya}\}$, where $a R b$ means “a follows b on the app.”


The following data was recorded :

User	Follows
Aditi	Bhavna & Charu
Bhavna	Aditi & Bhavna
Charu	Charu & Aditi
Divya	Charu & Divya

Social Network Case Study: Follower Relationships

Users: Aditi, Bhavna, Charu, Divya

$R: a \text{ follows } b$



On the basis of above information answer the following questions:

- (a) Represent the relation R as a set of ordered pairs and determine whether it is reflexive and symmetric.

Give reasons for your answer.

- (b) Check whether the relation R is transitive.

Hence, state whether R is an equivalence relation or not.

2

2